




3 2044 106 366 586

Library
Arnold Arboretum



of

Harvard University

Rebound February 1966



Digitized by the Internet Archive
in 2017 with funding from
BHL-SIL-FEDLINK

GULIELMI HUDSONI

Regiæ Societatis Socii et Pharmacopæi Londinensis.

4300
11

Flora Anglica,

EXHIBENS

PLANTAS

PER

Regnum Angliæ sponte crescentes,

DISTRIBUTAS

Secundum Systema Sexuale :

CUM

DIFFERENTIIS SPECIERUM,

SYNONYMIS AUTORUM,

NOMINIBUS INCOLARUM,

SOLO LOCORUM,

TEMPORE FLORENDI,

OFFICINALIBUS PHARMACOPÆORUM.

LONDINI:

Impensis AUCTORIS : Prostant venales apud J.
NOURSE in the Strand, et C. MORAN in Co-
vent-Garden.

M.DCC.LXII.

OFFICE OF THE SECRETARY OF THE ARMY

WASHINGTON, D. C.

Flora Anglica

NORTH AMERICA

PLANT

Region

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

...

NOBILISSIMO VIRO

HUGONI COMITI

NORTHUMBRIÆ,

ORDINIS PERISCELIDIS EQUI, &c.

ARTIUM TUM UTILIUM,

TUM ELEGANTIORUM,

JUDICI ET PATRONO,

HOC OPUS HUMILLIME DEDICAT

Gul. Hudson.



PRÆFATIO.



AUCIS lectorem præfari nec-
cessarium duxi, tum ut eum
admonerem quid in hac Flo-
ra expectare debeat, tum ut
notas quasdam compendii gra-
tiâ usurpatas explicarem.

Fundamentum hujus operis quoad *Catalo-
gum* plantarum Raii synopsis agnoscitur, et
agnosci debet; quantulum enim intactum
reliquit vir ille diligentissimus, accuratissi-
mus, et addere licet, sagacissimus, confiten-
tibus tum popularibus suis, tum exteris; qui
floram ejus inter optimas et perfectissimas
uno ore numerant? si nullum igitur post ejus
tempora augmentum accepisset ipsa scientia
botanica nihil pene pro plantis anglicanis dig-
noscendis ulterius desideraretur, quam nova

synopseos editio, cujus exemplaria hodie apud bibliopolas omnino non inveniuntur. At exortum est his diebus novum fidus, quod orbi botanico lucem affudit ne in somnis quidem antea visam; magnus scilicet ille Linnæus, qui partes plantarum minutissimas, vel prius neglectas, vel non omnino observatas in conspectum producendo, plene illustrando, et rite nominando fundamentum jecit novæ methodo simplici, compendiosæ, certæ; quam et constantiâ plane admirandâ toti regno vegetabili adhuc cognito feliciter applicuit.

Hunc igitur virum egregium quoad *methodum* per omnia, quatenus licuit, secutus sum.

Essentiales characteres ex Systemate naturæ edit. ultim. adduxi; quando autem novum genus occurrebat, characterem naturalem ad ejus exemplar, prout in *Genera plantarum* exhibetur adjunxi.

Quoad Synonyma, ut vocantur, sive Characteres Specificos, primo loco eos adhibui quos idem author in *Species plantarum* dedit; quotiescunque certum fuit eum plantam descripsisse; aliter novum characterem, ejus vestigiis insistens construxi; postea synonyma adjeci ex Bauhini pinace, Raio, Gerardo, Parkinsono et variis authoribus hodiernis deprompta.

Nomina plerumque Anglica ex Raio, Gerardo, &c. apposui; quædam mutavi exemplo
et

et hortatu Domini Stillingfleet motus, ut generica obtinerem, quod optandum esset per omnia, ut confusio quæ hodie regnat finis aliquando imponatur; quando autem nulla occurrebant meo jure, ut opinor, imposui.

Quoad loca natalia plantarum satis duxi indicasse solum, quo maxime vulgares sponte proveniunt; locum vero rariorum semper notavi ex Raii Synopsi ultima Editione et ex observationibus sive propriis, sive amicorum.

Plantarum naturam quoad durationem indicavi Signis sequentibus; viz. ☉ annuam, ♂ biennem, ♀ perennem innuit.

Hoc signum ♀ fruticem vel arborem designat.

Numeri I, II, III, &c. tempus florendi indicant, viz. Januarium, Februarium, Martium, &c.

Plantas cultas non autem indigenas omnino omisi, ut Avenam sativam, Triticum sativum, Solanum tuberosum, &c.

Plantæ quæ sine culturâ non raro inveniuntur extra hortorum pomæria, sed quæ ab origine propagationem eorum rejectamentis debeat asterisco * distinguuntur.

Hanc occasionem lubens amplector gratias agendi amicis, qui mihi plantas pro eorum benevolentia suppeditarunt inter quos nominandi sunt Domini Alchorne, Bolton, Collinson, Lyons, Miller, Watson, M. D. Wilmer, M. D. præcipue vero D. Stillingfleet, qui et hoc et aliis amicitiae officiis privatim et publice functus est.

Cæterum quamvis nihil sciens prætermis-
si quod confere potuit elucidando rem subjec-
tam, non labori peperci, nullam occasionem
neglexi libros consulendi, herbaria sicca e-
volvendi et plantas vivas examinandi et des-
secandi, ut opus omnibus numeris absolu-
tum prodiret; conscius sum tamen me nul-
lo modo scopum attigisse, præsertim quoad
classẽ cryptogamium, quæ ad hæc usque
tempora quantum scio, obscura manet, et
mihi plurimum molestiæ creavit. In uno
istius classis ordine, viz. Fungorum me mul-
tum deficere agnosco; nec sine aliis Dilleni-
is, aliis Micheliis extricare se quisquam po-
terit, cum hunc ordinem experientia edoctus
affirmare possum, ut Linnæi verbis utar, eti-
amnum Chaos esse, nescientibus Botanicis in
his, quid sit Species, quid Varietas.





Nomina Auctorum Explicata.

ACT. parif. Memoires de l'Academie Royale des Sciences. Paris. 4to.

Act. Lond. Acta Philosophica Regiæ Societatis Londinensis. Lond. 4to.

Act. upf. Acta Regiæ Societatis Upsalienfis. Holmiæ. 4to.

Amm. St. rar. ruth. Joannis Ammani Stirpium rariorum in Imperio Ruthenico, &c. Petrop. 1739. 4to.

Barr. Icon. Jacobi Barrelieri Icones Plantarum per Galliam, Hispaniam et Italiam observatarum. Paris. 1714. fol.

Blackf. spec. bot. J. Blackstone Specimen Botanicum. Lond. 1746. 8vo.

Blackf. cat. pl. ejusdem Catalogus Plantarum circa Harefield. Lond. 1737.

Baub. pin. et pr. Caspari Bauhini Pinax et Prodrromus Theatri Botanici. Basil. 1671. 4to.

Baub.

NOMINA AUCTORUM EXPLICATA.

Bauh. th. Ejusdem Theatrum Botanicum. *ibid.*
1658.

Bauh. hist. Joannis Bauhini Historia Plantarum universalis. 3. Ebrod. 1650. fol.

Bocc. pl. rar. Pauli Boccone Plantarum rariorum Siciliae Icones et Descriptiones. Oxon. 1674. 4to.

Bocc. mus. Ejusdem musæum Plantarum Rariorum. Venet. 1697. 4to.

Boerb. ind. Hermannii Boerhaave Index alter Plantarum. Lugdb. 1720. 4to.

Breyn. cent. Jacobi Breynii Centuria Plantarum Gedani. 1678. fol.

Breyn. cent. pr. Ejusdem Prodomus Fasciculus Plantarum. *ibid.* 1689. 4to.

Buddl. hort. succ. Adami Buddleii Hortus Siccus Plantarum Angliæ.

Buxb. en. pl. hal. Joannis Christiani Buxbaum Enumeratio Plantarum in Agro Hallensi. Halæ Magdeb. 1721. 8vo.

Buxb. cent. Ejusdem Plantarum minus cognitarum Centuriæ quatuor. I. et II. 1728. III. 1729. IV. 1733. 4to. Petrop.

Cæsalp. Andreas Cæsalpinus de Plantis. Florent. 1583. 4to.

Cat. giff. Joannis Jacobi Dillenii Catalogus Plantarum circa giffam. Giffæ. 1719. 8vo.

Cat

NOMINA AUCTORUM EXPLICATA.

Cat. nott. Car. Deering Catalogus Plantarum circa Nottinghamiam. Nottingh. 1738. 8vo.

Cat. pl. angl. Joannis Raii Catalogus Plantarum Angliæ. Lond. 1688. 12mo.

Cat. cant. Ejusdem Catalogus Plantarum circa Cantabrigiam. Cant. 1633. 12mo.

Clus. Caroli Clusii Rariorum Stirpium per Hispaniam observatarum. Antw. 1601. fol.

Col. ecph. Fabii Columnæ Ecphrasis I. et II. minus cognitarum rariorumque Stirpium. Romæ. 1616. 4to.

Comm. cat. Joannis Commelini Catalogus Plantarum Hollandiæ. Amstel. 1685. 12mo.

Dalib. parif. M. Dalibardi Floræ Parisinæ prodromus. Paris. 1750. 8vo.

Dill. musc. Joannis Jacobi Dillenii Historia Muscorum. Oxon. 1741. 4to.

Dalech. lugd. Jacobi Dalechampii Historia generalis Plantarum. Lugd. 1584. fol.

Dod. pempt. Remberti Dodonæi Stirpium Historiæ Pemptades Sex. Antw. 1616.

Fl. jen. Henrici Bernhardi Rupprii Flora Jenensis. Francofurt. 1726. 8vo.

Fl. sibir. Joannis Georg. Gmelini Flora Siberica. 2. Petrop. 1750. 4to.

Fl. suec. Caroli Linnæi Flora Suecica. Ed. I. Lugdb. 1745. II. Stockh. 1755. 8vo.

NOMINA AUCTORUM EXPLICATA.

Fl. lapp. Ejusdem Flora Lapponica. Amst. 1737.
8vo.

Fl. pruss. Joannis Loefelii Flora Prussica. Regimont.
1703. 4to.

Fuchf. Leonardi Fuchsii Historia Plantarum. Basil.
1542. fol.

Ger. Em. Joannis Gerardi Historia Plantarum a Thoma Johnson Emaculata. Lond. 1633. et 1636. fol.

Gron. virg. Joannis Fred. Gronovii Flora Virginica. 2.
Leid. 1739. 8vo.

Gled. fung. Joan. Gottlieb. Gleditchii Methodus Fungorum. Berol. 1753. 8vo.

Guett. stamp. D. Guettardus Observations sur les Plantes. 2. Paris. 1747. 12mo.

Hist. ox. Roberti Morisoni et Jacobi Bobarti Historia Plantarum Oxoniensis. 3. Oxon. 1680.

Hall. helv. Alberti Halleri Enumeratio Stirpium Helveticarum. Gotting. 1742. fol.

Hort. reg. monspel. Petri Magnol. Hortus Regius Montpelienfis. Monsp. 1697. 8vo.

Johnsf. Thomæ Johnsoni Itinera Botanica. Lond. 1632.
1634. 8vo.

Lin. gen. pl. Caroli Linnæi Genera Plantarum. Lugdb.
1752. 8vo.

Lin. phil. bot. Ejusdem Philosophia Botanica. Holmiæ.
1751. 8vo.

Lin.

NOMINA AUCTORUM EXPLICATA.

- Lin. hort. upf.* Ejusdem Hortus Upsaliensis. Holmiæ. 1748. 8vo.
- Lin. hort. cliff.* Ejusd. Hortus Cliffortianus. Amstel. 1737. fol.
- Lin. virid. cliff.* Ejusd. Viridiarum Cliffortianum. Amstel. 1737. 8vo.
- Lin. mat. med.* Ejusd. Materia Medica. Holmiæ. 1749. 8vo.
- Lin. it. oeland.* Ejusd. Iter Oelandicum. Holmiæ. 1745. 8vo.
- Lin. it. W-goth.* Ejusd. Iter Westrogothicum. Holmiæ. 1745. 8vo.
- Lin. it. scan.* Ejusd. Iter Scanicum. Holmiæ. 1751. 8vo.
- Lin. amoen. acad.* Ejusd. Amoenitates Academicæ. I. Lipsiæ. 1749. II. Holmiæ. 1751. III. Amstel. 1756. IV. et V. Lugdb. 1760. 8vo.
- Lin. spec. pl.* Ejusd. Species Plantarum. Holmiæ. 1753.
- Lob. ic. Matthiæ Lobelii* Icones Plantarum. Antw. 1581. 4to.
- Lob. ob. et adv.* Ejusdem Observationes et Adversaria. Lond. 1605. fol.
- Magn. bot. monsp.* Petri Magnol Botanicum Monspeliense. Monspel. 1686. 8vo.
- Mart. cent.* Joannis Martyn Centuria Plantarum. Lond.
- Mart. tourn.* Ejusd. Tournefort's History of Plants. 2. Lond. 1732.
- Mentz. pug. ic.* Christiapi Mentzelii Index Plantarum et Pugillus. Berol. 1696. fol.

Merr.

NOMINA AUCTORUM EXPLICATA.

- Merr. pin.* Christophori Merret Pinax Rerum Naturalium Britannicarum. Lond. 1667. 8vo.
- Mich. nov. gen.* Petri Antonii Michelii Nova Plantarum Genera. Florent. 1729. 4to.
- Park.* Joannis Parkinsoni Theatrum Botanicum. Lond. 1640. fol.
- Park. par.* Ejusd. Paradisus Terrestris. Lond. 1656. fol.
- Phyt. Brit.* Phytologia Britannica. Lond. 1650.
- Pet. herb. br.* Jacobi Petiveri Herbarium Britannicum. Lond. fol.
- Pet. Gaz.* Ejusd. Gazophylaceum. 2 vols. fol. Lond. 1711.
- Pet. conc. gr.* Ejusd. Concordia Graminum. Lond. fol.
- Pet. mus.* Ejusd. Musæum. Lond. 1703. 8vo.
- Pluk. alm et phyt.* Leonardi Pluknetii Almagestum Botanicum et Phytographia. Lond. fol. diverso tempore edita.
- Pont.* Julii Pontederæ Compendium Tabularum Botanicarum. Patavii. 1720. 4to.
- R. Syn.* Joannis Raii Synopsis Methodica Stirpium Britannicarum. Ed. I. 1690. II. 1696. III. Lond. 1724. 8vo.
- R. hist.* Ejusd. Historia Plantarum. 3. Lond. 1686. et 1704. fol.
- R. meth.* Ejusd. Methodus Plantarum emendata et aucta. ibid. 1703. 8vo.
- Riv. irr. mon. ic.* Augusti Quirini Rivini Ordines Plantarum Flore Irregulari Monopetalo, Tetrapetalo et Pentapetalo. Lipsiæ 1691. et 1699. fol.
- Rox.

NOMINA AUCTORUM EXPLICATA.

Roy. lugdb. Adriani Van Royen Floræ Leidenfis Prodromus. Lugdb. 1740. 8vo.

Sauv. monsp. Fr. Sauvages Flora Monspeliensis. Hagæ. 1751. 8vo.

Sch. agr. Joannis Jacobi Scheuchzeri Agrostographia, five Graminum Juncorum Cyperorum Cyperoidum iisque affinium Historia, Tigurii. 1719. 4to.

Sibb. scot. il. Roberti Sibbaldi Scotia Illustrata. Edinb. 1684 fol.

Spig. Ifag. Andreæ Spigelii Ifagoge in Rem Herbariam. Lugdb. 1633.

Tab. Jacobi Theodori Tabernæmontani Icones Plantarum. Franc. 1690. 4to.

Trag. Hieronymi Tragi Stirpium Historia. Argentor. 1552. 4to.

Turn. Gulielmi Turneri Historia Plantarum. Lond. 1551.

Tourn. Josephi Pitton Tournefort Institutiones Rei Herbariæ. Paris. 1700. 4to.

Vaill. paris. Sebastiani Vaillant Botanicon Parisiense. Lugdb. et Amstel. 1727. fol.



ERRATA.

PA G. 5. lin. 30. pro calycibus lege calycibus. p. 19. l. 4. palustre. p. 22. l. 6. muticis. p. 29. l. 30. aristatæ. p. 32. l. 13. pratensis. p. 38. l. 26. nemoralis. p. 39. l. 6. paniculato. p. 44. l. 21. et in pratis. p. 47. l. 30. prato. p. 48. l. 7. salinas et antiquas. ib. cis. p. 52. l. 4. subovata. p. 60. l. 14. Græs. p. 69. l. 12. pinguibus. p. 75. l. 21. ne una. p. 78. l. 6. deciduus. p. 79. l. 12. Belladonna. p. 90. l. 1. petiolis. p. 95. l. 19. capitales. p. 96. l. 9. umbella. p. 97. l. 2. Involucra umbellis. p. 98. l. 5. radiatæ. ib. l. 6. suborbiculatæ. p. 104. l. 10. subhumidas. p. 105. l. 28. adde, Anglis; Hemlock Dropwott. p. 109. l. 1. cerefolium. p. 123. l. 13. flores. p. 136. l. 2. Eboracensis, Westmorlandicis, Cumberlandicis. p. 410. l. 26. terminali. p. 144. l. 2. filamenta. p. 145. l. 34. chamælea. p. 156. l. 3. coloratus. p. 161. l. 16. unifloro. p. 167. l. 9. polysperma. p. 168. l. 1. paniculato. ib. 40. Triplow. p. 169. l. 3. murum. ib. 26. cucullatis. p. 170. l. 33. minimum. p. 179. l. 20. Lythrum. p. 186. l. 6. prominentibus. p. 188. l. 4. beam. p. 189. l. 24. pomis. basi. p. 195. l. 37. præsertim ubi &c. p. 210. l. 2. unguis. p. 220. l. 16. Nep. p. 225. l. 26. floralibus. p. 226. l. 31. lateribus. p. 229. l. 24. bilabiatis. p. 242. l. 19. brevior. p. 250. l. 29. Iræo. p. 255. l. 13. codded. p. 256. l. 28. fugacissimis. p. 259. l. 22. Apulus. p. 268. l. 1. in littoribus. p. 287. l. 6. lupulinum. ib. 19. desfecteus. p. 309. l. 14. aristis. ib. 17. flosculosis. p. 318. l. 10. folii. p. 332. l. 17. dele ad. p. 342. l. 26. linearis. p. 344. l. 23. petiolatis. p. 354. l. 22. Betula. p. 361. l. 7. dele *. ib. 14. adde *. ib. 21. adde *. p. 370. l. 19. Germina. p. 372. l. 13. tria. p. 384. l. 16. Pteris. p. 399. l. 14. individuo. ib. 21. incubitu. p. 404. l. 39. aggeres. p. 406. l. 15. scoparium. p. 410. l. 10. adde short Bryum. p. 437. l. 2. foliolo. p. 440. l. 5. Anthera. p. 448. l. 26. ater. p. 490. l. 13. fluviatilis. p. 495. l. 32. truncis.

MONANDRIA MONOGYNIA.

Classis I.

MONANDRIA MONOGYNIA.

SALICORNIA *Lin. Gen. plant.* 10.

Calyx ventricosus, integer. *Petala* nulla, *Semen* unicum.

SALICORNIA articulis apice crassioribus obtusis. *europæa*.
Lin. mat. med. 8. *Sp. plant.* 3.

Salicornia geniculata annua. *Tourn.* 51. *R. Syn.* 136.

Kali geniculatum annum. *Bauh. pin.* 289. *Ger. Em.* 535. *Park.* 280.

β Kali geniculatum perenne fruticosius procumbens. *R. Syn.* 136.

γ Salicornia myosuroides procumbens, furculis longissimis. *R. Syn.* 137.

δ Salicornia ramosior procumbens, foliis brevibus purpurascens. *R. Syn.* 137.

ε Salicornia erecta, foliis brevibus cupressiformibus. *R. Syn.* 137.

Anglis, Marsh Sampire, Jointed Glasswort, or Saltwort.

Habitat in littoribus maritimis frequens. ☉ VIII. IX.

HIPPURIS *Lin. Gen. pl.* 11.

Calyx nullus, *Petala* nulla, *Stigma* simplex. *Semen* unicum.

HIPPURIS. *Fl. lapp.* 1. *Fl. suec.* 2. *Sp. Pl.* 4. *vulgaris*.

Limnopeuce. *Cord. hist.* 150. *R. Syn.* 136.

Equisetum palustre, brevioribus foliis, polyspermum.
Bauh. pin. 15.

Cauda equina fœmina. *Ger. Em.* 1114.

Equisetum palustre alterum brevioribus setis. *Park.* 1200.

MONANDRIA DIGYNIA.

Anglis Mare's-tail.

Habitat in fontibus, paludibus, solo profundo limoso,
in stagnis prope Hornsey. 24. V.

DIGYNIA.

CALLITRICHE *Lin. Gen. pl.* 13.

Calyx nullus. *Petala* duo. *Semen* unicum ovale, nudum.

verna.

1. CALLITRICHE foliis superioribus ovalibus, floribus androgynis. *Lin. Syst. Nat.* 844.

Stellaria. Cat. Giff. app. 119. t. 6. *R. Syn.* 289.

Stellaria aquatica. Bauh. pin. 141. *Park.* 1282.

Alfne palustris ferpyllifolia. Ger. Em. 614.

- β *Stellaria minor et repens. Cat. Giff.* 120. *R. Syn.* 289.

Anglis, vernal Starwort, Star-headed Water Chickweed.

Habitat in fossis et aquis quietis ubique. ☉. IV. --- VIII.

Autumnalis.

2. CALLITRICHE foliis omnibus linearibus apice bifidis, floribus hermaphroditis. *Lin. Syst. Nat.* 844.

Stellaria aquatica longifolia. Pet. H. br. t. 6. f. 4.

R. Syn. 290.

Lenticula aquatica angustifolia, folio in apice dissecto.

Fl. pr. 140. t. 38.

Anglis, Autumnal Starwort.

Habitat cum priore frequens. ☉. IX.

Classis

Classis II.

DIANDRIA
MONOGYNIA.* *Flores inferi, monopetali, regulares.*

LIGUSTRUM. 18.

*Corolla quadrifida. Bacca tetrasperma.*LIGUSTRUM. *Hort. cliff. 6. Fl. suec. 5. Sp. pl. vulgare.*
76. *J. Bauh. hist. I. 528. Ger. Em. 714. Park.*
1446. *R. Syn. 465.*Ligustrum germanicum. *Bauh. pin. 475.**Anglis, Privet, or Prim.**Habitat in sylvis et sepibus. h. V.*** *Flores inferi, monopetali, irregulares. Fructus capsulares.*

VERONICA. 25.

*Cor. quadrifida rotata : lacinia infima angustiore. Capsula bilocularis.** *Spicatæ.*I. VERONICA spica terminali, foliis oppositis crenatis obtusis; caule adscendente simplicissimo. *Sp. pl. 10. Fl. suec. 10.**Veronica spicata minor. Bauh. pin. 247. R. Syn. 279.**Veronica recta minima. Ger. Em. 527.**Veronica erecta angustifolia. Park. 550.**Anglis, Upright spiked male Speedwell, or Fluëllin.**Habitat in pratis et pascuis. In several Closes adjoining to Newmarket Heath, beyond Botolpham. R. Syn. in pascuis circa Horseheath Hall, in comitatu Cantabrigienfi. D. Miller. 4 VI.*

- hybrida.* 2. VERONICA spica terminali, foliis oppositis obtuse ferratis, caule erecto. *Sp. pl.* 11.
 Veronica spicata Cambrobritannica, bugulæ subhirsuto folio. *R. Syn.* 278. *t.* 11.
Anglis, Welsh Speedwell.
 Habitat ad latera montis cujusdam, Craig Wreidhin dicti in comitatu Montis Gomerici Walliæ, D. Lhwyd invenit 4. VII.
- officinalis.* 3. VERONICA spicis lateralibus pedunculatis, foliis oppositis, caule procumbente. *Sp. pl.* 11.
 Veronica mas supina et vulgatissima. *Bauh. pin.* 246.
R. Syn. 281.
 Veronica vera et major. *Ger. Em.* 626.
 Veronica mas vulgaris et supina. *Park.* 550.
Anglis, male Speedwell.
 Habitat in pascuis sterilibus et ericetis. 4. V.
Pharm. Veronicæ Herba.
 ** *Corymboso-racemosæ.*
- serpyllifolia.* 4. VERONICA racemo terminali subspicato, foliis ovatis glabris crenatis. *Sp. pl.* 12.
 Veronica pratensis serpyllifolia. *Bauh. pin.* 247.
 Veronica pratensis minor. *Park.* 551. *R. Syn.* 279.
 Veronica minor. *Ger. Em.* 627.
 Veronica fœmina quibusdam, aliis Betonica Pauli serpyllifolia. *Bauh. hist.* III. 366.
Anglis, Paul's Betony, little or smooth Speedwell.
 Habitat in pratis et pascuis passim. 4. V.
- Becabunga.* 5. VERONICA racemis lateralibus, foliis ovatis planis, caule repente. *Fl. suec.* 14. *Sp. pl.* 12.
 Veronica aquatica rotundifolia, Becabunga dicta minor. *R. Syn.* 280.
 Anagallis aquatica minor, folio subrotundo. *Bauh. pin.* 252.
 Anagallis aquatica vulgaris, sive Becabunga. *Park.* 1236.
 Anagallis, sive Becabunga. *Ger. Em.* 620.
Anglis, common Brooklime.
 Habitat in fossis et rivulis. 4. VI.
Pharm. Becabungæ Herba.

6. VERONICA racemis lateralibus, foliis lanceolatis *Anagalliferratis*, caule erecto. *Sp. pl.* 12. *Fl. suec.* 13. *lis.* Δ.
 Veronica aquatica longifolia media. *R. Syn.* 280.
Anagallis aquatica minor folio oblongo. Bauh. pin.
 252.
Anagallis aquatica folio oblongo crenato. Park. 1237.
Anagallis aquatica minor. Ger. Em. 620.
Anglis, long-leaved Water-speedwell.
Habitat in fossis et palustribus. 2. VII.
7. VERONICA racemis lateralibus, pedicellis pendulis, *scutellata*.
 foliis linearibus integerrimis. *Sp. pl.* 12. *Fl. suec.* 17.
 Veronica aquatica angustifolia minor. *R. Syn.* 280.
Anagallis aquatica angustifolia scutellata. Bauh. pin.
 252.
Anglis, narrow-leaved Water-speedwell.
*Habitat in palustribus, in ericeto Hampstediano et Putne-
 jano, 2. VI.*
8. VERONICA floribus racemosis lateralibus, foliis *chamæ-*
ovatis rugosis dentatis sessilibus, caule debili. *Sp. drys.*
pl. 13.
 Veronica foliis oppositis plicatis dentatis, scapis ex alis
 inferioribus laxè spicatis. *Hort. Cliff.* 8.
Chamædryis sylvestris. Ger. Em. 657. *R. Syn.* 281.
Chamædryis spuria minor rotundifolia. Bauh. pin.
 249.
Chamædryis spuria sylvestris. Park. 107.
Anglis, wild Germander.
Habitat in pratis et pascuis. 2. V.
9. VERONICA racemis lateralibus paucifloris calycibus *montana*.
hirsutis, foliis ovatis rugosis crenatis petiolatis, cau-
 le debili. *Cent. 1. Syst. N.* 849.
 Veronica *chamædryoides* foliis pediculis oblongis in-
 fidentibus. *R. Syn.* 281.
Chamædryi spuriae affinis, rotundifolia scutellata.
Bauh. pin. 249.
Anglis, stalked Speedwell. *Pet. H. Br. t.* 51. f. 4.
Habitat in sylvis humidis et ad sepes, in Charlton Wood,
the lower part, plentifully. Mr. J. Sherard. R.
Syn. 2. V. *** *Pe.*

*** *Pedunculis unifloris.*

arvensis. 10. VERONICA floribus solitariis, foliis cordatis incis-
fis sessilibus.

Veronica floribus solitariis, foliis cordatis incisfis pe-
dunculo longioribus. *Sp. pl.* 13. *Fl. suec.* 20.

Veronica flosculis singularibus, cauliculis adhærenti-
bus. *R. Syn.* 279.

Alfine veronicæ foliis, flosculis cauliculis adhærenti-
bus. *Bauh. pin.* 250.

Alfine foliis veronicæ. *Ger. Em.* 613.

Alfine foliis subrotundis veronicæ. *Park.* 762.

Anglis, Speedwell or Chickweed.

Habitat in agris et muris antiquis. ☉. V.

agrestis. 11. VERONICA floribus solitariis, foliis cordatis incisfis
petiolatis.

Veronica floribus solitariis, foliis cordatis incisfis pe-
dunculo brevioribus. *Sp. pl.* 13. *Fl. suec.* 16.

Veronica floribus singularibus, in oblongis pediculis,
Chamædryfolia. *R. Syn.* 279.

Alfine foliis Triflaginis. *Ger. Em.* 616. *Park.* 764.

Alfine chamædryfolia, flosculis pediculis oblongis infi-
dentibus. *Bauh. pin.* 250.

Anglis, Germander-speedwell, or Chickweed.

Habitat in arvis ruderalis, cultis frequens. ☉ V.

hederifolia. 12. VERONICA floribus solitariis, foliis cordatis planis
quinelobis. *Sp. pl.* 13. *Fl. suec.* 21.

Veronica flosculis singularibus, Hederulæ folio, Mor-
sus gallinæ minor dicta. *R. Syn.* 280.

Alfine Hederulæ folio. *Bauh. pin.* 250.

Alfine Hederulæ folio minor. *Park.* 760: 762.

Alfine hederacea. *Ger. Em.* 616.

Anglis, Ivy-leaved Speedwell, or small Henbit.

Habitat in agris ruderalis, cultis frequens. ☉. IV. V.

triphyllus. 13. VERONICA floribus solitariis, foliis digitato-par-
titis pedunculo brevioribus. *Sp. pl.* 14. *Fl. suec.* 22.

Veronica foliis alternis: inferioribus quinque partitis,
superioribus tripartitis, floribus solitariis. *Hort.*
cliff. 9.

Veronica

Veronica flosculus singularibus, foliis laciniatis, erecta.

R. Syn. 280.

Alfine triphyllos cœrulea. Bauh. pin. 250.

Alfine triphyllos five laciniata. Park. 760.

Alfine recta. Ger. Em. 612.

Anglis, Trifid Speedwell.

Habitat in arvis arenosis, at Rowton in Norfolk, betwixt the town and the highway, twelve miles before you come to Norwich; and at Methwold in Suffolk, between the two windmills and the warren lodge, &c.

J. Willisel. R. Syn. ☉. V. VI.

PINGUICULA. 28.

Cor. ringens, calcarata. Cal. bilabiatus, quinquefidus. Caps. unilocularis.

1. *PINGUICULA nectario cylindraceo longitudine pe- vulgaris.*
tali. *Fl. lapp.* 11. *Fl. suec.* 21. *Sp. pl.* 17. *Roy.*
lugdb. 304.

Pinguicula Gefneri. Bauh. Hist. III. 546. *R. Syn.* 281.

Pinguicula five sanicula Eboracensis. Park. 532. *Ger. Em.* 788.

Sanicula montana flore calcari donato. Bauh. pin. 243.

Anglis, Butterwort, or Yorkshire Sanicle.

Habitat in uliginosis in comitatibus Eboracensi, Westmorlandico, Cumberlandico, &c. on Shooter's Hill, and the little bog at Charleton. R. Syn. 4. V.

2. *PINGUICULA nectarii apice incrassato. Sp. pl.* 17. *lusitanica.*
Pinguicula flore minore carneo. R. Syn. 281. *Hist.*
I. 752.

Viola palustris, Pinguicula dicta; lusitanica. Grif.
lusit.

Anglis, Cornwal Butterwort.

Habitat in Cornubia locis palustribus circa Kilkhampton, et alibi. R. Syn. 4. V.

DIANDRIA MONOGYNIA:

UTRICULARIA. 29.

Corolla ringens, calcarata. *Cal.* diphyllus, æqualis.
Caps. unilocularis.

vulgaris. 1. UTRICULARIA nectario conico. *Fl. lapp.* 14. *Sp.*
pl. 18. *Fl. suec.* 24.

Lentibularia. *Riv. mon* 78. *R. Syn.* 286.

Millefolium aquaticum lenticulatum. *Bauh. pin.* 141.

Millefolium aquaticum flore luteo galericulato. *Bauh.*
hist. III. 783. *Park.* 1258.

Millefolium palustre galericulatum. *Ger. Em.* 828.

Anglis, Common-hooded Milfoil.

Habitat in aquis stagnantibus, paludibus et fossis, in stagnis prope Hornsey. 4. VI. VII.

minor. 2. UTRICULARIA nectario carinato. *Fl. suec.* 29.
Sp. pl. 18.

Lentibularia minor. *Pet. H. Br. t.* 36. *f.* 12. *R.*
Syn. 286.

Millefolium palustre galericulatum minus, flore minore. *Pluk. Ph. t.* 99. *f.* 6.

Anglis, lesser-hooded Milfoil.

Habitat in fossis et paludibus limosis. Found by Mr.
Dent on Feversham-moor in Cambridgeshire. Both
sorts observed by Mr. Lawson in the Ditches by the
Causeway over the Moss to the Fell-end, near Witherslack,
Westmorland; in the River on Hounslow
Heath, by Mr. Dandridge. *R. Syn.* 4. VI. VII.

*** *Flores inferi, monopetali, irregulares, Fructus*
gymnospermi.

LYCOPUS. 31.

Cor. quadrifida: lacinia unica emarginata. *Stam.*
distantia. *Sem.* quatuor, retusa.

europæus. LYCOPUS foliis sinuato-ferratis. *Sp. pl.* 21. *Fl.*
suec. 31.

Lycopus palustris glaber. *Tourn.* 191. *R. Syn.* 236.

Marrubium palustre glabrum. *Bauh. pin.* 230.

Marrubium aquaticum. *Ger. Em.* 700. *vulgare Park.*
 1230.

Anglis,

Anglis, Water Horehound.

Habitat *ad ripas fluviorum et lacuum, solo sabuloso.*

4. VII.

S A L V I A. 36.

Cor. inæqualis. Filamenta transverse pedicello affixa.

1. SALVIA foliis cordato-oblongis crenatis: summis *pratenf*
amplexicaulibus, verticillis subnudis, corollis galea
glutinosi. *Sp. pl.* 25. *Fl. suec.* 32.

Scleara pratenfis foliis ferratis, flore cæruleo. *Tourn.*
179. *R. Syn.* 237.

Horminum pratenfe foliis ferratis. *Bauh. pin.* 238.

Horminum sylvestre vulgare. *Park.* 55.

Horminum sylvestre Fuchsii. *Ger. Em.* 760.

Anglis, Meadow Clary.

Habitat *in septo Cobhamiano in Effexia, a D. Wat-*
sio observatus. 4.

2. SALVIA foliis sinuato-ferratis, corollis calyce angust-*verbenacea* X
rioribus. *Sp. pl.* 25.

Horminum sylvestre Lavendulæ flore. *Bauh. pin.*
239. *Park.* 57. *R. Syn.* 237.

Horminum sylvestre. *Ger. Em.* 760.

Anglis, common English wild Clary.

Habitat *in pratis et pascuis frequens.* 4. V. *

Pharm. Hormini semina.

**** Flores superi.

C I R C Æ A. 24.

Corolla dipetala. Calyx diphyllus superus. Sem. unum
biloculare.

1. CIRCÆA caule erecto, racemis pluribus. *Sp. pl.* 9. *lutetiana.*
Fl. suec. 6.

Circæa lutetiana. Lob. ic. 266. *Ger. Em.* 351. *R.*
Syn. 289.

Circæa lutetiana major. Park. 351.

Solanifolia Circæa dicta major. Bauh. pin. 168.

Anglis, Enchanters Nightshade.

Habitat *in locis umbrosis humidis et ad sepes passim.* 4.

VI. VII.

2. C I R-

* *About Sudbury in Suffolk; Andover in Hants.*

DIANDRIA MONOGYNIA.

- x alpina. 2. CIRCÆA caule adscendente, racemo unico. *Sp. pl.*
 9. *Fl. suec.* 7.
 Circæa calyce colorato. *Fl. lapp.* 3.
 Circæa minima foliis brevioribus dentatis Hall. *helv.*
 456.
 Circæa lutetiana minor. *Park.* 351.
 Circæa minima. *Column. Ecphr.* 2. p. 79. t. 80.
 Solanifolia Circæa alpina. *Bauh. pin.* 168.
Anglis, Mountain Enchanters Nightshade.
 Habitat *ad radices montium in Comitatus Westmor-*
landico, Eboracensi, &c. circa Dallam Tower in
agro Westmorlandico. 4. VIII. At the foot of
Broadway-Hills, before one comes into Broadway
the **D I G Y N I A** *-- left side of the*

ANTHOXANTHUM. 40.

- Cal.* Gluma bivalvis, uniflora. *Cor.* Gluma bival-
 vis, acuminata. *Sem.* unicum.
odoratum. 1. ANTHOXANTHUM spica oblonga ovata, flosculis
 subpedunculatis arista longioribus. *Sp. pl.* 28. *Fl.*
suec. 33.
 Gramen vernum spica brevi laxa. *R. Syn.* 389. 398.
 Gramen pratense spica flavescente. *Bauh. pin.* 3.
Th. 44.
 Gramen Anthoxanthum spicatum. *Bauh. Hist.* II.
 466.
 Gramen alopecurum vernum pratense, spica flavesc-
 ente. *Hist. Ox.* III. p. 193. f. 8. t. 7. f. 25.
Anglis, vernal, or Spring Grass.
 Habitat *in pratis et pascuis* 4. V.

B U F O N I A. 41.

- Cal.* Tetraphyllus. *Cor.* 4petala. *Capf.* unilocula-
 ris bivalvis disperma.
tenuifolia. 1. BUFONIA. *Sp. pl.* 123.
 Alfinoides. *R. Syn.* 346.

Herniaria

DIANDRIA DIGYNIA.

11

Herniaria angustifolium gramineo folio, erecta. *Magn. Hort.* 97. t. 97.

Alfina polygonoides tenuifolia, flosculis ad longitudinem caulis velut in spicam dispositis. *Pluk. Ph.* t. 75. f. 3.

Anglis, Bastard Chickweed.

Habitat in maritimis circa Bostoniam agri Lincolnien-
sis. On Hounslow-heath. Mr. Doody. *R. Syn.*
4. V.

Classis

Classis III.

TRIANDRIA
MONOGYNIA.

* Flores superi.

VALERIANA. 43.

Cal. nullus. *Cor.* monopetala, basi hinc gibba, super. *Sem.* unicum.

officinalis. 1. VALERIANA floribus triandris, foliis omnibus pinnatis. *Mat. Med.* 21. *Sp. pl.* 31. *Fl. suec.* 34. Valeriana sylvestris major. *Bauh. pin.* 164. *Fl. lapp.* 14. *Ger. Em.* 1075. *Park.* 122. *R. Syn.* 200.

β Valeriana sylvestris major montana. *Bauh. pin.* 164. *Anglis*, Great wild Valerian.

Habitat in sepibus, sylvis, paludibus, frequens. β in pascuis montosis. 4. VI.

Pharm. Valerianæ Radix.

dioica. 2. VALERIANA floribus triandris dioicis, foliis radicalibus ovatis, caulinis pinnatis.

Valeriana floribus triandris dioicis, foliis pinnatis integerrimis. *Sp. pl.* 31. *Fl. suec.* 35. *Syst. Nat.* 860.

Valeriana palustris minor. *Bauh. pin.* 164.

Valeriana sylvestris minor. *Park.* 122. *Lob.* 715. *R. Syn.* 200.

Valeriana minor. *Ger. Em.* 1075. *Riv. mon.* 3.

x β Valeriana sylvestris seu palustris minor altera. *R. Syn.* 200.

Anglis, small wild or marsh Valerian. Both varieties in London. *Exham*

Habitat in pascuis humidis. 4. VI.

Locusta. 3. VALERIANA floribus triandris, caule dichotomo, foliis linearibus. *Sp. pl.* 33. *Fl. suec.* 36.

Valerianella arvensis præcox humilis semine compresso. *Mor. Umb.* *R. Syn.* 201.

Vale-

- Valeriana campestris inodora major. *Bauh. pin.* 165.
 Lactuca agnina. *Ger. Em.* 310. *Park.* 812.
 β Valerianellæ vulgaris species major serotina. *Mor. prælud.* 319.
 γ Lactuca agnina seu Valerianella foliis ferratis. *R. Syn.* 201.
Anglis, Lambs-lettuce, or Corn-fallet.
 Habitat in arvis et agris. ☉. IV. V.

C R O C U S. 63.

- Cor.* Sexpartita, æqualis. *Stigmata* convoluta.
 *1. CROCUS spatha univalvi radicale, corollæ tubo *fativus*.
 longissimo. *Sp. pl.* 35.
Crocus, *Bauh. hist. II.* 637. *Ger. Em.* 151. *fativus*.
Bauh. pin. 651.
Anglis, Saffron.
 Habitat in agro Cantabrigienfi, et Essexienfi Saffron-
 walden oppidum colitur. *R. Syn.* 4. VIII.
Pharm. Croci stigma.

I R I S. 57.

- Cor.* Sexpartita. *Petalis* alternis reflexis. *Stigmata* petaliformia.
 1. IRIS corollis imberbibus, petalis interioribus stigma- *Pseudac-*
 te minoribus, foliis ensiformibus. *Syst. Nat.* 863. *rus*.
Sp. pl. 38. *Fl. suec.* 37.
Iris palustris lutea. *Ger. Em.* 50. *R. Syn.* 374.
Acorus adulterinus. *Bau. pin.* 34.
Acorus palustris, five Pseudo-iris et *Iris lutea palustris*.
Park. 1219.
 β *Iris palustris pallida*. *R. Syn.*
Anglis, yellow water Flower-de-luce.
 Habitat in udis, fluviis, paludosis frequens. 4. VII.
Pharm. *Acori palustris Radix*.
 2. IRIS corollis imberbibus, petalis interioribus paten- *fœtidiss-*
 tissimis, caule unangulato, foliis ensiformibus. *Syst.* *ma.*
Nat. 863. *Sp. pl.* 39.
Gladiolus fœtidus. *Bauh. pin.* 30.
Xyris. *Ger. Em.* 60. seu spatula fœtida. *Park.* 256.
Anglis, stinking Gladdon, or Gladwyn.

Habitat

TRIANDRIA MONOGYNIA.

Habitat in sylvis et ad sepes, prope Hornsey; in sylva prope Charleton in Cantio. 4. VI. VII.

Pharm. Iris foetidæ Radix.

*** Flores Graminei valvulis Glumæ calycinæ.

SCHOENUS. 60.

Glumæ paleaceæ, univalves, congestæ. Cor. o. sem. unicum, subrotundum.

* Culmo tereti.

Mariscus. 1. SCHOENUS culmo tereti, foliis margine dorsoque aculeatis. *Fl. suec.* 38. *Sp. pl.* 42.

Cyperus longus inodorus germanicus. *Bauh. pin.* 14.

Cyperus longus inodorus sylvestris. *Ger. Em.* 29. *Park.* 1263. *R. Syn.* 426.

Pseudo-cyperus palustris foliis et carina ferratis. *Sch. agrost.* 376.

Anglis, long-rooted bastard Cyperus.

Habitat in paludibus; on Hinton Moor, near Cambridge, plentifully; also in boggy places by the river Thame at Dorsethill, near Tamworth, in Warwickshire; and by the sea side between Pensans and Marketjeu in Cornwall. 4. VII. VIII.

nigricans. 2. SCHOENUS culmo tereti nudo, capitulo ovato, involucri diphylli valvula altera subulata longa. *Sp. pl.* 43. *Fl. suec.* 39. *Dalib. parif.* 113.

Juncus lævis minor, panicula glomerata nigricante. *Cat. pl. An.* 174. *R. Syn.* 430. *Hist. Ox.* III. 243. t. 10. f. 28.

Gramen sparteum nigro capitulo. *Merr. pin.*

Anglis, round black-headed Bog-rush.

Habitat in paludibus cespitosus, et turfosis. 4. VI.

ferrugineus. 3. SCHOENUS culmo tereti nudo, spica duplici, involucri valvula majore spicam æquante. *Sp. pl.* 43. *Fl. suec.* 40.

Juncello accedens graminifolia plantula capitulis Armeriæ proliferæ. *Lbwyd. R. Syn.* 430.

Gramen Cyperoides minimum Caryophylli proliferi capitulo simplici squamato. *Hist. Ox.* III. 246. t. 12. f. 40. Anglis,

Anglis, brown bastard Cyperus.

Habitat in *paludibus turfosis*, on the mountains of Wales,
R. Syn. in Westmorlandia frequens. 4. VII.

** *Culmo triquetro.*

4. SCHOENUS culmo subtriquetro nudo, spica disticha, *compressus.*
 involucro monophyllo. *Sp. pl.* 43.

Gramen cyperoides spica simplici compressa disticha.
Pluk. Ph. t. 34. *f.* 9. *Sch. Agrost.* 490. *Hist. Ox.*
III. 245. *R. Syn.* 425.

Anglis, compressed bastard Cyperus.

Habitat in *paludibus cespitosis*, in comitatu Westmor-
 landico frequens; about Chiffelhurst in Kent; and
 in the Rill near Dulwich Wells. *R. Syn.* 4. VII.

5. SCHOENUS culmo subtriquetro folioso, floribus fasciculatis, foliis setaceis. *Sp. pl.* 44. *Fl. suec.* 41.

Cyperus minor palustris hirsutus, paniculis albis, paleaceis. *Hist. Ox. III.* 239. *t.* 9. *f.* 39. *R. Syn.*
 427.

Gramen cyperoides albis glumis. *Merr. pin.*

Gramen junceum leucanthemum. *Ger. Em.* 30.

Anglis, white flowered Rush-grass.

Habitat in *palustribus*. Mr. Doody observed it on Bagshot Heath; Mr. Petiver in a low bog between Wickham and Croydon. *R. Syn. in comitatibus*,
Eboracensi, Lancastriensi ubique. 4. VII.

C Y P E R U S. 61.

Glumæ paleaceæ, distiche imbricatæ. *Cor. o. Sem.*
 unicum nudum.

1. CYPERUS culmo triquetro folioso, umbella foliosa *longus.*
 supra decomposita; pedunculis nudis, spicis alternis. *Roy. lugdb.* 50. *Mat. med.* 29. *Sp. pl.* 45.

Cyperus longus. *Ger. Em.* 30. odoratus. *Park.* 145.
R. Syn. 425.

Cyperus odoratus radice longa. f. Cyperus Off. *Bauh.*
pin. 14.

Anglis, sweet Cyperus, or English Galingale.

Habitat in *paludibus*. Found by Mr. Newton. in the
isle

isle of Purbeck, Dorsetshire, near a chapel on the side that looks towards Portland island. R. Syn. 4. VII.

Pharmac. Cyperi longi Radix.

SCIRPUS. 62.

Glumæ paleaceæ, undique imbricatæ. Cor. o. Sem. unicum imberbe.

** Spica unica.*

palustris. 1. SCIRPUS culmo tereti nudo, spica subovata imbricata. *Sp. pl.* 47.

Scirpus equiseti capitulo majori. *Tourn.* 528. *R. Syn.* 428. *Sch. agrost.* 360.

Juncus equiseti capitulis, major. *Bauh. pin.* 12.

Juncus aquaticus capitulis equiseti. *Park.* 1196.

Juncus minor capitulis equiseti. *Ger. Em.* 34.

Anglis, Club-rush.

Habitat in fossis, paludibus, fluviiis frequens. 4. VII.

cespitosus. 2. SCIRPUS culmo striato nudo, spica bivalvi terminali longitudine calycis, radicibus squamula interstinctis. *Fl. suec.* 43, *Sp. pl.* 48.

Scirpus montanus capitulo breviori. *Tourn.* 528. *Sch. agr.* 363. *t.* 7. *f.* 18. *R. Syn.* 429.

Juncus parvus palustris, cum parvis capitulis equiseti. *Pluk. Ph.* *t.* 406.

Anglis, Dwarf Club-rush.

Habitat in paludibus cespitosus; in pratis udis et locis palustribus circa Middleton vicum in agro Warvicensi.

Mr. Doody observed it in very dry places among Heath, and near a foot high. *R. Syn.* 4. VII.

acicularis. 3. SCIRPUS culmo tereti nudo setiformi, spica ovata bivalvi, seminibus nudis. *Fl. suec.* 44. *Sp. pl.* 48. *Dalib. parif.* 17.

Scirpus minimus capitulis equiseti. *Cat. Giff.* 165. *R. Syn.* 429.

Juncellus omnium minimus capitulis equiseti. *Hist. Ox.* III. 145. *t.* 10. *f.* 37. *Pluk. Ph.* *t.* 40. *f.* 7.

Anglis, the least upright Club-rush.

Habita

Habitat in palustribus ; in ericeto Hounsflemano versus Hampton, invenit D. Doody. R. syn. 4. VIII.

4. SCIRPUS culmis teretibus nudis alternis caule folioso *fluitans*, flaccido. *Sp. pl.* 48.

Scirpus foliis linearibus planis alternatim fasciculatis, spica terminali. *Guet. Stamp.* 141.

Scirpus equiseti capitulo minor. *Sch. agr.* 365. t. 7. f. 20. R. Syn. 431.

Juncellus, capitulis equiseti, minor fluitans. *Bauh. pin.* 12.

Gramen junceum clavatum minimum. R. *Hist.* 1310. *Hist. Ox.* III. 230. t. 10. f. 31. *Pluk. Ph.* t. 35. f. 1.

Anglis, floating Club-rush.

Habitat in fossis et paludibus, in the ponds of Wandsworth Common, next the new churchyard in Surrey. 4. VII. VIII.

** *Culmo tereti polystachio.*

5. SCIRPUS culmo tereti nudo, spicis ovatis pluribus *lacustris*, pedunculatis terminalibus. *Fl. suec.* 46. *Sp. pl.* 48.

Scirpus palustris altissimus. *Tourn.* 528. R. Syn. 428.

Juncus maximus s. Scirpus major. *Bauh. pin.* 12. *theatr.* 178.

Juncus lævis maximus. *Park.* 1191.

Juncus aquaticus maximus. *Ger. Em.* 35.

β Juncus s. Scirpus medius. *Bauh. pin.* 12.

γ Juncus aquaticus major caule carinato. *Doody.* R. Syn. *Anglis*, Bull-rush.

Habitat in stagnis et fluviis frequens. 4. VII. VIII.

6. SCIRPUS culmo tereti nudo, spicis subglobosis glo- *Holofcho-* meratis pedunculatis, involucro diphylo inæquali *enus*, mucronato. *Sp. pl.* 49.

Scirpus maritimus, capitulis rotundioribus glomeratis. *Tourn.* 528. R. Syn. 429.

Juncus acutus maritimus, capitulis rotundis. *Bauh. pin.* 11.

Juncus acutis maritimus alter. *Park.* 1196.

C

Anglis,

TRIANDRIA MONOGYNIA.

Anglis, Round-headed Bull-rush.

Habitat in litoribus maritimis. Found by Mr. Stephens in Brounton Boroughs in Devonshire; in Somersetshire and Hampshire. *Peta conc. gr. R. Syn.* 429. 4. VII.

- setaceus.* 7. SCIRPUS culmo nudo setaceo, spicis lateralibus subfoliariis sessilibus. *Sp. pl.* 49. *Fl. suec.* 45. Scirpus foliaceus humilis. *C. Giff.* 158. *R. Syn.* 430. Juncellus omnium minimus. Chamæschænos. *Lob. adv.* 44. *Hist. Ox. III.* 232 t. 10. f. 23. Juncellus inutilis sive Chamæschænos. *Bauh. pin.* 12. pr. 22. Gramen junceum minimum capitulo squamoso. *Bauh. pin.* 6. pr. 13. Gramen junceum maritimum exile Plimmofstii. *Park.* 1270. ic. 1271. *descr.* *Anglis*, the least Rush. Habitat in arenosis humidioribus. ☉. VII. VIII.

*** Culmo triquetro, Panicula nuda.

- mucronata.* 8. SCIRPUS culmo triquetro nudo acuminato, panicula conglomerata nuda. *Sp. pl.* 50. *tus.* Juncus acutus maritimus caule triangulo. *Bauh. pin.* 11. pr. 22. *Fl.* 175. *Park.* 1194. *Hist. Ox.* 232. t. 10. f. 20. Juncus acutus maritimus caule triquetro, rigido, mucrone pungente. *Pluk. Ph.* t. 40. f. 1. *R. Syn.* 429. β. Juncus acutus maritimus, caule triquetro maximo molli, procerior nostras. *Pluk. Ph.* t. 40. f. 2. *Anglis*, pointed Bull-rush. Habitat in litoribus maritimis; ad ripam Thamefis fluvii tam infra quam supra Londinum. *R. Syn.* 4. VII. VIII.

**** Culmo triquetro, Panicula foliacea.

- maritimus.* 9. SCIRPUS culmo triquetro, panicula conglobata foliacea, spicularum squamis trifidis; intermedia subulata. *Sp. pl.* 51. *Fl. suec.* 47.

Cyperus

Cyperus culmo triquetro, panicula foliacea, pedunculis simplissimis spicis confertis. *Roy. lugdb.* 50.

Gramen cyperoides panicula sparfa majus. *Bauh. pin.* 6.

Gramen cyperoides palustris panicula sparfa majus. *Park.* 1266. *R. Syn.* 425.

β *Cyperus rotundus inodorus Anglicus.* *Bauh. pin.* 14. *Anglis*, round rooted bastard *Cyperus*.

Habitat in littoribus maritimis, et ad ripas fluviorum frequens. 4. VIII.

10. SCIRPUS culmo triquetro folioso, umbella foliacea, pedunculis nudis supra-decompositis, spicis confertis. *Sp. pl.* 51. *Fl. suec.* 48. *sylvaticus.*

Gramen cyperoides miliaceum *Bauh. pin.* 6. *Th.* 90.

Pseudo-cyperus miliaceus. *Park.* 1171.

Cyperus gramineus *Bauh. Hist. II.* 504. *R. Syn.* 426. *miliaceus.* *Ger. Em.* 30.

Anglis, Millet *Cyperus-grafs.*

Habitat in aquis, et sylvis humidis inter Hampstead et Highgate, by the Thame side near Tamworth in Warwickshire; in a brook near Haverford-west in Pembrokehire. Mr. Dale observed it by the river Blackwater, at a Mill below Bocking in Essex. *R. Syn.* 4. VII.

ERIOPHORUM. 63.

Glumæ paleaceæ, undique imbricatæ. Cor. o. Sem. 1. lana longissima cinctum.

1. ERIOPHORUM culmis teretibus, foliis planis, spicis pedunculatis. *Sp. pl.* 52. *Fl. suec.* 49. *Dalib. on. paris.* 18. *polystachis.*

Eriophorum spicis pendulis. *Fl. lapp.* 28. *Roy. Lugdb.* 51.

Gramen pratense tomentosum panicula sparfa. *Bauh. pin.* 4.

Gramen tomentarium. *Ger. Em.* 29.

Gramen junceum lanatum, vel juncus bombycinus vulgaris. *Park.* 1271.

Linagrostis. *Tab. Hist.* 599. *R. Syn.* 435. *Anglis*, Cotton-grafs.

TRIANDRIA MONAGYNIA.

Habitat in pratis humidis, paludibus, cespitosis. 4. VI. VII.

- vagina-*
tum. 2. ERIOPHORUM culmis vaginatis teretibus, spica
paleacea. *Sp. pl.* 52. *Fl. suec.* 50. *Dalib. pa-*
ris. 18.
Juncus capitulo lanuginoso. f. Schœnolaguros. *Bauh.*
pin. 12.
Gramen juncoides lanatum alterum Danicum. *Park.*
1271.
Anglis, Hare's-tail-rush.
Habitat in paludibus alpinis, solo cespitose, in comitatu
Westmorlandico frequens. On the great bog beyond
Joan Cole's, towards Croydon, Mr. Sherard. *R.*
Syn. 4. II.—IV.

NARDUS. 65.

Calyx nullus. *Cor.* bivalvis.

- stricta.* 1. NARDUS spica setacea recta. *Sp. pl.* 53. *Fl. suec.*
52.
Gramen sparteum juncifolium. *Bauh. pin.* 5. *R. Syn.*
393.
Spartum parvum Batavicum et Anglicum. *Park.* 1199.
Spartum nostras parvum lobelio. *Ger. Em.* 43.
Anglis, Mat-grafs.
Habitat in palustribus et ericetis frequens. 4. VI. VII.

DIGYNIA.

PHALARIS. 69.

Cal. bivalvis, carinatus, longitudine æqualis, corol-
lam includens.

- canarien-*
sis. * 1. PHALARIS panicula subovata spiciformi, glumis
carinatis. *Sp. pl.* 54.
Phalaris major femine albo. *Bauh. pin.* 28. *Hist. Ox.*
186. t. 3. f. 1.
Phalaris. *Ger. Em.* 86.
Anglis, manured Canary grafs.
Habitat in ruderalis et ad vias. ☉. VI.—IX.

2. PHA-

2. PHALARIS spica ovata ; calycis valvulis dorso cili- *arenaria.*
 atis, culmo ramoso.
 Phleum spica ovata ciliata, caule ramoso. *Sp. pl.* 60.
 Gramen typhinum maritimum minus. *R. Syn.* 398.
Pluk. Ph. t. 33. *f.* 8.
 Gramen typhinum Danicum minus. *Park.* 1170.
Anglis, Sea Canary-grafs.
 Habitat in arena mobili juxta mare. ☉. VII.
3. PHALARIS panicula oblonga ventricosa. *Sp. pl.* 55. *arundina-*
 Arundo foliis planis panicula spicata, spiculis con- *cea.*
 fertis. *Hort. cliff.* 26.
 Gramen arundinaceum acerosa gluma nostras. *Park.*
 1273; *Hist. Ox.* III. 203. *t.* 6. *f.* 14. *R. Syn.*
 400.
 Gramen aquaticum paniculatum latifolium. *Bauh.*
pin. 6.
 β Gramen arundinaceum acerosa gluma Jersejanum.
D. Sherard.
Anglis, Reed Canary-grafs.
 Habitat ad ripas fluviorum. 4. VII.

P A N I C U M. 70.

Cal. trivalvis ; valvula tertia minima.

1. PANICUM spica tereti, involucellis bifloris fascicu- *viride.*
 lato-pilosis, seminibus nervosis. *Syst. Nat.* II. 870.
 Gramen paniceum spica simplici lævi. *R. Syn.* 393.
 Gramen panici effigie spica simplici. *Ger. Em.* 17.
 Gramen paniceum spica simplici ἐλυμάγῳσις. *Bauh.*
pin. 8. *Park.* 1154.
 β Gramen paniceum spica aspera. *Bauh. pin.* 8.
Anglis, Green Panick-grafs.
 Habitat in arvis prope Battersea copiose. ☉. VII.
 VIII.
- 2 PANICUM spicis alternis conjugatisque, spiculis sub- *Crusgalli.*
 divis, glumis aristatis hispidis, rachi quinquangu-
 lari. *Sp. pl.* 56.
 Gramen paniceum spica divisa. *Bauh. pin.* 8. *R.*
Syn. 394.

TRIANDRIA DIGYNIA.

Panicum vulgare. Ger. Em. 85.

Panicum sylvestre Herbariorum. Park. 1154.

Anglis, Loose Panick-grafs.

Habitat cum priore. ☉. VIII.

sanguinalis. 3. *PANICUM* spicis digitatis basi interiore nodosis, flosculis geminis nuticis, vaginis foliorum punctatis. Sp. pl. 57.

Gramen dactylon latiore folio. Bauh. pin. 8. Th. 114.

Ischæmon sylvestre latiore folio. Park. 1178.

Ischæmon vulgare. Ger. Em. 27.

Anglis, Cock's-foot Panick-grafs.

Habitat in arvis inter Battersea et Wandsworth; plentifully in the ploughed fields about Elden in Suffolk.

T. Willifell. R. Syn. ☉. VIII.

dactylon. 4. *PANICUM* spicis digitatis patentibus; basi interiore villosis, floribus solitariis, sarmentis repentibus. Sp. pl. 58.

Gramen Dactylon, folio arundinaceo, majus aculeatum. Bauh. pin. 7.

Gramen Dactyloides radice repente. Ger. Em. 28.

Gramen canarium ischæmi paniculis. Park. 1179.

Gramen repens cum panicula graminis Mannæ. Bauh. Hist. II. 459.

Anglis, creeping Panick-grafs.

Habitat in arenosis. Found by Mr. Newton on the sandy shores between Pensans and Marketjeu in Cornwall plentifully. R. Syn. 4. VII.

PHLEUM. 71.

Cal. bivalvis, linearis truncatus, apice bicuspidato.
Cor. inclusa.

pratense. 1. *PHLEUM* spica cylindrica longissima; glumis ciliatis.

Pleum spica cylindrica longissima. Fl. lapp. 26. Fl. suec. 50. Sp. pl. 54.

Gramen typhoides maximum spica longissima. Bauh. pin. 4. Th. 49. Hist. Ox. III. 188. t. 4. f. 1.

Gramen

- Gramen typhinum majus seu primum. *Ger. Em.* 11.
R. Syn. 398.
- Gramen typhinum medium f. vulgatissimum. *Park.*
 1170.
- β Gramen typhinum minus. *Ger. Em.* 11. *Park.* 1170.
Anglis, Meadow Cat's-tail-grafs.
Habitat in pratis et pascuis. 11. VII.
2. PHLEUM panicula cylindrica spiciformi, glumis gla- *panicula-*
 bris, culmo subramoso. *tum.*
Anglis, branched Cat's-tail-grafs.
Habitat in pratis infra King's Weston prope Bristol-
um. 11. VII.
3. PHLEUM spica ovato-oblonga basi sterili, culmo *nodosum*,
 subrepente, radice bulbosa. *Syst. Nat.* 871.
 Gramen nodosum spica parva. *Baub. pin.* 2. pr. 3.
Anglis, bulbose Cat's-tail-grafs.
Habitat in pascuis et ad vias ubique. 11. VII. VIII.

ALOPECURUS. 72.

Calyx bivalvis. *Corolla* univalvis.

1. ALOPECURUS panicula cylindrica spiciformi, cul- *pratenfis*,
 mo erecto.
 Alopecurus culmo spicato erecto. *Roy. lugdb.* 54. *Fl.*
succ. 59. *Sp. pl.* 60.
 Gramen alopecuroides majus. *Ger. Em.* 10.
 Gramen phalaroides spica molli sive Germanicum.
Baub. pin. 4.
 Gramen phalaroides majus, sive Italicum. *Baub.*
pin. 4. *Park.* 1164.
Anglis, Meadow Fox-tail-grafs.
Habitat in pratis et pascuis ubique. 11. V.
2. ALOPECURUS spica cylindrica longissima glumis *myosuroides*,
 glabris; culmo suberecto.
 Gramen myosuroides majus, spica longiore, aristis
 rectis. *R. Syn.* 397.
 Gramen typhoides spica angustiore. *Baub. pin.* 4.
Th. 53.

TRIANDRIA DIGYNIA.

Gramen alopecuroides spica longa majus et minus.
Park. 1169.

Gramen alopecuroides minus. *Ger. Em.* 10.

β Gramen myosuroides minus, spica brevior aristis recurvis. *R. Syn.* 397.

Anglis, Field Fox-tail-grass.

Habitat in arvis, et ad vias. 4. VII. VIII.

bulbosus. 3. ALOPECURUS spica cylindrica, culmo erecto, radice bulbosa.

Gramen myosuroides nodosum. *R. Syn.* 397.

Anglis, bulbous Fox-tail-grass.

Habitat in pascuis et pratis. In the first field next the road before you go into Northfleet. *Dr. Wilmer.* 4. VI.

geniculatus. 4. ALOPECURUS culmo spicato infracto. *Fl. suec.* 60. *Sp. pl.* 60.

Gramen aquaticum geniculatum spicatum. *Bauh. pin.* 3. *Th.* 41. *R. Syn.* 396.

Gramen fluviatile spicatum. *Ger. Em.* 14.

Gramen aquaticum spicatum. *Park.* 1275.

β Gramen fluviatile album. *Tab. Icon.*

Anglis, Flote Fox tail grass.

Habitat in aquosis. 4. VI.—VIII.

S T I P A. 84.

Calyx bivalvis, uniflorus. *Cor.* valvula exteriori aristata terminali : basi articulata.

pennata. 1. STIPA aristis lanatis. *Sp. pl.* 78.

Gramen sparteum pennatum. *Bauh. pin.* 5. *Th.* 70. *R. Syn.* 393.

Spartum Austriacum pennatum. *Clus. Hist.* 2. p. 221. *Ger. Em.* 42.

Anglis, Feather-grass.

Habitat in montosis. Found by *Dr. Richardson* in company with *Thomas Lawson*, on the Limestone rocks hanging over a little valley called Longleadale, about six miles north of Kendal in Westmoreland. *R. Syn.* 4. VII. VIII.

DACTYLIS. 80.

Cal. bivalvis, acutus; altera valvula majore.

1. DACTYLIS spicis sparsis secundis scabris numerosis. *cynosuroides.*
Læfl. hisp. 115. Sp. pl. 71.
Spartum Effexianum, spica gemina clausa. Pet. conc.
gr. 35. R. Hist. III. App. 248. Syn. 393.
Gramen sparteum ferotinum, capite bifido vel gemino.
Merr. pin.
Anglis, Smooth Cock's-foot-grass.
Habitat in paludibus. At Crixey Ferry in Essex. Mr.
Buddle found it in the marshes on the river Wallfleet,
near Farnbridge Ferry in Dengey Hundred, in Es-
sex; and Mr. J. Sherard in the Swail, near the
mouth of Feversham Creek, about August and Sep-
tember. R. Syn. ☉.

2. DACTYLIS panicula secunda glomerata. *Sp. pl. glomeratus.*
71. Fl. 87.
Poa flosculus confertis uno versu dispositis. Hort.
cliff.
Gramen spicatum folio aspero. Bauh. pin. 3. pr. 59.
Th. 45.
Gramen pratense spica multiplici rubra. Park. 1161.
Gramen panicula torosa pratense asperum. Hist. Ox.
III. 202. t. 6. f. 38.
Calamagrostis torosa panicula. Park. 1182.
Anglis, rough Cock's-foot-grass.
Habitat in pratis et pascuis. 4. VI.—VIII.

MILIUM. 73.

Cal. bivalvis, uniflorus, corollam subexcedens. Stigmata penicilliformia.

1. MILIUM floribus paniculatis dispersis muticis. *Sp. pl. effusum.*
61. Fl. suec. 61.
Gramen miliaceum. Lob. ic. 3. Ger. Em. 6. Bauh.
hist. II. 462. R. Syn. 402.
Gramen sylvaticum panicula miliacea sparsa. Bauh.
pin. 8.
Gramen miliaceum vulgare. Park. 1153.

Anglis,

TRIANDRIA DIGYNIA.

Anglis, Millet-grafs.

Habitat in *sylvis humidis frequens*. ☉. VI. VII.

AGROSTIS. 74.

Cal. bivalvis, uniflorus, corolla paulo minor. *Stigmata* longitudinaliter hispida.

* *Aristate*.

*spicaven-
ti.*

1. AGROSTIS petalo exteriori arista recta stricta longissima, panicula patula. *Syst. Nat.* 872. *Sp. pl.* 61. *Fl. suec.* 62.

Gramen segetum panicula arundinacea. *Bauh. pin.* 3. *Tb.* 35.

Gramen agrorum. *Lobelii. Bauh. hist. II.* 461.

Gramen agrorum venti spica. *Park.* 1158.

Gramen arundinaceum. *Ger. Em.* 5.

Gramen miliaceum majus, glumis aristatis, spadiceis et pallidis. *Pet. conc. gr. n.* 122. 123. *R. Syn.* 405.

Anglis, Silky Bent-grafs.

Habitat in *arvis passim*. ☉. VII.

canina.

2. AGROSTIS calycibus coloratis, petalorum arista dorsali recurva. *Fl. suec.* 65. *Sp. pl.* 62.

Gramen supinum canenum paniculatum, folio varians. *Bauh. pin.* 1.

Gramen paniculatum supinum, ad infima culmorum genicula, foliorum capillarium fasciculis donatum. *Sch. agr.* 141.

Anglis, brown Bent-grafs.

Habitat in *pratibus humidis frequens*. 4. VII. VIII.

rubra.

3. AGROSTIS panicula lanceolata spiciformi, calyce flosculo triplo longiore, petalorum arista dorsali recurva.

Agrostis panicula parte florente patentissima, petalo exteriori glabro terminato arista tortili recurva. *Fl. suec.* 64. *Sp. pl.* 62.

Panicum ferotinum arvense spica pyramidata. *Tourn.* 515. *R. Syn.* 394.

Gramen ferotinum spica laxa pyramidalis. *R. hist.* 1288:

Gramen

Gramen alopecuro accedens, ex culmi geniculis spicas cum petiolis longiusculis promens. *Pluk. phyt. t. 33. f. 6.*

Anglis, red Bent-grafs.

Habitat in pratis et pascuis. In Insula Shepey copiose.

☉. VII.

**** Muticæ.**

4. AGROSTIS paniculæ ramulis divaricatis muticis, cul- *stolonifera* mo repente, calycibus æqualibus. *Fl. suec. 65. Sp. pl. 62.*

Agrostis culmo repente foliis radicalibus brevior, folii suprema vagina ventricosa. *Fl. suec. ed. I. 61.*

Gramen montanum miliaceum minus, radice repente. *R. Syn. 402.*

Gramen caninum supinum minus. *Bauh. pin. 1. Sch. agr. 128.*

- β Gramen miliaceum folio angustissimo. *Pet. conc. gr. 116.*

- γ Gramen miliaceum angustifolium, glumis perexiguis. *Pet. conc. gr. 117.*

Anglis, creeping Bent-grafs.

Habitat in pratis et pascuis subhumidis sed rarius. 4. VIII.

5. AGROSTIS panicula coarctata mutica, calycibus æ- *palustris.* qualibus hispidiusculis coloratis ; culmo repente.

Gramen miliaceum majus, panicula spadicea. *Pet. conc. gr. 118. R. Syn. 464.*

- β Gramen miliaceum majus panicula viridi. *Pet. conc. gr. 119.*

Anglis, Marsh Bent-grafs.

Habitat in fossis et locis uliginosis frequens. 4. VII.

6. AGROSTIS panicula capillari patente, calycibus *capillaris.* subulatis æqualibus hispidiusculis coloratis flosculis muticis. *Roy. lugdb. 59. Dalib. parif. 23. Sp. pl. 62.*

Gramen montanum panicula spadicea delicatiore. *Bauh. pin. 3.*

Gramen miliaceum locustis minimis, panicula fere arundinacea. *R. meth. em. 177. Syn. 402.*

Gramen

Gramen pratense vulgare panicula fere arundinacea.

Baub. hist. II. 461.

Anglis, Fine Bent-grafs.

Habitat in pratis et pascuis ubique. 4. VIII.

sylvatica. 7. AGROSTIS panicula coarctata mutica, calycibus æqualibus, corolla ante florescentiam calyce brevior, postea duplo longior.

Gramen miliaceum sylvestre glumis oblongis. *Pet. oonc. gr. 121. R. Syn. 404.*

Anglis, Wood Bent-grafs.

Habitat in sylvis humosis. First discovered by Mr. Buddle in Bishop's-wood, Hampstead. R. Syn. in sylvula Hornsey-wood dicta. 4. VIII.

minima. 8. AGROSTIS panicula mutica filiformi. *Sp. pl. 63.*
Nardus panicula spicata, floribus solitariis alternis
Guett. stamp. 155. Dalib. parif. 155.

Gramen minimum paniculis elegantissimis. *Baub. pin. 2. Sch. agr. 40. Hist. Ox. III. 200. t. 2. f. 10.*

Gramen minimum. *Baub. hist. II. 465. Dalech. hist. 425.*

Gramen minimum anglo-britanicum. *Lob. ill. 20. R. Syn. pl. dub.*

Anglis, small Bent-grafs.

Habitat in Wallia D. Stillingfleet invenit.

*** Flores biflori vagi.*

A I R A. 75.

Cal. bivalvis, biflorus, flosculi absque interjecto rudimento.

** Muticæ.*

cristata. 1. AIRA panicula spicata, floribus muticis pedunculo longioribus glumis acuminatis inæqualibus. *Roy. Lugdb. 61. Sp. pl. 63.*

Gramen spica cristata subhirsutum. *Baub. pin. 2. Hist. Ox. III. 194. t. 4. f. 7.*

Gramen pumilum spica purpureo-argentea molli. *R. Syn. 396.*

Anglis,

Anglis, crifted Hair-grafs.

Habitat in pratis montanis *sterilibus*. On Blackheath.

Pet. conc. gr. 4. VII. VIII.

2. AIRA panicula coarctata, floribus pedunculatis muti- *cærulea*.
cis convoluto-subulatis, foliis planis. *Syst. Nat.*
873. *Sp. pl.* 63. *Fl. suec.* 67.

Poa spiculis subulatis, panicula rara contracta. *Fl.*
lapp. 29.

Gramen pratense ferotinum panicula longa purpuraf-
cente. *R. Syn.* 404. *Hist. Ox. III.* 201. *t.* 5.
f. 22.

Gramen pratense spica Lavendulæ. *Merr. Pin.*

Anglis, Purple Hair-grafs.

Habitat in locis uliginosis, cespitosis, sterilissimis, fre-
quens. 4. VIII.

3. AIRA panicula patente, floribus muticis lævibus ca- *aquatica*.
lyce longioribus, foliis planis. *Syst. Nat.* 873. *Fl.*
suec. 68. *Sp. pl.* 64.

Aira culmo inferiore repente, flosculis muticis calyce
longioribus, altero pedunculato. *Roy. lugdb.* 60.

Gramen miliaceum aquaticum. *R. Syn.* 402. *Sch.*
agr. 218.

Gramen caninum supinum paniculatum dulce. *Bauh.*
pin. 2. *Bauh. hist. II.* 459.

Gramen miliaceum fluitans suavis saporis. *Merr. pin.*

Gramen exile tenuifolium Canario simile. f. Gramen
dulce. *Park.* 1174.

Anglis, Water Hair-grafs.

Habitat in aquosis et ad ripas fluviorum. 4. VI. VII.

** *Aristate.*

4. AIRA foliis planis, panicula patente, petalis basi vil- *cæspitosa*.
losis aristatisque; arista erecta brevi. *Fl. suec.* 70.
Sp. pl. 64.

Gramen fegetum altissimum panicula sparsa. *Bauh.*
pin. 3. *Hist. Ox. III.* 199. *t.* 5. *f.* 1.

Gramen fegetum panicula speciosa. *Park.* 1158.

Gramen fegetale. *Ger. Em.* 5.

Gramen

Gramen miliaceum segetale majus. *Pet. conc. gr.*

114. *R. Syn.* 403.

Anglis, Turfy Hair-grafs.

Habitat in pratis et sylvis humidis frequens. 4. VII. VIII.

flexuosa. 5. AIRA foliis setaceis, culmis subnudis, panicula divaricata pedunculis flexuosis. *Fl. suec.* 71. *Sp. pl.* 65.

Aira panicula rara calycibus fuscis. *Fl. lapp.* 49.

Gramen paniculatum locustis parvis purpuro-argenteis majus et perenne. *R. Syn.* 407.

Gramen avenaceum panicalatum alpinum, foliis capillaceis brevibus, locustis purpuro-argenteis et aristatis *Sch. agr. t.* 4. *f.* 4.

Gramen avenaceum capillaceum minoribus glumis. *Bauh. pin.* 10.

montana. 3. Aira foliis setaceis panicula angustata, flosculis basi pilosis aristatis; arista tortili longiore. *Fl. suec.* 72. *Sp. pl.* 65.

Gramen nemorosum paniculis albis, capillaceo folio. *Bauh. pin.* 7. *pr.* 14. *th.* 97. *Hist. Ox.* III. 300. *t.* 7. *f.* 6. *R. Syn.* 407.

Anglis, Mountain Hair-grafs.

Habitat in ericetis et pascuis sterilibus. 4. VII. VIII.

setacea. 6. AIRA foliis setaceis, panicula secunda coarctata, aristis flosculis duplo longioribus recurvis.

Anglis, small leaved Hair-grafs.

Habitat in ericeto Strattoniensi in comitatu Norfolk. D. Stillingfleet reperit. 4. VII. VIII.

canescens. 7. AIRA foliis setaceis; summo paniculam inferne obvolvante. *Roy. lugdb.* 60. *Fl. suec.* 74. *Sp. pl.* 65.

Aira foliis setaceis aristis ex basi glumarum calycem æquantibus. *Hort. cliff.* 495.

Gramen foliis junceis, radice jubata. *Bauh. pin.* 5. *Th.* 72.

Gramen miliaceum maritimum molle. *Pet. conc. gr.* 126. *R. Syn.* 405.

Anglis, Grey Hair-grafs.

Habitat in arenosis maritimis. 4. VII.

8. AIRA foliis setaceis; vaginis angulatis floribus pa-
miculato-spicatis, flosculis basi aristatis. *Fl. suec.*

75. *Sp. pl.* 65.

Gramen parvum præcox panicula (potius spica) laxa
canescente. *R. Syn.* 407. *Pluk. ph. t.* 33. *f.* 9.

Gramen minimum spica brevi habitiore nostrum. *R.*

Syn. ed. 2. *p.* 252.

Anglis, early Hair grafs.

Habitat in arenosis et sterilibus frequens. ☉. V. VI.

9. AIRA foliis setaceis, panicula divaricata, floribus a-
ristatis distantibus. *Sp. pl.* 66. *caryophyl-
lea.*

Aira foliis subulatis, aristis a basi flosculorum tortili-
bus, flosculo altero pedunculato. *Roy. lugdb.* 60.

Dalib. parif. 26.

Gramen paniculatum, locustis purpureo-argenteis an-
nuum. *R. Syn.* 407.

Gramen paniculatum purpureo-argenteum, locustis
parvis annum. *Hist. Ox. III.* 200. *t.* 5. *f.* 11.

Caryophyllus arvensis glaber minimus. *Baub. pin.*

105.

Anglis, Silver Hair-grafs.

Habitat in pascuis arenosis ubique. ☉. VII.

MELICA. 76.

Calyx bivalvis, biflorus. *Rudimentum* floris inter
flosculos.

1. MELICA petalis imberbibus, panicula nutante sim-
plici. *Sp. pl.* 66. *nutans.*

Melica floribus sub culmo pendulis. *Fl. lapp.* 36.

Gramen avenaceum nemorense, glumis rarioribus
exfusco xerampelinis. *R. Syn.* 403.

Gramen avenaceum locustis rarioribus. *Baub. pin.* 10.

Gramen avenaceum spica mutica rariore gluma. *Hist.*
Ox. III. t. 7. *f.* 49.

Gramen locustis rubris. *Park.* 1151. *Baub. hist.* II.
434.

- β Gramen avenaceum locustis rubris montanum. *Baub.*
pin. 10.

Anglis,

TRIANDRIA DIGYNIA.

Anglis, Melic-grafs.

Habitat in *sylvis subhumidis frequens*. 4. VI. VII.

*** *Flores multiflori vagi*.

B R I Z A. 7.

Cal. bivalvis, multiflorus. *Spicula* disticha; valvulis cordatis.

media.

1. BRIZA spiculis ovatis, calyce flosculis brevior. *Fl. suec.* 86. *Sp. pl.* 70.
Gramen tremulum. *Bauh. hist.* II. 469. majus.
Bauh. pin. 2. *R. Syn.* 412.
Gramen tremulum f. phalaris media Anglica prima an secunda. *Park.* 1165.
Phalaris pratense. *Ger. Em.* 86.
Anglis, middle Quaking-grafs, Cow-quakes, and Ladies-hair.
Habitat in *pratis et pascuis*. 4. VI.

minor.

2. BRIZA spiculis triangulis, calyce flosculis longior. *Sp. pl.* 70.
Gramen tremulum minus, panicula parva. *Bauh. pin.* 2. *Hist. Ox.* III. 208. t. 6. f. 47.
Gramen tremulum minus, panicula ampla, locustis parvis triangulis. *R. Syn.* 412.
Anglis, small Quaking-grafs.
Habitat in *pascuis*. In *Jersey*. Dr. Sherard. *R. Syn.* 4. VII.

P O A. 77.

Cal. bivalvis, multiflorus. *Spicula* ovata: valvulis ovato-acuminatis.

- aquatica.* 1. POA panicula diffusa, spiculis sexfloris linearibus. *Fl. suec.* 81. *Sp. pl.* 67.
Poa panicula diffusa, spiculis sexfloris linearibus muticis compressis. *Act. stockh.* 1741. p. 185.
Gramen paniculatum aquaticum latifolium. *Bauh. pin.* 3. *Hist. Ox.* III. 201. t. 6. f. 25.
Gramen palustre paniculatum altissimum. *Bauh. pin.* 3.

Gramen

Gramen aquaticum majus. *Ger. Em.* 6. 411.

Anglis, Reed Meadow-grafs.

Habitat *ad ripas fluviorum, et in paludibus.* 4. VII.
VIII.

2. POA panicula diffusa, spiculis subtrifloris basi pubescentibus, culmo erecto tereti. *Syst. Nat.* 874. *Sp. pl.* 67.

Gramen pratense paniculatum medium. *Bauh. pin.* 2. *R. Syn.* 409.

Gramen pratense minus. *Bauh. hist. II.* 542. *Park.* 1156. *Ger. Em.* 2.

Anglis, common Meadow-grafs.

Habitat *in pratis et pascuis ubique.* 4. VI.—VIII.

3. POA panicula diffusa, spiculis quinquefloris, glabris pratensis, culmo erecto tereti. *Fl. suec.* 82. *Sp. pl.* 67.

Gramen pratense paniculatum majus latiore folio, Poa Theophrasti. *Bauh. pin.* 2. *th.* 28. *R. Syn.* 409.

Gramen pratense. *Ger. Em.* 2. vulgatus. *Park.* 1156. *Anglis*, great Meadow-grafs.

Habitat *in pratis et pascuis prope Mary-le-bon.* 4. VI.
VII.

4. POA panicula coarctata secunda, culmo obliquo compresso. *Fl. suec.* 84. *Sp. pl.* 68.

Poa spiculis ovatis compressis muticis, panicula secunda contracta. *Act. stockh.* 1741. p. 184.

Gramen pratense medium culmo compresso, *Bud. H. S. R. Syn.* 409.

Gramen murorum, radice repente. *Bauh. pin.* 2, *pr.* 2.

Gramen paniculatum, radice repente, culmo compresso. *Vaill. parif.* 91. *t.* 18. *f.* 5.

Anglis, creeping Meadow-grafs.

Habitat *in locis siccis, testis, muris.* On the walls about Eltham, Mr. J. Sherard. *R. Syn. in pratis prope Mary-le-bon.* ①. VI.

- angustifolia.* 5. POA panicula diffusa spiculis quadrifloris pubescentibus, culmo erecto tereti. *Fl. suec.* 81. *Sp. pl.* 67.
 Poa spiculis ovato-oblongis, foliis subulatis. *Fl. lapp.* 52.
 Gramen pratense paniculatum majus angustiore folio.
Bauh. pin. 2. *pr.* 5. *Hist. Ox.* III. 201. *t.* 5. *f.* 19.
Anglis, narrow leaved Meadow-grafs.
 Habitat in sylvis et sepibus. ☉. VII.
- bulbosa.* 6. POA panicula secunda patentiuscula, spiculis quadrifloris. *Sp. pl.* 70.
 Poa culmorum basi tuberosa. *Guet. stamp.* I. 168.
Dalib. paris. 29.
 Gramen arvense, panicula crispa. *Bauh. pin.* 3. *Barr. ic.* 703.
 Gramen arvense, panicula crispa longiore nostras.
Pluk. alm. 176.
Anglis, bulbose Meadow-grafs.
 Habitat in pratis et pascuis; prope Clapham in comitatu Surrienfi. 4. VI.
- setacea.* 7. POA panicula diffusa, spiculis trifloris pubescentibus, foliis radicalibus setaceis.
Anglis, Hair-leaved Meadow-grafs.
 Habitat in pascuis siccis prope Mary-le-bon. ☉. V. VI.
- annua.* 8. POA panicula diffusa angulis rectis, spiculis obtusis culmo obliquo compresso. *Fl. suec.* 85. *Sp. pl.* 68. *Dalib. paris.* 28.
 Gramen pratense minus seu vulgatissimum. *R. Syn.* 408.
 Gramen pratense minimum album et rubrum. *Park.* 1156. *Ger. Em.* 3.
 Gramen pratense paniculatum minus. *Bauh. pin.* 3. *Th.* 31.
Anglis, annual Meadow-grafs, or Suffolk-grafs.
 Habitat in pratis et pascuis ubique. ☉. IV.—IX.
- nemoralis.* 9. POA panicula attenuata, spiculis subifloris mucronatis scabris, culmo incurvo. *Sp. pl.* 69.
 Gramen

Gramen sylvaticum, panicula tenui e raris locustis composita. *Dill. lat. giff. 57.*

Gramen paniculatum angustifolium alpinum, locustis rarioribus et angustioribus non aristatis. *Sch. agr. 164. t. 2. f.*

Anglis, Wood Meadow-grafs.

Habitat in *sylvis et umbrosis*. ☉. VI.

10. POA panicula subspicata, spiculis secundis coarctata-*maritima*,
tis, foliis convolutis.

Gramen paniculatum maritimum vulgatissimum. *R.*

Syn. 409.

β Gramen caninum maritimum paniculatum. *R. Syn. 410.*

Anglis, Sea Meadow-grafs.

Habitat in *littoribus maritimis ubique*. 4. VI. VII.

11. POA panicula lanceolata subramosa, paniculis alter-*rigida*,
nis secundis.

Poa panicula lanceolata subramosa, floribus alternis secundis. *Roy. lugdb. 62. Guett. stamp. 62.*

Gramen loliaceum murorum duriusculum spica erecta rigida. *Hist. Ox. III. 182. t. 2. f. 9.*

Gramen panicula multiplici. *Bauh. pin. 3. pr. 6. Park. 1157.*

Gramen exile duriusculum in muris et aridis proveniens. *R. Syn. 410.*

Anglis, Hard Meadow-grafs.

Habitat in *muris, tectis, et arenosis frequens*. ☉. VII.

12. POA spicata, spiculis alternis sessilibus, subsexfloris. *loliacea*,

Gramen pumilum Loliaceo simile. *R. Syn. 395.*

Gramen exile duriusculum maritimum foliolis circumvolutis, veluti juncis brevibus. *Pluk. phyt. t. 33. f. 7.*

Anglis, spiked Meadow-grafs.

Habitat in *arenosis maritimis frequens*. 4. V. VI.

TRIANDRIA DIGYNIA.

FESTUCA. 82.

Cal. bivalvis, *Spicula* oblonga, teretiuscula, glumis acuminatis.

* *Panicula secunda.*

ovina.

1. FESTUCA panicula secunda coarctata aristata, culmo tetragono nudiusculo, foliis setaceis. *Fl. suec.* 91. *Sp. pl.* 73.

Poa spiculis ovato angustis aristato-acuminatis. *Fl. lapp.*

Gramen capillaceum, locustellis pennatis non aristatis. *Pluk. Ph. t.* 34. *f.* 2. *R. Syn.* 410.

Gramen loliaceum, foliis brevibus junceis, minus. *Hist. Ox. III.* 182. *t.* 3. *f.* 13.

Gramen foliis junceis brevibus majus, radice nigra. *Bauh. pin.* 5. *Sch. agr.* 276.

vivipora.

- β Festuca spiculis viviparis. *Fl. suec.* 913.

Gramen sparteum montanum, spica foliacea graminea majus et minus. *R. Syn.* 410. *t.* 22. *f.* 1.

Anglis, Sheep's Fescue-grass.

Habitat in pascuis siccioribus, præsertim glareosis; β in montibus Eboracensibus, Westmorlandicis, Cambro-britanicis, frequens. 4. VI. VII.

duriuscula

2. FESTUCA panicula nutante inferne ramosa, spicis adscendentibus hispidis, foliis setaceis. *Roy. lugdb.* 68. *Sp. pl.* 74.

Gramen pratense, panicula duriore laxa unam partem spectante. *R. Syn.* 413. *t.* 19. *f.* 1. *Sch. agr.* 285.

Gramen tenue duriusculum et pene junceum. *Bauh. hist. II.* 463.

Anglis, hard Fescue-grass.

Habitat in pratis et pascuis siccioribus. 4. VI.

rubra.

3. FESTUCA panicula secunda scabra spiculis sexfloris aristatis; flosculo ultimo mutico, culmo semitereti. *Fl. suec.* 92. *Sp. pl.* 73.

Gramen alpinum pratense, panicula duriore laxa spadicæa, locustis majoribus. *Sch. agr.* 287.

Anglis, Purple Fescue-grass.

Habitat in pascuis siccis. 4. VI.

4. FESTUCA

4. FESTUCA panicula secunda, spiculis erectis, calycis altera valvula integra, altera acuminata. *Sp. pl.* 75. *bromoides.*
Festuca spicis erectis ad unum latus, palea altera calycina minima, altera acuminata. *Roy. lugdb.* 68.
Gramen paniculatum bromoides minus, paniculis aristatis unam partem spectantibus. *R. Syn.* 415.
hist. 1287. *Pluk. phyt. t.* 33. *f.* 10.
Anglis, barren Fescue-grafs.
Habitat in *pascuis arenosis*. ☉. V. VI.

5. FESTUCA panicula spicata nutante, calycibus minutissimis muticis, floribus scabris: aristis longis; *myuros.*
Syst. Nat. 876. *Sp. pl.* 74.
Gramen murorum spica longissima. *Ger. Em.* 31.
Hist. Ox. III. 215. *t.* 7. *f.* 43. *R. Syn.* 415.
Gramen spica nutante longissima. *Park.* 1162.
Anglis, Wall Fescue-grafs.
Habitat in *muris et locis agrorum sterilioribus et siccioribus*. ☉. VI.

6. FESTUCA panicula erecta, spiculis linearibus muticis, foliis planis. *pratensis.*
Gramen paniculatum elatius, spicis longis muticis et squamosis. *R. Syn.* 411.
Anglis, Meadow Fescue-grafs.
Habitat in *pratis et pascuis*. ☿. VI. VII.

** *Panicula æquali.*

7. FESTUCA panicula secunda ramosa erecta, spiculis *elatior*, subaristatis; exterioribus teretibus. *Fl. suec.* 94.
Sp. pl. 75.
Gramen loliaceum, spica divisa, pratense majus. *Hist. Ox.* III. 184. *t.* 2. *f.* 15.
Gramen paniculatum nemorosum, latiore folio glabrum, panicula nutante non aristata. *R. Syn.* 411.
β Gramen arundinaceum aquaticum panicula avenacea.
R. Syn. 411.
Anglis, Tall Fescue-grafs.
Habitat in *pratis et pascuis sed rarius*. ☿. VII.

- decumbens* 8. FESTUCA panicula erecta, spiculis subovatis muticis, calyce flosculis majore, caule decumbente. *Fl. suec.* 93. *Sp. pl.* 75.
 Gramen avenaceum parvum procumbens, paniculis non aristatis. *R. Syn.* 2d. ed. 262. 3d. ed. 408.
Pluk. ph. t. 34. f. 1.
 Gramen triticeum palustre humilius, spica mutica brevior. *Hist. Ox.* III. 177. e. 1. f. 6.
Anglis, small Fescue-grafs.
 Habitat in pascuis sterilibus subhumidis. 4. VIII.
- fluitans.* 9. FESTUCA panicula ramosa erecta, spiculis subsessilibus teretibus muticis. *Fl. suec.* 95. *Sp. pl.* 75.
 Gramen aquaticum cum longissima panicula. *Bauh. hist.* II. 490. *R. Syn.* 412.
 Gramen fluviatile. *Ger. Em.* 14. *Park.* 1275.
 Gramen aquaticum fluitans, multiplici spica. *Bauh. pin.* 2.
Anglis, Flote Fescue-grafs.
 Habitat in fossis et aquosis ubique. 4. VI. VII.
- foliacea.* 10. FESTUCA spicata, spiculis alternis sessilibus compressis muticis.
Anglis, spiked Fescue-grafs.
 Habitat ad vias et in pascuis sed rarius; prope Vauxhall. 4. VI.
- sylvatica.* 11. FESTUCA spicata, spiculis alternis subdistichis sessilibus aristatis.
 Festuca culmo alternatim spicato, spicis teretibus. *Fl. lapp.* 29. *Roy. lugdb.* 67.
 Festuca graminea nemoratis latifolia mollis. *Bauh. pin.* 10.
 Gramen avenaceum dumetorum spicatum. *R. Syn.* 394.
Anglis, Wood Fescue-grafs.
 Habitat in sylvis et sepius frequens. 4. VII.

B R O M U S. 83.

Cal. bivalvis. *Spicula* oblonga, teres, disticha; aristata infra apicem.

1. BROMUS panicula patente, spiculis ovatis: aristis *secalinus*.
rectis. *Fl. suec.* 96. *Sp. pl.* 76.

Festuca culmo paniculata, spicis compresso-ovatis.
Fl. lapp. 28.

- β Bromus panicula erecta coarctata. *Sp. pl.* 78. hordea-
Festuca avenacea hirsuta, paniculis minus sparsis. *R. ceus*.
Syn. 413.

Gramen avenaceum pratense, panicula squamata et
villosa. *Hist. Ox. III.* 213. *t. 7. f.* 18.

- γ *Festuca* avenacea spicis habitioribus glumis glabris. *R.*
Syn. 414.

Gramen avenaceum segetale majus gluma turgidiore.
Hist. Ox. III. 212. *t. 7. f.* 16.

- δ *Gramen* avenaceum pratense, gluma tenuiore glabra.
Hist. Ox. III. 213. *t. 7. f.* 19.

- ε *Festuca* avenacea spicis strigosioribus, e glumis glabris
compactis. *R. Syn.* 414.
Anglis, Field Brome-grass.

Habitat in pratis et pascuis. γ. δ. in arvis. ☉. V. VI.

2. BROMUS panicula nutante, spiculis ovato oblongis. *arvensis*.
Fl. suec. 97. *Sp. pl.* 77.

Bromus culmo paniculato, spicis compressis. *Fl.*
lapp. 27.

Bromus herba. *Dod. p.* 540.

Festuca graminea effusa juba. *Bauh. pin.* 192.

Festuca elatior paniculis minus sparsis, locustis oblongis
strigosis, aristis purpureis splendentibus. *R.*
Syn. 414.

Anglis, Corn Brome-grass.

Habitat inter segetes prope Southamptoniam D. Sherard
invenit. *R. Syn. prope* Gravesend copiose. ♀. VII.

3. BROMUS panicula simplici erecta, spiculis ovato-erectus.
oblongis, foliis involutis pilosis.

Anglis, upright Brome-grass

Habitat in cretaceis circa Rochester, Dartford, and
Gravesend, in Cantio. 4. VI.

- ciliatus.* 4. BROMUS panicula nutante, foliis utrinque vaginif-
que subpilosis glumis ciliatis. *Sp. pl.* 76.
Festuca avenacea sterilis spicis erectis. *R. Syn.* 413.
Pluk. Pb. t. 299. *f.* 2.
Festuca avenacea sterilis, pediculis brevioribus et spicis
erectis. *Hist. Ox. III.* 212. *t.* 7. *f.* 13.
Anglis, Wall Brome-grafs.
Habitat in tectis muris et pascuis siccissimis. ☉. V.

- sterilis.* 5. BROMUS panicula patula, spiculis oblongis distichis,
glumis subulato aristatis. *Sp. pl.* 77.
Festuca avenacea sterilis elatior seu Bromos Dioscori-
dis. *Bauh. pin.* 9. *Th.* 146. *Hist. Ox. III.* 212.
t. 7. *f.* 11. *R. Syn.* 412.
Bromos herba five avena sterilis. *Park.* 1147.
Bromos sterilis. *Ger. Em.* 16.
Anglis, barren Brome-grafs.
Habitat in sylvis, sepibus et ad margines agrorum fre-
quens. ☉. VI. VII.

- giganteus.* 6. BROMUS panicula nutante, spiculis quadrifloris a-
ristis brevibus. *Fl. succ.* 99. *Sp. pl.* 77. *It. scan.*
394.
Gramen sylvaticum glabrum panicula recurva. *Vaill.*
parif. 93. *t.* 18. *f.* 3.
Gramen avenaceum glabrum, panicula e spicis raris
strigosis composita, aristis tenuissimis. *R. Syn.* 415.
Gramen bromoides aquaticum latifolium, panicula
sparsa tenuissime aristata. *Sch. agr.* 264. *t.* 5.
f. 17.
Anglis, tall Brome-grafs.
Habitat in sylvis et sepibus subhumidis. 4. VIII.

- ramosus.* 7. BROMUS panicula ramosa nutante scabra, spiculis
linearibus decemfloris arista longioribus, foliis sca-
bris.
Festuca graminea nemoralis latifolia mollis. *Bauh.*
pin. 9. *th.* 144.

Gramen

TRIANDRIA DYGINIA.

41

Gramen avenaceum dumetorum panicula sparfa. *R. Syn.* 415.

Gramen avenaceum dumetorum, juba longiore spica divisa. *Hist. Ox. III.* 213. *t.* 7. *f.* 27.

Anglis, Wood Brome-grafs.

Habitat in sylvis et sepibus frequens. ☉. VII. VIII.

8. BROMUS culmo indiviso, spicis alternis subseffilibus *pinnatus*. teretibus, flosculis arista longioribus.

Bromus culmo indiviso, spicis alternis subseffilibus teretibus. *Fl. suec.* 100. *Sp. pl.* 78.

Gramen spica brizæ majus. *Baub. pin.* 9. *pr.* 19. *th.* 133. *R. Syn.* 392.

Anglis, spiked Brome-grafs.

Habitat in pratis siccis. In coppices and hedges about Oxford. On all the heaths and commons for twenty miles on this side York. *R. Syn.* Circa Dover, in Cantio frequens. 4. VI.

A V E N A. 85.

Cal. bivalvis, multiflorus; arista dorsali contorta.

1. AVENA calycibus trifloris, receptaculo calycem ex-nuda. cedente petalis dorso aristatis. *Amoen. acad.* III. 401.

Avena nuda. *Baub. pin.* 23. *hist. II.* 433. *Ger. Em.* 77. *Park.* 1134. *R. Syn.* 389.

Anglis, naked Oats or Pilecorn.

Habitat in arvis sed rarius. In extrema Cornubia prope Belerium, promontorium copiose feritur. *R. Syn.* in Staffordshire, Dr. Plot. ☉. VII.

2. AVENA panicula patente, calycibus trifloris flosculis *fatua*. basi pilosis. *Fl. suec.* 101. *Sp. pl.* 80.

Avena sylvestris pilosa, aristis recurvis. *Hist. Ox. III.* 209. *t.* 7. *f.* 5.

Festuca utriculis lanugine flavescentibus. *Baub. pin.* 10.

Ægilops quibusdam aristis recurvis, s. Avena pilosa. *Baub. hist. II.* 433. *R. Syn.* 389.

Ægilops bromoides. *Ger. Em.* 77. *Belgarum. Park.* 1148.

Anglis,

Anglis, bearded Oat-grafs.

Habitat in arvis, prope Stockwell in Comitatu Surri-
ensi. 2. VIII.

pratensis. 3. AVENA calycibus quinquefloris, panicula spicata,
foliis involutis.

Avena calycibus quinquefloris, panicula spicata. *Fl.*
suec. 104. *Sp. pl.* 80.

Gramen avenaceum montanum spica simplici, aristis
recurvis. *R. Syn.* 405. t. 21. f. 1.

Anglis, Meadow Oat-grafs.

Habitat in ericetis, pratis montosis præsertim solo creta-
ceo. 4. VII.

pubescens. 4. AVENA calycibus subtrifloris basi pilosis, foliis planis
pubescentibus.

Gramen avenaceum hirsutum, f. glabrum panicula
purpureo-argentea splendente. *R. hist.* 1909. *Syn.*
406. t. 21. f. 2.

Anglis, rough Oat-grafs.

Habitat in pratis et pascuis siccioribus. *Supra Ban-*
stead Downs copiose. 4. VI.

elatior. 5. AVENA calycibus bifloris, flosculo hermaphrodito
mutico, masculo aristato. *Fl. lapp.* 98. *suec.* 102.
Sp. pl. 79.

Gramen nodosum avenacea panicula. *Baub. pin.* 2.
pr. 3. *R. Syn.* 406.

Gramen caninum. nodosum. *Ger. Em.* 23. f. bulbosum
vulgare. *Park.* 1175.

β Gramen avenaceum elatius, juba longa splendente. *R.*
meth. em. 179.

Anglis, Tall Oat-grafs.

Habitat in pratis et pascuis. 4. VII. VIII.

flavescens. 6. AVENA panicula laxa, calycibus trifloris brevibus,
flosculis omnibus aristatis. *Roy. lugdb.* 66. *Sp. pl.*
80. *Fl. suec.* 103.

Gramen avenaceum pratense elatius, panicula flavesc-
cente, locustis parvis. *R. Syn.* 407.

Gramen avenaceum, spica sparsa flavescente, locustis
parvis. *Raii. hist. Ox.* III. 215. t. 7. f. 42.

Anglis,

Anglis, yellow Oat-grafs.

Habitat in pratis et pascuis ubique. 4. VII.

A R U N D O. 87.

Cal. bivalvis. Flosculi congesti, lana cincti.

1. ARUNDO calycibus quinquefloris, panicula laxa. *phragmites*.
Roy. lugdb. 66. *Gron. virg.* 137. *Sp. pl.* 81. *Fl. su-*
ec. 105.

Arundo vulgaris five phragmites Dioscoridis. *Bauh.*
pin. 17.

Arundo vallatoria. *Ger. Em.* 36. *R. Syn.* 401.

Harundo vulgaris five vallatoria. *Park.* 1208.

Anglis, common Reed-grafs.

Habitat in fluviis, Lacubus, fossis, ubique. 4. VII.

2. ARUNDO calycibus unifloris, culmo ramoso. *Fl. Calama-*
lapp. 42. *suec.* 107. *Sp. pl.* 82. *grostis.*

Gramen arundinaceum panicula molli spadicea majus.

Bauh. pin. 7. *Sch. agr. pr. t.* 5. *R. Syn.* 401.

Gramen tomentorium arundinaceum. *Ger. Em.* 9.

Calamagrostis f. Gramen tomentosum. *Park.* 1182.

Anglis, branched Reed-grafs.

Habitat in sylvis et sepibus subhumidis. *Inter New-*
ington & Hornsey. 4. VI. VII.

3. ARUNDO calycibus unifloris, panicula erecta, foliis epigejos.
subtus glabris. *Sp. pl.* 81. *Fl. suec.* 106.

Arundo panicula coarctata erecta integra, foliis inferne
glabris. *Fl. suec. ed. 1.* 101.

Gramen arundinaceum paniculatum montanum, pa-
nicula spadiceo viridi, semine papposo. *Sch. agr.*
124.

Calamagrostis minor glumis rufis et viridibus. *Pet.*
conc. gr. n. 69.

Anglis, small Reed-grafs.

Habitat circa Oundle in comitatu Leicesterienfi. *R.*
Syn. 4. VII.

4. ARUNDO calycibus unifloris, foliis involutis mu- *arenaria.*
cronato-pungentibus. *Sp. pl.* 82. *Fl. suec.* 108.

Gramen

TRIANDRIA DIGYNIA.

Gramen sparteum spicatum foliis mucronatis longioribus vel spica secalina. *Bauh. pin. 5. th. 67. R. Syn. 393.*

Spartum marinum nostras. *Park. 1198.*

Spartum anglicanum. *Ger. Em. 42.*

Anglis, Sea Reed-grafs.

Habitat in littoribus maritimis. 4. VI. VII.

*** Flores multiflori, infidentes Receptaculi elongati dentibus.

L O L I U M. 90.

Involucrum monophyllum, fixum. Cal. o. Spicula multiflora.

perenne. 1. LOLIUM spicis muticis, radice perenni. *Hort. cliff. 24. Sp. pl. 83.*

Gramen loliaceum angustiore folio et spica. *Bauh. pin. 9. R. Syn. 395.*

Lolium rubrum. *Ger. Em. 78. f. Phoenix Park. 1145.*

β Gramen loliaceum paniculatum. *R. Syn. 395.*

Anglis, perennial Darnel-grafs.

Habitat ad vias et pratis ficcioribus ubique. 4. VI.

temulentum. 2. LOLIUM spica aristata, radice annua. *Syst. nat. 879. Sp. pl. 83.*

Lolium album. *Ger. Em. 78. Park. 1146. R. Syn. 396.*

Gramen loliaceum spica longiore. *Bauh. pin. 9. th. 121.*

Anglis, annual Darnel-grafs.

Habitat in arvis. ☉. VII. VIII.

E L Y M U S. 91.

Involucrum diphyllum. Spiculæ geminæ:

arenarius. 1. ELYMUS foliis mucronato-pungentibus. *It. scan. 336. Sp. pl. 83.*

Secale spiculis geminatis. *Fl. suec. 1, 106.*

Gramen caninum maritimum spica triticea nostras. *R. Syn. 390. Sch. agr. 6.*

Gra-

Gramen caninum maritimum spicatum. *Bauh. pin.* 1.

Gramen caninum geniculatum maritimum spicatum.

Park. 1277.

Anglis, Sea Lyme-grafs.

Habitat in littoribus maritimis frequens. 4. V. VI.

TRITICUM. 94.

Cal. bivalvis, subtriflorus, sessilis, obtusiusculus,
acutus.

1. TRITICUM calycibus subulatis trifloris acuminatis. *repens.*
Sp. pl. 86.

Gramen spica triticea repens vulgare, caninum dictum. *R. Syn.* 390.

Gramen caninum. *Ger. Em.* 23. vulgatius. *Park.* 1173. *Hist. Ox.* III. 178. t. 1. f. 8.

Gramen caninum arvense f. Gramen Dioscoridis.
Bauh. pin. 1. th. 7.

Anglis, common Wheat-grafs, Dog's-grafs, Quick-grafs, or Couch-grafs.

Habitat in agris ubique. 4. VI.—VIII.

Pharmac. Graminis canini Radix.

2. TRITICUM calycibus subulatis quadrifloris aristatis. *caninum.*
Sp. pl. 86.

Elymus spiculis rectis involucro destitutis; infimis geminis. *Fl. suec.* 112. *Syst. Nat.* 879.

Gramen caninum aristatum radice non repente, sylvaticum. *R. Syn.* 390.

Gramen caninum non repens elatius, spica aristata.
Hist. Ox. III. 177. t. 1. f. 2.

Anglis, bearded Wheat-grafs.

Habitat in sylvis et sepibus. In Stokenchurch woods plentifully; circa Croydon, D. Doody; near Settle in Yorkshire, Mr. J. Sherard. *R. Syn. in via* inter Greenwich & Woolwich. 4. VI. VII.

3. TRITICUM calycibus truncatis quinquefloris, foliis junceum involutis. *Cent. pl.* 13. *Syst. Nat.* 880.

Gramen maritimum, spica loliacea foliis pungentibus nostras. *Pluk. Ph.* t. 33. f. 4. a.

Gramen

TRIANDRIA DIGYNIA.

- Gramen phoenicoides foliis convolutis, junceis ac pungentibus alterum. *Bauh. hist. II. 478.*
- β Gramen loliaceum maritimum supinum spica crassiore. *Tourn. 516. R. Syn. 391.*
- γ Gramen caninum maritimum spica foliacea. *Bauh. pin. 2.*
Anglis, Sea Wheat-grafs.
Habitat in littoribus maritimis passim. 4. VI. VII.

HORDEUM. 93.

Involucrum hexaphyllum, triflorum.

- murinum.* 1. HORDEUM flosculis lateralibus masculis aristatis. *Sp. pl. 85.*
Hordeum spurium vulgare. Park. 1147.
 Gramen secalinum et secale sylvestre. *Ger. Em. 73.*
R. Syn. 391.
 Gramen hordeaceum minus et vulgare. *Bauh. pin. 9.*
- β Gramen secalinum. *Ger. Em. 29.*
- γ Gramen secalinum palustre et maritimum. *R. Syn. 392.*
Anglis, Wall Barley-grafs, or Way-bennet.
Habitat ad vias et muros, β in pratis humidis, γ in pratis maritimis. ⑤. IV. ad VIII.

SECALE. 92.

Involucrum diphyllum, biflorum.

- villosum.* 1. SECALE glumaram ciliis villosis, squamis calycinis cuneiformibus. *Sp. pl. 84.*
 Gramen secalinum majus sylvaticum. *Hist. Ox. III. 180. R. Syn. 393.*
 Gramen hordeaceum montanum spica strigosiori brevius aristata. *Sch. agr. pr. t. 1.*
Anglis, Wood Rye-grafs.
Habitat in sylvis et sepibus. In Stokenchurch woods in Oxfordshire, plentifully, Mr. Bobart. In the woods a mile west from Petersfield, Mer. pin. In the high woods by Hambleton, in the road from Henley

TRIANDRIA DIGYNIA.

47

ley to Great Marlborough, Mr. J. Sherard and Mr. Rand. *R. Syn.* ☉. VII.

CYNOSURUS. 81.

Cal. bivalvis, multiflorus; *Recept.* unilaterale.

1. CYNOSURUS bracteis pinnatifidis. *Sp. pl.* 72. *Fl. cristatus.* *suec.* 88.

Gramen cristatum. *Baub. hist.* II. 468. *Ger. Em.* 29. *R. Syn.* 398.

Gramen cristatum anglicum. *Park.* 1159.

Gramen pratense cristatum, f. Gramen spica cristata læve. *Baub. pin.* 3. *th.* 43.

- β Gramen cristatum quadratum f. quatuor cristatum glumarum versibus. *R. Syn.* 399.

Anglis, crested Dog-tail-grass.

Habitat in pratis et pascuis. 4. VIII.

2. CYNOSURUS bracteis pinnato-paleaceis aristatis. *echinatus.* *Roy. lugdb.* 64. *Sp. pl.* 72.

Gramen alopecuroides spica aspera. *Baub. pin.* 4.

Gramen alopecuroides spica alpera brevi. *Park.* 1168. *R. Syn.* 397.

Anglis, rough Dog tail grass.

Habitat in arenosis. Common on all the sandy grounds in Jersey; Mr. J. Sherard. ☉. VII.

3. CYNOSURUS bracteis integris. *Sp. pl.* 72. *Fl. cœruleus.* *suec.* 89.

Gramen glumis variis. *Baub. pin.* 10. *pr.* 21.

Gramen parvum montanum spica crassiore purpureo-cœrulea brevi. *R. Syn.* 399.

Anglis, blue Dog-tail-grass.

Habitat in pratis montosis. In prate Helf-fellnab, prope Kendal, et alibi in Westmorlandia. 4. VII.

4. CYNOSURUS panicula subspicata, flosculis simplicibus biaristatis. *Sp. pl.* 73. *Fl. suec.* 90.

Panicum floribus conglomerato-spicatis lævibus arista dimidio brevioribus. *Fl. suec. ed. l.* 54.

Alopecuros maxima anglica, *Park.* 1166. *R. Syn.* 396.

Alope-

TRIANDRIA DIGYNIA.

Alopecuros altera maxima anglica paludosa. Ger. Em. 88.

Gramen alopecuroides anglobritannum maximum. Bauh. pin. 4.

Anglis, bearded Dog-tail-grass.

Habitat in riguis herbidis comitatus Southamptoniæ proxime salinis et antiquis ædes Drayton vocatas, eis mare, duobus miliaribus Anglicis a Portsmouth ex adverso vectis insulæ plurima; itemque in udis fossis lacustribus Essexiensis comitatus juxta Thamelis amoenissima fluente. At Purfleet in Essex, over against the mill towards Raynam, on the other side of the great ditch; Mr. Newton. R. Syn. 4. VII.

TRIANDRIA.

MONTIA. 96.

Cal. diphyllus. Cor. monopetala, irregularis. Caps. unilocularis, trivalvis.

- fontana.* 1. MONTIA. *Fl. lapp. 30. suec. 115. Sp. pl. 87.*
Montia aquatica minor. Mich. gen. 18. t. 13. f. 2.
Cameraria arvensis minor. Dill. giff. 46.
Alfene parva palustris tricoccus, portulacæ aquaticæ similis. R. Syn. 252.
Portulaca arvensis. Bauh. pin. 288.
Anglis, Water Chickweed.
Habitat ad fontes et scaturagines. ☉. IV.

Classis

*Classis IV.*TETRANDRIA
MONOGYNIA.

* *Flores monopetali, monospermi, superi. Aggregati.*

DIPSACUS. 107.

*Cal. communis polyphyllus ; proprius superus. Re-
cept. paleaceum.*

1. DIPSACUS foliis sessilibus ferratis, aristis fructus re-*fullonum*,
curvis.

Dipsacus foliis sessilibus ferratis. *Sp. pl.* 97.

Dipsacus fativus. *J. Bauh. hist. III.* 73. *Bauh. pin.*
385. *Ger. Em.* 1167. *Park.* 983. *R. Syn.* 192.

Carduus fullonum. *Lob. ic.* 17.

Anglis, manured Teasel.

Habitat *ad sepes.* ♂. VII.

Pharmac. Dipsaci fativi *Radix.*

2. DIPSACUS foliis sessilibus ferratis, aristis fructus e-*sylvestris*,
rectis.

Dipsacus sylvestris aut Virga pastoris major. *Bauh.*
pin. 385.

Dipsacus sylvestris. *Ger. Em.* 1167. *Park.* 984. *R.*
Syn. 192.

Anglis, wild Teasel.

Habitat *in incultis frequens.* ♂. VII.

3. DIPSACUS foliis petiolatis appendiculatis. *Sp. pl.* 97. *pilosus*.
Dipsacus sylvestris capitulo minore, vel Virga pastoris
minor. *Bauh. pin.* 385.

E

Dip-

TETRANDRIA MONOGYNIA.

Dipsacus minor seu *virga pastoris*. *Ger. Em.* 1168.
R. Syn. 192.

Virga Pastoris. *Park.* 984.

Anglis, small wild Teasel.

Habitat in aquosis et ad sepes. 4. VIII.

SCABIOSA. 108.

Calyx communis polyphyllus ; *proprius* duplex superus. *Receptaculum* paleaceum.

* *Corollulis quadrifidis.*

fuccifa. 1. SCABIOSA corollulis quadrifidis æqualibus, caule simpliciter, ramis approximatis, foliis lanceolato-ovatis.

Fl. juec. 119. *mat. med.* 41. *Sp. pl.* 98.

Scabiosa radice fuccifa, flore globofo. *R. Syn.* 191.

Succisa glabra et hirsuta. *Bauh. pin.* 269.

Morsus Diaboli. *Ger. Em.* 726.

Morsus Diaboli vulgaris, flore purpureo. *Park.* 491.

Anglis, Devil's-bit.

Habitat in pratis et pascuis. 4. VI.—VIII.

arvensis. 2. SCABIOSA corollulis quadrifidis radiantibus, caule hispido foliis pinnatifidis hispidis. *Syst. Nat.* 889.
Sp. pl. 99.

Scabiosa caule et foliis hirsutis, imis ovatis, superioribus semipinnatis. *Hall. helv.* 669.

Scabiosa pratensis hirsuta. *Bauh. pin.* 269.

Scabiosa major vulgaris. *Ger. Em.* 719.

Scabiosa vulgaris pratensis. *Park.* 484.

Scabiosa major communior folio laciniato. *Bauh. hist.* III. 2.

Anglis, common field Scabious.

Habitat in pratis et arvis. 4. VIII.

Pharmacop. *Scabiosæ Herba.*

** *Corollulis quinquefidis.*

columbaria. 3. SCABIOSA corollulis quinquefidis radiantibus, foliis radicalibus ovatis crenatis, caulinis pinnatis setaceis.
Sp. pl. 99.

Scabiosa capitulo globofo major et minor. *Bauh. pin.* 270.

Scabiosa

TETRANDRIA MONOGYNIA.

51

Scabiosa minor campestris. *Park.* 487.

Scabiosa minor sive Columbaria. *Ger. Em.* 719.

Anglis, the lesser field Scabious.

Habitat in pascuis montosis, et siccioribus; supra Duper's hill prope Croydon. 24. VI. VII.

** Flores monopetali, monocarpii, inferi.

CENTUNCULUS. 135.

Cal. quadrifidus. *Cor.* quadrifida patens. *Stam.* brevica. *Capf.* unilocularis, circumscissa.

1. CENTUNCULUS. *Cat. giff.* 161. *app.* III. t. 5. *minimus.*

R. Syn. 1. *Fl. suec.* 136. *Sp. pl.* 116.

Anagallis, paludosa minima, foliis subrotundis alternis.

Vaill. Paris. 12. t. 4. f. 2.

Alfina palustris minima, flosculis albis, fructu coriandri exiguo. *Mentz. pug.* t. 4. f. 5.

Anglis, bastard Pimpernel.

Habitat in agris arenosis subhumidis. On the low marshy ground near the paper-mills on Hounslow-heath; D.

Watson. Blackf. spec. bot. in pascuo ante vicum

Chiffelhurst loco subudo. R. Syn. 6. VI.

PLANTAGO 133.

Cal. quadrifidus. *Cor.* quadrifida: limbo reflexo.

Stam. longissima. *Capf.* bilocularis circumscissa.

1. PLANTAGO foliis ovatis glabris, scapo tereti, spica majoris flosculis imbricatis. *Sp. pl.* 112. *Fl. suec.* 129.

Plantago latifolia vulgaris. Park. 493.

Plantago latifolia. Ger. Em. 419. sinuata. *Baub. pin.* 189.

2. *Plantago latifolia glabra minor. Baub. pin.* 189.

Anglis, great Plantain, or Waybread.

Habitat in pratis et ad vias ubique. 6. VI. VII.

Pharmac. Plantaginis Herba, Semina.

2. PLANTAGO foliis ovato-lanceolatis pubescentibus, media spica cylindrica, scapo tereti. *Sp. pl.* 113. *Hort. cliff.* 36. *Fl. suec.* 130.

Plantago major incana. Park. 493. *R. Syn.* 314.

Plantago incana. Ger. Em. 419.

TETRANDRIA MONOGYNIA.

Plantago latifolia incana. *Baub. pin.* 189.

Anglis, hoary Plantain.

Habitat in *pascuis glareosis et ad vias.* 4. VII. VIII.

lanceolata. 3. *PLANTAGO* foliis lanceolatis, spica subovato nuda, scapo angulato. *Fl. sue.* 131. *Sp. pl.* 113.

Plantago lanceolata. *Baub. hist.* III. 505.

Plantago angustifolia major. *Baub. pin.* 189.

Plantago quinquenervia. *Ger. Em.* 422. *major. Park.*

495.

Anglis, Ribwort, or Ribwort-plantain.

Habitat in *pratis et pascuis ubique.* 4. VI. VIII.

kefingii. 4. *PLANTAGO* foliis linearibus subdentatis, scapo tereti, spica cylindrica, bracteis carinatis membranaceis.

Plantago foliis linearibus subdentatis, scapo tereti, spica ovata, bracteis carinatis membranaceis. *Læf. hisp.* 124. *Sp. pl.* 115. ?

Plantago marina. *Ger. Em.* 423. *R. Syn.* 315. vulgaris. *Park.* 498.

Anglis, Sea-plantain.

Habitat in *palustribus maritimis.* 4. VII.

maritima. 5. *PLANTAGO* foliis femicylindræis integerrimis basi lanatis, scapo tereti. *Sp. pl.* 114. *Fl. suec.* 133.

Eaten by the
Scandinavians.

Plantago scapo spicato, foliis linearibus subtus convexis. *Fl. lapp.* 63.

**Plantago maritima major tenuifolia.* *Tourn.* 127.

Coronopus, *Ger. Em.* 425. *maritima major. Baub. pin.* 190.

Anglis, narrow-leaved Plantain.

Habitat in *littoribus maritimis inter Newcastle & Morpeth in Northumbria.* 4. VI. VII.

Coronopus. 6. *PLANTAGO* foliis linearibus pinnato-dentatis, scapo tereti.

Plantago foliis linearibus dentatis, scapo tereti. *Sp. pl.* 115.

Plantago foliis linearibus pinnato-dentatis. *Hort. cliff.* 37.

Plantago foliis laciniatis, *Coronopus dicta.* *R. Syn.* 315. *Coronopus*

Coronopus vulgaris five *Cornu Cervinum*. *Park.*
502.

Coronopus sylvestris hirsutior. *Bauh. pin.* 190.

Cornu Cervinum. *Ger. Em.* 427.

3. *Plantago gramineo folio hirsuto minor, capitulo rotundo brevi*. *R. Syn.* 316.

Anglis, Buckshorn-plantain, Star of the Earth.

Habitat in glareosis. ☉. VII. VIII.

Pharmac. *Coronopi Herba*.

7. *PLANTAGO foliis lineari-lanceolatis basi lanatis, montana. spica subrotunda, scapo tereti.*

Plantago alpina angustifolia. *Bauh. hist. III.* 506. *R. Syn.* 315.

Anglis, Mountain-plantain.

Habitat in rupibus Trivyldaugh *supra lacum* Lhyn
Bochlyn Sancti Parisii; *D. Lhwyd.* *R. Syn.* 4.
VII.

8. *PLANTAGO foliis subulatis, scapo unifloro. Syst. uniflora. Nat.* 896. *Fl. suec.* 135. *Fl. lapp.* 64. *Sp. pl.* 115.

Plantago palustris gramineo folio, monanthos parisiensis. Tourn. 128. *R. Syn.* 316.

Plantago floribus femineis sessilibus ad exortum scapi uniflori maris. Juss. act. 1742. *p.* 131. *t.* 7.

Gramen junceum five Holosteum minimum palustre, capitulis quatuor longissimis staminibus donatis. Hist. Ox. III. 230. *t.* 9. *f.* 30. *Pluk. Ph. t.* 35.
f. 2.

Anglis, Grass-leaved Plantain.

Habitat in palustribus, arenosis. Ad Hoseley Lough in Northumberland; D. Richardson. R. Syn. in ericeto Hounsflejo prope Witton. 4. VII.

** *Flores monopetali, monocarpi superi.*

SANGUISORBA. 136.

Cal. diphyllus. Germen inter calycem corollamque.

1. *SANGUISORBA spicis ovatis. Hort. cliff.* 39. *Fl. officinalis. suec.* 137. *Sp. pl.* 116.

TETRANDRIA MONOGYNIA.

Sanguiforba major flore spadiceo. *Bauh. hist. III. 2.*
120. *R. Syn. 203.*

Pimpinella sanguiforba major. *Bauh. pin. 160.*

Pimpinella major vulgaris. *Park. 582.*

Pimpinella sylvestris. *Ger. Em. 1045.*

β Sanguiforba minor. *Bauh. hist. III. 2. 113.*
Anglis, Burnet.

Habitat in pratis et pascuis subhumidis, β in pascuis
montosis. 4.

Pharm. Pimpinellæ Herba.

* Flores monopetali, dicocci, superi. Stellatæ.

RUBIA. 119.

Cor. monopetala, campanulata. Baccæ duæ, monospermæ.

anglica. 1. RUBIA foliis quaternis. *Roy. lugdb. 254. Sp. pl.*
109.

Rubia quadrifolia asperima lucida peregrina. *Herm.*
lugdb. 523.

Rubia sylvestris. *Park. 274.*

Anglis, wild Madder.

Habitat in rupe S. Vincentii, prope Bristolium, & in
sepibus in Devoniam frequens. 4. VII. VIII.

SHERARDIA. 112.

Cor. monopetala infundibuliformis, Semina bina, tridentata.

arvensis. 1. SHERARDIA foliis omnibus verticillatis, floribus
terminalibus. *Sp. pl. 102. Fl. suec. 120.*

Sherardia. *Dill. gen. 96. Hort. cliff. 33.*

Rubeola arvensis repens cærulea. *Bauh. pin. 334. pr.*
145. *R. Syn. 225.*

Rubeola minor pratensis cærulea. *Park. 276.*

Anglis, little Field-madder.

Habitat in arvis. ☉. V. VI.

ASPERULA.

A S P E R U L A. 113.

Cor. monopetala, infundibuliformis. *Sem.* duo globosa.

1. ASPERULA foliis octonis lanceolatis, floribus fasciculatis pedunculatis. *Fl. suec.* 121. *Sp. pl.* 103.
Asperula *Ger. Em.* 1224. *R. Syn.* 224. odorata. *Park.* 563.
Asperula f. *Rubeola montana odorata. Bauh. pin.* 334. *Anglis, Woodroof.*
Habitat in sylvis et umbrosis. 4. V.
Pharm. *Asperulæ odoratæ Herba.*
2. ASPERULA foliis quaternis linearibus, superioribus *cynanchica* oppositis caule erecto, floribus quadrifidis. *Syst. Nat.* 891.
Asperula foliis inferioribus senis, intermediis quaternis subulatis triquetris, floribus quadrifidis. *Sp. pl.* 104.
Rubeola foliis quaternis, feminibus glabris, floribus umbellatis. *Hall. helv.* 458.
Rubeola vulgaris quadrifolia lævis, floribus purpurantibus. Tourn. 130. *R. Syn.* 225.
Rubia cynanchica. Bauh. pin. 333. *hist. III.* 723.
Synanchica. Lugd. 1185. *Ger. Em.* 1120.
Asperula repens Gesneri f. Saxifraga altera Cæsalpini. Park. 453.
Anglis, Squinancy-wort.
Habitat in montosis siccioribus, præsertim cretaceis. 4. VII.

G A L I U M. 117.

Cor. monopetala, plana. *Sem.* bina subrotunda.

* *Fructus glabro.*

1. GALIUM foliis octonis linearibus sulcatis, ramis floriferis brevibus. *Hort. cliff.* 34. *Mat. med.* 46.
Sp. pl. 107. *Fl. suec.* 123.
Gallium caule erecto, foliis plurimis verticillatis linearibus. *Fl. lapp.* 61.
Gallium luteum. Bauh. pin. 335. *Ger. Em.* 1126.
Park. 564. *R. Syn.* 224.
E 4 *Anglis,*

TETRANDRIA MONOGYNIA.

Anglis, yellow Ladies Bedstraw, or Cheese-rening.
*Habitat in pratis ad vltarum agrorumque margines, locis
 siccis frequens.* 4. VII. VIII.
Pharmacop. Gallii Herba.

- X *Mollugo*. 2. **GALIUM** foliis octonis ovato-linearibus subserratis
 patentissimis mucronatis, caule flaccido, ramis pa-
 tentibus. *Sp. pl.* 107.
Mollugo vulgarior. *Park.* 565. *R. Syn.* 223.
Rubia sylvestris lævis. *Bauh. pin.* 333.
Rubia sylvestris. *Ger. Em.* 1118.
Anglis, wild Madder, or great bastard Madder.
Habitat in sepibus et ericetis frequens. 4. VI. VII.
- montanum* 3. **GALIUM** foliis senis obovatis mucronatis glabris,
 caule prostrato, pedunculis ternatis.
Mollugo montana minor Gallio albo similis. *R. hist.*
I. 482. *Syn.* 224.
Gallium album minus. *Pet. H. Br. t.* 30. *f.* 6.
Anglis, mountain Ladies Bedstraw.
Habitat in montosis. 4. VII.
- uligino-
sum.* 4. **GALIUM** foliis senis lanceolatis retrorsum serrato-
 aculeatis mucronatis rigidis, corollis fructu majori-
 bus. *Fl. suec.* 127.
Aparine palustris minor Parisiensis, flore albo. *Tourn.*
114. Vaill. parif. 14. *Fl. lapf.* 58. *R. Syn.* 225.
Rubia quædam minor. *Bauh. hist.* III. 2. 716.
Anglis, marsh Goose-grass.
*Habitat in ericetis et pascuis humidioribus frequens, in
 ericeto Hampstediano prope Londinum copiose.* 4.
 VII. VIII.
- erectum.* 5. **GALIUM** foliis octonis glabris lanceolatis mucrona-
 tis, caule suberecto, panícula terminali, pedunculis
 dichotomis.
Anglis, small mountain bastard Madder.
Habitat in pascuis montosis humidiusculis. 4. VIII.
- perfillum.* 6. **GALIUM** foliis octonis hispidis linearibus acuminatis
 subimbricatis, pedunculis dichotomis. *Sp. pl.* 106.
Rubeola saxatilis. *Bauh. pin.* 334. *pr.* 145.
Anglis,

Anglis, the least Ladies Bedstraw.

Habitat in montibus prope Kendal, in comitatu Westmorlandico, 4. VIII.

7. GALIUM foliis quaternis obovatis inæqualibus, caulibus diffusis. *Fl. suec.* 126. *Sp. pl.* 105.

Gallium palustre album. *Baub. pin.* 335.

Gallium album. *Ger. Em.* 1126. *Baub. hist.* III.

721.

Molluginis vulgationis varietas minor. *Park.* 565.

R. Syn. 224.

Anglis, white Ladies Bedstraw.

Habitat ad ripas fluviorum et in pratis humidioribus.

4. VII.

8. GALIUM foliis senis lanceolatis carinatis scabris retrorsum aculeatis, geniculis simplicibus, fructibus glabris. *Sp. pl.* 106.

Aparine femine læviore. *R. Syn.* 225. *hist.* 484.

Aparine lævis. *Park.* 567.

Anglis, Goose-grafs with smoother seeds.

Habitat in arvis sed rarius. ☉. VI.

** *Fructu hispido.*

9. GALIUM foliis octonis lanceolatis carinatis scabris Aparine. ✕
retrorsum aculeatis, geniculis villosis, fructu hispido.

Sp. pl. 108.

Aparine vulgaris. *Baub. pin.* 334. *Park.* 567.

Aparine. *Ger. Em.* 1122. *Baub. hist.* III. 713. *R.*

Syn. 225.

Anglis, Cleavers, or Goose-grafs.

Habitat in cultis et ruderalis ubique. ☉. V. VIII.

Pharm. Aparines *Herba.*

10. GALIUM foliis verticillatis, pedunculis bifloris fructibus hispidis. *Sp. pl.* 108. *parisiense.*

Gallium parisiense tenuifolium, flore atro purpureo.

Tourn. 664.

Aparine foliis lineari-lanceolatis acuminatis flaccidis, corollis fructu majoribus. *Roy. lugdb.* 255.

Aparine minima. *R. Syn.* 225. t. 9. f. 1.

Anglis,

Anglis, the least Goose-grafs.

Habitat in muris et pratis sterilibus subhumidis. 4. VII.

- boreale.* 11. GALIUM foliis quaternis lanceolatis trinerviis glabris, caule suberecto, feminibus hispidis. *Fl. lapp.* 60. *suec.* 124. *Sp. pl.* 108.
 Mollugo montana erecta quadrifolia. *R. Syn.* 224.
 Rubia erecta quadrifolia. *Bauh. hist.* III. 716.
 Rubia pratensis lævis, acuto folio. *Bauh. pin.* 333.
pr. 145.

Anglis, Crosswort-madder.

Habitat in pratis montosis, in comitatibus Westmorlandico & Eboracensi. 4. VI.—VIII.

Flores tetrapetali superi.

C O R N U S. 139.

Involucrum tetraphyllum sæpius. *Petala* supera quatuor. *Drupa* nucleo biloculari.

- sanguinea.* 1. CORNUS arborea, cymis nudis. *it. W-goth.* 32. *Sp. pl.* 117.
 Cornus umbellis involucro multoties longioribus. *Hort. cliff.* 38. *Roy. lugdb.* 249. *Hall. helv.* 463.
 Cornus femina. *Bauh. pin.* 447. *Ger. Em.* 1467. *Park.* 1521. *R. Syn.* 460.
Anglis, the female Cornel, Dog-berry-tree, Gatter-tree, or Prickwood.
 Habitat in sepibus et sylvis. 2. VI.

- herbacea.* 2. CORNUS herbacea, ramis binis. *Fl. lapp.* 65. *t.* 5. *f.* 3. *Sp. pl.* 118.
 Cornus pumila herbacea, Chamæpericlymenum dicta. *Dill. alth.* 108. *t.* 91.
 Chamæpericlymenum. *Park.* 1461. *Ger. Em.* 1296. *R. Syn.* 261.
 Periclymenum humile. *Bauh. pin.* 302.
Anglis, dwarf Honeyfuckle.
 Habitat in Northumbriæ montibus Chevioticis dictis. *R. Syn.* in Pickering moor in comitatu Eboracensi. 4. VI.

Flores

Flores incompleti inferi.

ALCHEMILLA. 153.

Calyx octofidus. *Cor.* nulla. *Sem.* conicum.

1. **ALCHEMILLA** foliis lobatis. *Fl. suec.* 141. *Sp. vulgaris.*
pl. 123. *Alchemilla* foliis simplicibus. *Fl. lapp.* 66.
Alchemilla. *Ger. Em.* 949. *R. Syn.* 158. *vulgaris.*
Bauh. pin. 319.
Alchemilla major vulgaris. *Park.* 538.
Anglis, Ladies Mantle.
Habitat in pratis et pascuis montosis. 4. VI.—VIII.
Pharmac. Alchemillæ Herba.

2. **ALCHEMILLA** foliis lobatis, plicatis, acute serratis, fericeis.
Alchemilla alpina pubescens minor. *Tourn.* 508.
Pluk. Ph. t. 240. *f.* 1.
Anglis, the least Ladies Mantle.
Habitat in montibus Westmorlandicis. 4. VII.

3. **ALCHEMILLA** foliis digitatis ferratis. *Fl. suec. alpina.*
142. Sp. pl. 123.
Alchemilla alpina pentaphyllos. *R. Syn.* 158.
Pentaphyllum petrosum, Heptaphyllum Clusii. *Ger.*
Em. 988.
Tormentilla alpina folio fericeo. *Bauh. pin.* 326.
Tormentilla argentea. *Park.* 393.
Anglis, Cinquefoil Ladies Mantle.
Habitat in montosis. In rupibus juxta stagnum Hulf-
water dictum inveni, non longe a Penrith Cumber-
landiæ oppido; non procul a Kendal in Westmor-
landia, in vallicula umbrosa Longsledale dicta. *D.*
Richardson. R. Syn. 4. VII.

DIGYNIA.

APHANES. 154.

Cal. octofidus. *Cor.* o. *Sem.* bina, nuda.

- arvensis.* 1. APHANES. *Hort. cliff.* 39. *Fl. suec.* 143. *Gron. virg.* 17. *Sp. pl.* 123.
Percepier Anglorum. *Lob.* 727. *Ger. Em.* 1594.
R. Syn. 159.
Polygonum felinoides. *Park.* 449.
Chærophyllo nonnihil similis. *Bauh. pin.* 152.
Alchemilla minima montana. *Col. Ec.* 145. t. 146.
Anglis, Pursley-piert.
Habitat in arvis et pascuis siccioribus et glareosis frequens. ☉. V.

TETRAGYNIA.

POTAMOGETON. 160.

Cal. o. *Petala* quatuor. *Stylus* nullus. *Sem.* quatuor.

- natans.* 1. POTAMOGETON foliis oblongo-ovatis petiolatis natantibus. *Sp. pl.* 126.
Potamogiton rotundifolium. *Bauh. pin.* 193. *R. Syn.* 148.
Potamogiton latifolium. *Ger. Em.* 821.
Fontalis major latifolia vulgaris. *Park.* 1254.
Anglis, broad leaved Pondweed.
Habitat in fluviis et stagnis passim. 4. VIII.
- perfoliatum.* 2. POTAMOGETON foliis cordatis amplexicaulibus. *Sp. pl.* 126.
Potamogiton perfoliatum. *R. hist. I.* 188. *Syn.* 149.
Potamogiton foliis latis splendentibus. *Bauh. pin.* 193.
Potamogiton. 3. *Dodonæi.* *Ger. Em.* 822.

Pota-

β Potamogiton rotundifolium alterum. *Lofæl. fl. pr.*

205. t. 15.

Anglis, perfoliated Pondweed.

Habitat in fluviis frequens. 4. VI. VII.

3. POTAMOGETON foliis lanceolatis planis in pe-
tiolos defineniibus. *Hort. cliff.* 40. *Fl. suec.* 147.
Sp. pl. 126.

Potamogiton aquis immerfum folio pellucido, lato,
oblongo, acuto. *R. Syn.* 148.

Potamogiton foliis angustis splendentibus. *Bauh. pin.*
193.

Potamogiton longis acutis foliis. *Ger. Em.* 822.

Anglis, long leaved Pondweed.

Habitat in fluviis et lacubus. 4. VI.

4. POTAMOGETON foliis lanceolatis alternis undu-
latis ferratis. *Roy. lugdb.* 212. *Fl. suec.* 148. *Sp. pl.*
126.

Potamogiton f. fontinalis crispa. *Bauh. hist.* III. 778.
R. Syn. 149.

Potamogeton foliis crispis f. *Lactuca ranarum.* *Bauh.*
pin. 193.

Tribulus aquaticus minor *Quercus floribus.* *Ger. Em.*
824.

Tribulus aquaticus minor prior. *Park.* 1248.

Anglis, the greater Water-caltrops.

Habitat in fluviis, fossis, stagnis ubique. 4. V. VI.

5. POTAMOGETON foliis lanceolatis oppositis sub-
undulatis. *Sp. pl.* 126.

Potamogeton f. Fontalis media lucens. *Bauh. hist.*
III. 777. *R. Syn.* 149.

Potamogeton longo ferrato folio. *Bauh. pin.* 193.

Tribulus aquaticus minor, *Moscatellæ floribus.* *Ger.*
Em. 823.

Tribulus aquaticus minor alter. *Park.* 1248.

β Potamogeton folio angusto pellucido fere gramineo.
Doody. R. Syn. 148.

Anglis, the lesser Water-caltrops, or Frog's-lettuce.

Habitat in fluviis leniter fluentibus. 4. VI.

- compressum.* 6. POTAMOGETON foliis linearibus obtusis caule compresso. *Fl. suec.* 149. *Sp. pl.* 127.
 Potamogeton caule compresso, folio graminis canini.
R. hist. I. *Syn.* 149.
Anglis, small branched Pondweed with a flat stalk.
Habitat in fluviis. In Cambridge and many other rivers. R. Syn. 4. VI. VII.
- gramineum.* 7. POTAMOGETON foliis lineari-lanceolatis alternis sessilibus stipula latioribus. *Fl. suec.* *Sp. pl.* 127.
 Potamogeton gramineum latiusculum foliis et ramificationibus dense stipatis. *Budd. H. Sic. Vol.* 27. *R. Syn.* 149. t. 4. f. 3.
 β Potamogeton gramineum latum. *Pet. h. br. t.* 5. f. 10.
Anglis, glass leaved Pondweed.
Habitat in fossis prope Deptford. R. Syn. 4.
- pusillum.* 8. POTAMOGETON foliis linearibus oppositis alternisque distantibus: basi patentibus, caule tereti. *Fl. suec.* 153. *Sp. pl.* 127. *Dalib. paris.* 56.
 Potamogeton pusillum, gramineo folio, caule tereti. *R. hist.* 190. *Syn.* 150.
 Potamogeton minimum capillaceo folio. *Baub. pin.* 193. *pr.* 101.
 Potamogeton gramineum angustifolium. *Pet. h. br. t.* 5. f. 11.
Anglis, small grass-leaved Pondweed.
Habitat in fossis et stagnis. 4. VI. VIII.
- pectinatum.* 9. POTAMOGETON foliis fetaceis parallelis approximates distichis. *Fl. suec.* 150. *Sp. pl.* 127. *Dalib. paris.* 55.
 Potamogeton millefolium, seu foliis gramineis ramosum. *R. Syn.* 150.
 Potamogeton gramineum ramosum. *Baub. pin.* 193.
 Millefolium tenuifolium. *Ger. Em.* 828.
Anglis, Fennel-leaved Pondweed.
Habitat in fluviis. In fluvio Serpentine river dicto in Hyde-park prope Londinum. 4. VI.

10. POTAMOGETON foliis linearibus alternis distinctis inferne vaginantibus. *Fl. suec.* 152. *Sp. pl.* 127.

Potamogiton maritimum ramosissimum, grandiusculis capitulis, capillaceo folio nostras. *Pluk. Ph. t.* 216. *f.* 5. *R. Syn.* 150.

Potamogeton foeniculaceum. *Pet. h. br. t.* 5. *f.* 13. *Anglis*, Sea Pondweed.

Habitat in fossis maritimis. In fossis pone Sheerness, abunde. *R. Syn.* 2. VIII.

R U P P I A. 161.

Cal. o. *Cor.* o. *Sem.* quatuor pedicellata.

1. RUPPIA. *Hort. cliff.* 436. *It. W-goth.* 186. *Fl. maritima.* *suec.* 154. *Sp. pl.* 54.

Buccaferra maritima foliis acutissimis. *Mich. gen.* 72. *t.* 35.

Potamogiton maritimum gramineis longioribus foliis, fructu fere umbellato. *Pluk. Alm.* 305. *R. Syn.* 135.

Potamogiton maritimum pusillum alterum, feminibus singulis longis pediculis infidentibus. *Pluk. Ph. t.* 248. *f.* 4.

Potamogiton maritimum pusillum alterum. *R. hist. I.* 190.

Anglis, Sea-grass.

Habitat in maritimis. In fossis in insula Shepey copiose. 2. VIII.

S A G I N A. 162.

Cal. tetraphyllus. *Petala* quatuor. *Capf.* quadrilobularis, quadrivalvis.

1. SAGINA ramis procumbentibus. *Fl. lapp.* 157. *suec.* *procumbens.* *Sp. pl.* 128.

Alfinella muscosa flore repens. *Cat. giff.* 81. *R. Syn.* 345.

Saxifraga graminea pusilla, flore parvo tetrapetala. *R. Syn.* 2. 206.

Saxifraga anglicana alsine folio. *Ger. Em.* 568.

Saxifraga pusilla graminea, flore parvo herbido et muscosa. *Merr. pin.*

Alsine

TETRANDRIA TETRAGYNIA.

Alfine saxifraga perexigua tenuifolia, flore parvo tetrapetalo radice reptatrice. *Pluk. alm.* 23.

Caryophyllus minimus muscosus nostras. *Park.* 1340.

β Saxifraga anglica Alfine folia annua. *Plot. hist. n. ox.* t. 9. f. 7.

γ Saxifraga graminea pusilla foliis brevioribus, crassioribus et succulentioribus. *R. Syn.* 345.

Anglis, Pearl-wort, or Chickweed Breakstone.

Habitat in muris, tectis, pratis arenosis ubique. ☉. VI.

erecta. 2. SAGINA caule erecto subunifloro. *Sp. pl.* 128.

Alfinella foliis caryophylleis. *Cat. giff.* 47. *R. Syn.* 344. t. 15. f. 4.

Alfine tetrapetalos Caryophylloides, quibusdam Holosteum minimum. *R. Syn. ed.* 2. 206.

Alfine verna glabra. *Vaill. paris.* 6. t. 3. f. 2.

Anglis, the least Stichwort.

Habitat in pascuis glareosis frequens. ☉. IV. V.

Classis

Classis V.

PENTANDRIA
MONOGYNIA.

* Flores monopetali, inferi tetraspermi. Asperifoliæ R.

MYOSOTIS. 165.

Cor. hypocrateriformis, quinquefida, emarginata :
fauce clausa fornicibus.MYOSOTIS feminibus nudis : foliorum apicibus *scorpioides*
callofis. *Syst. Nat.* 915. *Fl. suec.* 175. *Sp. pl.*
131.Myosotis foliis hirsutis. *Hort. cliff.* 45.Myosotis scorpioides hirsuta. *Park.* 691. *R. Syn.* 229.Myosotis scorpioides hirsuta. *Ger. Em.* 337.Echium scorpioides arvense. *Bauh. pin.* 254.β Myosotis scorpioides latifolia hirsuta. *Merr. pin. R.*
Syn. 229. t. 9. f. 2.γ Myosotis scorpioides hirta minor. *Merr. pin. R. Syn.*
229.Echium scorpioides minus flosculis luteis. *Bauh. pin.*
254.♂ Myosotis scorpioides palustris. *Ger. Em.* 337. *R. Syn.*
229.Echium scorpioides palustre. *Bauh. pin.* 254.*Anglis*, Mouse-ear scorpion-grass.Habitat in pratis exaridis ubique. β in fossis, et ad sca-
turagines et fontes. 4. IV.—VIII.

LITHOSPERMUM. 166.

Cor. infundibuliformis fauce perforata, nuda. *Cal.*
quinquepartitus.1. LITHOSPERMUM feminibus lævibus, corollis vix *officinalis*. X
calycem superantibus, foliis lanceolatis. *Syst. Nat.*
913. *Sp. pl.* 133.Lithospermum majus erectum. *Bauh. pin.* 258.Lithospermum minus. *Ger. Em.* 609.

F

Lithos-

Lithospermum vulgare minus. *Park.* 432.

Lithospermum feu *milium folis.* *Baub. hist.* 3. 590.
R. Syn. 228.

Anglis, Gromwell, Gromill or Graymill.

Habitat in locis glareosis et aridis. *Circa* *Bristolium copiose*; *in sylva* *Charlton wood dicta in Cantio*; *Dr. Wilmer.* 4. V. VI. *Near Hadleigh in Suffolk.*
Pharmac. *Lithospermi semina.*

purpureo
cæruleum.

2. *LITHOSPERMUM* feminibus lævibus, corollis calycem multoties superantibus. *Hort. cliff.* 46. *Sp. pl.* 132.

Lithospermum minus repens latifolium. *Baub. pin.* 258.

Lithospermum majus. *Ger. Em.* 609.

Lithospermum vulgare majus. *Park.* 258.

Lithospermum majus Dodonæi, flore purpureo, semine *Anchusæ.* *Baub. hist.* III. 572. *R. Syn.* 229.

Anglis, the lesser creeping Gromwell.

Habitat in pascuis montosis. *On the top of a bushy hill near Denbigh town in Wales, on the north side of the town; also in Somersetshire, not far from Taunton.*
R. Syn. 4. VI.

X *arvense.*

3. *LITHOSPERMUM* feminibus rugosis, corollis vix calycem superantibus. *Fl. suec.* 160. *Sp. pl.* 132.

Lithospermum arvense radice rubra. *Baub. pin.* 258.
Park. 432. *R. Syn.* 227.

Anchusa degener facie Milii folis. *Ger. Em.* 610.

Anglis, bastard Alkanet.

Habitat in arvis frequens. ☉. V. VI.

A N C H U S A. 167.

Cal. infundibuliformis fauce clausa fornicibus.

sempervirens.

1. *ANCHUSA* pedunculis diphyllis capitatis. *Sp. pl.* 134.

Buglossum latifolium sempervirens. *Baub. pin.* 256.

Borrago sempervirens. *Ger. Em.* 797. *R. Syn.* 227.
Anglis, evergreen Alkanet.

Habitat in ruderatis. *Near Hornsplace near Rochester in Kent,* *R. Syn.* *Juxta Vauxhall prope Londinum.* 4. V. VI.

CYN OGLOSSUM. 168.

Cor. infundib. fauce clausa fornicibus. *Semina* depressa, interiore tantum latere stylo affixa.

CYNOGLOSSUM flaminibus corolla brevioribus, foliis lato-lanceolatis tomentosis sessilibus. *Fl. succ.* 162. *Sp. pl.* 134. *officinale*. x

Cynoglossum foliis lanceolato-ovatis, corollis calycem æquantibus. *Roy. lugdb.* 406. *Mat. med.* 60.

Cynoglossum majus vulgare. *Bauh. pin.* 257. *Ger. Em.* 804. *Park.* 511.

β Cynoglossum folio virenti. *Bauh. hist. III.* 600. *R. Syn.* 226.

Anglis, Hound's-tongue.

Habitat in ruderatis et ad vias. β In the London road, between Kelveden and Witham in Essex; but more plentifully about Braxted by the way-sides. At South-end by Eltham, Mr. J. Sherard. In a hedge facing the road on Stamford-hill, between Newington and Toddenham. *R. Syn.* 4. VI.

Pharmac. Cynoglossi Radix.

PULMONARIA. 169.

Cor. infundib. fauce pervia. *Cal.* prismatico-pentagonus.

1. PULMONARIA foliis radicalibus ovato-cordatis *officinalis*. scabris. *Sp. pl.* 135. *Mat. med.* 62. *Hall. helv.* 516.

Pulmonaria foliis Echii. *Ger. Em.* 808.

Symphytum maculosum f. Pulmonaria latifolia. *Bauh. pin.* 259.

Pulmonaria latifolia. *Park.* 248.

Anglis, Bugloss-cowslips, or long-leaved Sage of Jerusalem.

Habitat in nemorosis. In a wood by Holbury-house, in the new forest, Hampshire. *R. Syn.* 4. V.

2. PULMONARIA calycibus abbreviatis, foliis ovatis, *maritima*. caule ramoso procumbente. *Sp. pl.* 136.

Cerithe foliis ovatis petiolatis. *Hort. cliff.* 48. *Roy. lugdb.* 408.

PENTANDRIA MONOGYNIA.

Cerinthē maritima procumbens. Dill. *elth.* 75. t. 65.
f. 75.

Echium marinum. Phyt. Br. Cat. Angl. R. Syn. 228.

Buglossum marinum. Pet. H. Br. t. 29. f. 3.

Cynoglossum procumbens glaucophyllum maritimum
nostras floribus purpureo-cæruleis seminibus lævibus,
Pl. ph. t. 172. f. 3.

Buglossum dulce ex Insulis Lancastriæ. Park. 765.
Anglis, Sea Bugloss.

Habitat in maritimis et arena mobili. Near White-
haven in Cumberland, and against Bigger in the isle
of Walney, in Lancashire, Mr. Lawson. Near
Trefarthen in Anglesea, and in abundance by the ri-
ver Euyfni, in the way from Dinardinlle to Clynog in
Carnarvonshire, Mr. Lhwyd. R. Syn. 4. VII.

SYMPHYTUM. 170.

Corollæ limbus tubulato-ventricosus : fauce clausa
radiis subulatis.

- X officinale. I. SYMPHYTUM foliis ovato-lanceolatis decurrenti-
bus. *Mat. med.* 61. *Fl. suec.* 165. *Sp. pl.* 136.
Symphytum majus vulgare. Park. 523.
Symphytum, consolida major. *Bauh. pin.* 259. *Ger.*
Em. 660.

- β *Symphytum flore purpureo.* *Merr. pin.*
Anglis, Comfrey.
Habitat in aquis et ad ripas fluviorum. 4.
Pharmacop. Consolidæ majoris Radix.

BORAGO. 172.

Corolla rotata : fauce radiis clausa.

- X officinalis. * I. BORAGO foliis omnibus alternis, calycibus paten-
tibus. *Sp. pl.* 137.
Borrigo hortensis. *Ger. Em.* 797. *R. Syn.* 228.
Borrigo floribus cæruleis et albis. *Bauh. hist. III.*
574.
Buglossum latifolium, Borrigo. *Bauh. pin.* 256.
Anglis, Borage.
Habitat in ruderatis viis, muris, frequens. 4. VI. VIII.
Pharm. Borriginis Folia, Flores.

A S P E R U G O. 173.

Cal. fructus compressus lamellis plano parallelis,
sinuatis.

- I. ASPERUGO. *Fl. lapp.* 76. *Fl. suec.* 159. *Sp. pl.* *procumbens.*
138. *Hall. herb.* 522.

Buglossum sylvestre, caulibus procumbentibus. *Baub.*
pin. 257. *Hist. Ox.* III. 439. t. 26. f. 13.

Borago minor sylvestris. *Park.* 765.

Aparine major. *Plinii Ger. Em.* 1122.

Anglis, small wild Bugloss, great Goose-grass, and
German Madwort.

Habitat in ruderatis quinquibus. By Boxley in Suffex,
and in the Holy Island. *R. Syn.* 228. *prope* Purfleet
in Essex; D. Alchorne. ☉. IV. V.

L Y C O P S I S. 174.

Corollæ tubus incurvatus.

- I. LYCOPSIS foliis lanceolatis hispidis, calycibus flo. *arvensis.*
rescentibus erectis. *Fl. lapp.* 77. *Fl. suec.* 167.
Sp. pl. 139.

Buglossum sylvestre minus. *Baub. pin.* 256. *Park.*
765.

Buglossa sylvestris minor. *Ger. Em.* 799. *R. Syn.*
227.

Anglis, small wild Bugloss.

Habitat in arvis et ad vias frequens. ☉. VI.—IX.

E C H I U M. 175.

Corolla irregularis, fauce nuda.

- I. ECHIUM caule simplici erecto, foliis caulinis lan. *vulgare.*
ceolatis hispidis, floribus spicatis lateralibus stami-
nibus corollam æquantibus. *Mill. Dict. Hort. edit.* 6.

Echium caule simplici erecto, foliis caulinis lanceo-
latis hispidis, floribus spicatis lateralibus. *Hort.*
cliff. 43. *Roy. lugdb.* 407. *Sp. pl.* 139.

Echium vulgare. *Baub. pin.* 254. *Baub. hist.* III.
586. *Park.* 414. *Ger. Em.* 802. *R. Syn.* 227.

Anglis, Vipers Bugloss.

F 3

Habitat

PENTANDRIA MONOGYNIA.

Habitat in arvis et ad vias ; in agro Cantiano frequens.
 4. VII.

Pharmacop. Echii Herba, Flores.

- anglicum. 2. ECHIUM caule simplici erecto, foliis lanceolato-linearibus hispidis, floribus spicatis lateralibus. Staminibus corolla longioribus. *Mill. Dict. hort. edit. 6.*
 Echium alterum sive Lycopsis Anglica. *Merr. pin. R. Syn. 228.*

Lycopsis Anglica Lobelii.

Anglis, English Vipers Bugloss.

Habitat in arvis pascuis sterilioribus et ad vias frequens.
 4. VIII.

- italicum. 3. ECHIUM corollis subæqualibus vix calycem excedentibus: margine villosis. *Syst. Nat. 916. Sp. pl. 139.*

Lycopsis. *Bauh. pin. 255. Park. 519. Ger. Em. 802. R. Syn. 227.*

Anglis, Wall Vipers Bugloss.

Habitat in arenosis. In the isle of Jersey, on the sandy grounds, near S. Hilary, plentifully. 4. VII.

* *Fl. monopetali, inferi, angiospermi.*

PRIMULA. 180.

Involucrum umbellulæ. *Corollæ* tubus cylindricus :
 ore patulo.

- vulgaris. 1. PRIMULA foliis dentatis rugosis, scapo unifloro.
 Primula veris vulgaris. *Park. 536. R. Syn. 284.*
 Primula veris minor. *Ger. Em. 781.*
 Verbasculum sylvarum majus singulari flore. *Bauh. pin. 241.*
 Anglis, common Primrose.
 Habitat in sylvis sepibus et ericetis ubique. 4. IV. V.

- peris. 2. PRIMULA foliis dentatis rugosis, scapo multifloro,
 Primula foliis dentatis rugosis. *Sp. pl. 142.*
 Primula veris major. *Ger. Em. 781. R. Syn. 284.*
 Verbasculum pratense odoratum. *Bauh. pin. 241.*

Para-

PENTANDRIA MONOGYNIA.

71

Paralyfis vulgaris pratensis flore flavo simplici odorato.
Park. th. 535. *Par.* 244.

β Verbasculum pratense aut sylvaticum inodorum. *Baub. pin.* 241.

Primula pratensis inodora lutea. *Ger. Em.* 780. *R. Syn.* 284.

Anglis, α Pagils or Cowslips. β Great Cowslips or Oxslips.

Habitat in pratis 24. IV. V.

Pharmacop. Paralyfis Flores.

3. Primula foliis crenatis glabris, florum limbo plano. *farinosa.*
Fl. suec. 172. *Sp. pl.* 143.

Primula veris flore rubro. *Ger. Em.* 783.

Verbasculum umbellatum alpinum minus. *Baub. pin.* 247. *R. Syn.* 285.

Paralyfis minor flore rubro. *Park. par.* 246.

Anglis, Bird's-eye.

Habitat in paludibus montosis in comitatibus Westmorlandico et Eboracensi copiose. 24. V.

M E N Y A N T H E S. 185.

Corolla hirsuta. Stigma bifidum. Caps. unilocularis.

1. MENYANTHES foliis ternatis. *Fl. suec.* 173. *lapp. trifolia.* 80. *Hall. belv.* 487. *Sp. pl.* 145.

Menianthe palustre triphyllum latifolium et angustifolium. *Tourn.* 117. *R. Syn.* 285.

Trifolium palustre. *Baub. hist. II.* 389. *Baub. pin.* 327.

Trifolium paludosum. *Ger. Em.* 1194. *Park.* 1212.

Anglis, Marsh Trefoil, or Buck-bean.

Habitat in palustribus et aquosis frequens. 24. VI. VII.

Pharmacop. Trifolii paludosi Herba.

2. MENYANTHES foliis cordatis integerrimis, corollis ciliatis. *Sp. pl.* 145. *Nymphoides.*

Nymphæa lutea minor flore fimbriato. *Baub. hist. II.* 772. *Baub. pin.* 194. *R. Syn.* 368.

Anglis, fringed Water-lilly.

Habitat in fluviis et fossis majoribus. 24. VI. VII.

PENTANDRIA MONOGYNIA.

HOTTONIA. 186.

Cor. hypocrateriformis. *Stam.* tubo corollæ imposita.
Caps. unilocularis.

palustris. 1. HOTTONIA pedunculis verticillato-multifloris. *Syst.*
Nat. 919

Hottonia. *Boerb. ind. alt.* 207. *R. Syn.* 285. *Sp. pl.*
 145.

Millefolium aquaticum seu *Viola aquatica* caule nudo.
Baub. pin. 141.

Millefolium aquaticum floridum seu *Viola aquatica.*
Park. 1256.

Viola palustris. *Ger. Em.* 826.

Anglis, Water-violet.

Habitat in fossis et paludibus frequens. 4. VII. VIII. V

LYSIMACHIA. 188.

Cor. rotata. *Caps.* globosa, mucronata, decem-
 valvis.

* *Pedunculis multifloris.*

× *vulgaris.* 1. LYSIMACHIA paniculata racemis terminalibus. *Fl.*
suec. 175. *Sp. pl.* 146.

Lyfimachia foliis lanceolatis, racemo composito ter-
 minali. *Virid. cliff.* 415. *Roy. lugdb.* 416.

Lyfimachia lutea. *Baub. hist. II.* 901. *Ger. Em.* 474.
R. Syn. 282.

Lyfimachia lutea major. *Baub. pin.* 245. *vulgaris*
Park. 544.

β *Lyfimachia lutea foliis brevioribus et obtusis, floribus*
in summitate congestis. *R. Syn.* 282.

Anglis, yellow Willow-herb, or Loosestrife.

Habitat in paludibus nemorosis et ad ripas fluviorum. 4.
 VI. VII.

Pharmac. Lyfimachia Herba.

* *hirsiflora.* 2. LYSIMACHIA racemis lateralibus pedunculatis. *Fl.*
suec. 176. *Sp. pl.* 147.

Lyfimachia ex alis foliorum thyrsifera. *Fl. lapp.* 82.

Lyfimachia lutea flore globofo. *Ger. Em.* 475. *Park.*
 544. *R. Syn.* 283.

Lyfimachia

Lyfimachia bifolia flore globofo luteo. *Baub. pin.* 245.
Anglis, tufted Loofeftrife.

Habitat in paludibus et ad ripas fluviorum sed rarius. *Brickhills*
Found by Mr. Dodsworth in the East Riding of Ely
 Yorkshire. In Nynnechtiked, Anglesea, Mr.
 Lhwyd. By Kings-Langley in Hertfordshire. *Ph.*
Br. R. Syn. 4. VI.

** *Pedunculis unifloris.*

3. *LYSIMACHIA* foliis ovatis acutis, floribus solitariis, nemorum.
 caule procumbente. *Hort. cliff.* 52. *Roy. lugdb.*
 416. *Sp. pl.* 148.

Anagallis lutea. *Ger. Em.* 618. nemorum. *Baub. pin.*
 252.

Anagallis flore luteo. *Park.* 558.
Anglis, yellow Pimpernell of the Woods.

Habitat in sylvis et umbrosis humidioribus. 4. V. VI.

4. *LYSIMACHIA* foliis subcordatis, floribus solitariis Nummu- x
 caule repente. *Fl. suec.* 177. *Sp. pl.* 148. *Hall. laria.*
Helv. 481.

Nummularia. *Ger. Em.* 630. vulgaris *Park.* 555.

Nummularia major lutea. *Baub. pin.* 309.

Anglis, Moneywort.

Habitat in pratis et pascuis humidis. 4. VI.

5. *LYSIMACHIA* foliis ovatis acutiusculis, pedunculis tenella. x
 folio longioribus, caule repente. *Sp. pl.* 148.

Lyfimachia foliis orbiculatis planis, floribus solitariis
 rubellis caule repente. *Sauv. monsp.* 135.

Nummularia minor flore purpurascente. *Baub. pin.*

310. *Park.* 555. *Ger. Em.* 630. *R. Syn.* 283.

Anglis, purple Moneywort. *Bagshot heath* 8 1/2 miles from London.

Habitat in paludibus cespitosis. 4. VII. VIII.

A N A G A L L I S. 189.

Corolla rotata. *Caps.* circumscissa.

1. *ANAGALLIS* foliis indivisis, caule procumbente. *arvensis.*
Sp. pl. 148. *Fl. suec.* 178.

Ana-

PENTANDRIA MONOGYNIA.

Anagallis flore phæniceo. *Bauh. pin.* 252. *Park.* 558.
R. Syn. 282.

Anagallis mas. *Ger. Em.* 617.

β Anagallis phænicea foliis amplioribus ex adverso quaternis. *Pluk. alm.* 29. *R. Syn.* 282.

γ Anagallis fæminea. *Ger. Em.* 617. *R. Syn.* 282. cæruleo flore. *Bauh. pin.* 252.

♂ Anagallis terrestris flore albo. *R. Syn.* 282.

Anglis, male Pimpernel, γ female Pimpernel.

Habitat in arvis, et locis arenosis, γ in arvis inter Camberwell et Stockwell. ☉. V.—VIII.

CONVOLVULUS. 198.

Cor. campanulata, plicata. *Stigm.* duo. *Capf.* bilocularis: loculis dispermis.

* *Caule volubili.*

arvensis. 1. CONVOLVULUS foliis sagittatis utrinque acutis, pedunculis unifloris. *Fl. juec.* 181. *Sp. pl.* 153.
Dalib. parif. 65.

Convolvulus minor vulgaris. *Park.* 171. *R. Syn.* 275.

Convolvulus minor arvensis. *Bauh. pin.* 294.

Smilax lævis minor. *Ger. Em.* 861.

β *Convolvulus arvensis minimus.* *D. Dent.* *R. Syn.* 276.

γ *Convolvulus flore minimo ad unguem fere secto.* *R. Syn.* 276.

Anglis, small Bindweed.

Habitat in arvis et ad vias ubique. 4. VI. VII.

sepium. 2. CONVOLVULUS foliis sagittatis postice truncatis, pedunculis tetragonis unifloris. *Syst. nat.* 922.
Sp. pl. 152.

Convolvulus major albus. *Bauh. pin.* 294. *Park.* 163. *R. Syn.* 275.

Smilax lævis major. *Ger. Em.* 861.

Anglis, great Bindweed.

Habitat in sepibus subhumidis. 4. VII. VIII.

** *Caule*

** *Caule prostrato.*

- 3 CONVULVULUS foliis reniformibus, pedunculis *Soldanella*.
 unifloris. *Sp. pl.* 159. *Mat. med.* 81.
Convolvulus maritimus *Soldanella dicta.* *R. Syn.* 276.
Soldanella marina. *Ger. Em.* 861.
Soldanella maritima minor. *Bauh. pin.* 295.
Soldanella vulgaris. *Park.* 161.
Anglis, Sea Bindweed. Scottish Scurvy-grass.
Habitat in littoribus maritimis frequens. 4. VII.

POLEMONIUM. 200.

Cor. quinquepartita. *Stam.* squamis basin corollæ
 claudentibus inserta. *Stigma* trifidum. *Capsf.* tri-
 locularis, supera.

1. POLEMONIUM foliis pinnatis floribus terminalibus *cæruleum*.
 erectis calycibus tubo corollæ longioribus. *Syst. Nat.*
 925. *Sp. pl.* 162.
Valeriana cærulea. *Bauh. pin.* 164.
Valeriana græca. *Ger. Em.* 1076. *Park.* 122.
Anglis, great Valerian, Ladder to Heaven, and Jacob's
 Ladder.
Habitat in dumetis infra Malham-cove, eundo ab oppido
ad Malham-cove dextrorsum, ad sinistram enim re-
una invenitur planta; Dr. Richardson, R. Syn. 280.
 4. VI. *in the Hills between Morton in marsh & Broadway.*

VERBASCUM. 217.

Cor. rotata, subinæqualis. *Capsf.* bilocularis, bivalvis.

1. VERBASCUM foliis decurrentibus utrinque tomen- *Thapsus*.
 tosis. *Vir. cliff.* 13. *Mat. med.* 90. *Fl. suec.* 195.
Sp. pl. 177.
Verbascum caule simplici superne floribus sessilibus
clavato, foliis utrinque lanigeris. *Hort. cliff.* 55.
Verbascum mas latifolium luteum. *Bauh. pin.* 239.
R. Syn. 287.
Verbascum album vulgare, sive Thapsus barbatus
communis. *Park.* 60.
Thapsus barbatus. *Ger. Em.* 773.

Anglis,

PENTANDRIA MONOGYNIA.

Anglis, great white Mullein, High-taper, Cow's Lungwort.

Habitat in locis cretaceis, arenosis, sterilissimis passim.
 ♂. VII.

Pharmac. Verbasci Folia.

Lychnitis. 2. VERBASCUM foliis cuneiformi-oblongis. *Hort. ups.* 45. *Sp. pl.* 177.

Verbascum foliis ovato-acutis subtus villosis, crenatis, spicis laxis lateralibus et terminalibus. *Hort. cliff.* 54.

Verbascum flore albo parvo. *Bauh. hist.* III. 873.
R. Syn. 287.

Verbascum Lichnitis flore albo parvo. *Bauh. pin.* 240.

Verbascum mas foliis longioribus. *Park.* 60.

Verbascum Lichnitis Matthioli. *Ger. Em.* 775.

β Verbascum pulverulentum flore luteo parvo. *Bauh. hist.* III. 872. *R. Syn.* 287.

Anglis, Hoary Mullein, or white flowered Mullein.

Habitat in locis arenosis et cretaceis, in agro Cantiano frequens. β circa mœnia Nordovici urbis; et ad Buriam oppidum. *R. Syn.* ♂. VII.

nigrum.] 3. VERBASCUM foliis cordato-oblongis petiolatis.
Mat. med. 91. *Sp. pl.* 178.

Verbascum foliis ovatis crenatis subtus tomentosis caule angulato. *Vir. cliff.* 13. *Roy. lugdb.* 421. *Dalib. parif.* 91.

Verbascum nigrum flore parvo, apicibus purpureis.
Bauh. hist. III. 873. *R. Syn.* 288.

Verbascum nigrum flore ex luteo purpurascente. *Bauh. pin.* 240.

Verbascum nigrum. *Ger. Em.* 775. vulgare *Park.* 61.

Verbascum nigrum salvifolium luteo flore. *Lob. adv.* 241. 466.

Anglis, Sage-leaved black Mullein.

Habitat ad vias in agro Cantiano passim. 24. VII.

Blattaria. 4. VERBASCUM foliis amplexicaulibus oblongis glabris, pedunculis solitariis. *Hort. ups.* 46. *Sp. pl.* 178.

Verbascum foliis glabris ferrato-dentatis, radicalibus sinuatis. *Roy. lugdb.* 421. *Sauv. monsp.* 276.

Blattaria

Blattaria lutea. *Baub. hist. III. 874. R. Syn. 288.*
Blattaria lutea folio longo laciniato. *Baub. pin. 240.*
Blattaria flore luteo. *Park. 64.*
Blattaria Plinii. *Ger. Em. 776.*
Anglis, yellow Moth Mullein.
Habitat in pascuis glareosis. Prope Charlton in Cantio.
 ○. VI.

V I N C A. 261.

- Cor. contorta Folliculi duo, erecti. Semina nuda.*
1. VINCA caulibus procumbentibus, foliis lanceolato-*minor.*
 ovatis. *Sp. pl. 209.*
Vinca Pervinca minor. *Ger. Em. 894. R. Syn. 268.*
Vinca Pervinca vulgaris. *Park. 380.*
Clematis daphnoides minor. *Baub. pin. 301. Baub.*
hist. II. 130.
Anglis, Periwinkle.
Habitat in sepibus et sylvis sed rarius. In sylvula
*prope Hampsteadum. 4. V. **
Pharm. Vincæ Herba.
2. VINCA caulibus erectis, foliis ovatis. *Sp. pl. 209. major.*
Clematis daphnoides major. *Baub. pin. 302. R. Syn.*
268.
Clematis daphnoides latifolia, f. Vinca Pervinca ma-
major. *Park. 380.*
Clematis daphnoides, f. Pervinca major. *Ger. Em.*
894.
Anglis, the greater Periwinkle.
Habitat in sylvis et sepibus passim. 4. V.

H Y O S C Y A M U S. 219.

- Cor. infundibuliformis, obtusa. Stamina inclinata.*
Caps. operculata, bilocularis.
1. HYOSCYAMUS foliis amplexicaulibus. *Hort. cliff. niger.*
56. Mat. med. 86. Fl. suec. 199. Sp. pl. 179.
Hyoscyamus vulgaris. *Baub. hist. III. 627. R. Syn.*
274.
Hyoscyamus niger. *Ger. Em. 353. f. vulgaris. Baub.*
pin. 169. Park. 362.

Anglis,
 * By the road-side, on the left hand, as one ascends
 the hill in going from Enham to Andover, plen-
 tifully.

PENTANDRIA MONOGYNIA.

Anglis, common Henbane.

Habitat *ad Pagos, vias, fossas, ruderata.* ☉. VI.

Pharmac. Hyoscyami Radix, Semina, Herba.

D A T U R A. 218.

Cal. infundibul. plicata. *Cal.* tubulosus, angulatus, defiduus. *Capsf.* tetravalvis.

Stramonium. * I. DATURA pericarpis spinosis erectis ovatis. *Fl. suec.* 198. *Sp. pl.* 179.

Stramonium fructu spinoso oblongo, flore albo. *Tourn.* 119.

Solanum pomo spinoso oblongo, flore calathoide, Stramonium vulgo dictum. *R. Syn.* 266.

Solanum foetidum, pomo spinoso oblongo, flore albo. *Bauh. pin.* 164.

Stramonium spinosum. *Ger. Em.* 349.

Anglis, Thorny Apple.

Habitat *in ruderatis circa Londinum passim.* ☉. VII.

Pharmac. Stramonii Herba.

S O L A N U M. 224.

Cor. rotata. Antheræ subcoalitæ, apice poro gemino dehiscentes. *Bacca* bilocularis.

nigrum. I. SOLANUM caule inermi herbaceo, foliis ovatis dentato-angulatis, umbellis nutantibus. *Mat. med.* 94. *Fl. suec.* 200. *Sp. pl.* 186.

Solanum vulgare. *Park.* 346. *R. Syn.* 264.

Solanum officinarum. *Bauh. pin.* 166.

Solanum hortense. *Ger. Em.* 339. f. vulgare baccis nigris. *Bauh. hist.* III. 608.

β Solanum hortense f. vulgare baccis luteis. *Bauh. hist.* III. 608.

Anglis, Garden or common Nightshade.

Habitat *in ruderatis.* ☉. VI. VII.

Pharmac. Solani Herba, Fructus.

Dulcamara. 2. SOLANUM caule inermi frutescente flexuoso, foliis superioribus hastatis, racemis cymosis. *Mat. med.* 95. *Fl. suec.* 201. *Sp. pl.* 71.

Solanum

Solanum lignosum f. *Dulcamara*. *Park.* 350. *R. Syn.* 265.

Solanum scandens f. *Dulcamara*. *Baub. pin.* 167.

Amara Dulcis. *Ger. Em.* 350.

β. *Solanum lignosum* f. *Dulcamara marina*. *R. Syn.* 265. *Anglis*, Woody Nightshade, or Bitter-sweet.

Habitat in *sepibus humidis passim*. 4. VI. VII.

Pharmac. *Dulcamaræ*. *Radix, Folia, Cortex.*

A T R O P A. 222.

Cor. campanulata. *Stamina* distantia. *Bacca* globosa, bilocularis.

1. ATROPA caule herbaceo, foliis ovatis integris. *Sp. Balladonna.* *pl.* 181.

Balladonna. *Clus. Pann.* 504. *R. Syn.* 265.

Solanum lethale. *Park.* 346. *Ger. Em.* 340.

Solanum melanocerasus. *Baub. pin.* 166.

Anglis, Deadly Nightshade, or Dwale.

Habitat in *sylvis, sepibus, ruderatis*. In Chuckstone, near Rochester, in Kent; about Fulborn in Cambridgeshire, Sutton Cofield in Warwickshire. *R. Syn.* Among the Box going down to Darking, Surry. Dr. Wilmer. By the Road-side, under the Park Wall, between Temsford-mills and Welwyn, Hertfordshire. Dr. Watson. *Bl. spec.* 93. 4. VI. VII.

** *Flores monopetali superi.*

S A M O L U S. 205.

Cor. rotata. *Stamina* munita squamulis corollæ. *Capf.* unilocularis.

1. SAMOLUS. *Hort. cliff.* 51. *Fl. suec.* 192. *Sp. pl. valerandi.* 171.

Samolus Valerandi. *Baub. hist. III.* 791. *R. Syn.* 283.

Anagallis aquatica rotundo folio non crenato. *Baub. pin.* 283.

Anagallis aquatica rotundifolia. *Ger. Em.* 620.

Anagallis aquatica 3. *Lob. folio subrotundo non crenato.* *Park.* 1237.

Anglis, round-leaved Water-pimpernel.

Habitat

PENTANDRIA MONOGYNIA.

Habitat in palustribus, in pratis humidis inter Greenwich et Woolwich. 4. VI.

LONICERA. 210.

Cor. monopetala, irregularis. Bacca polysperma, bilocularis.

- Periclymenum.* 1. LONICERA capitulis ovatis imbricatis terminalibus; foliis omnibus distinctis. *Mat. med.* 68. *It. scan.* 305. *Fl. suec.* 193. *Sp. pl.* 173.
Caprifolium Germanicum. *Dod. pempt.* 411. *R. Syn.* 458.
Periclymenum. *Ger. Em.* 891.
Periclymenum f. *caprifolium* vulgare. *Park.* 1460.
Periclymenum non perfoliatum Germanicum. *Bauh. pin.* 302.
 β *Periclymenum* foliis quercinis. *Merr. pin. Anglis*, common Honeysuckle, or Woodbine.
 Habitat in sepibus. 2. V. VII.
Pharmac. *Caprifolii. Folia, Flores.*

CAMPANULA. 201.

Cor. campanulata fundo clauso valvis staminiferis. Stigma trifidum. Caps. infera, poris lateralibus dehiscens.

* *Foliis lævioribus angustioribus.*

- rotundifolia.* 1. CAMPANULA foliis radicalibus reniformibus, caulinis linearibus. *Fl. lapp.* 83. *Fl. suec.* 184. *Sp. pl.* 163. *Dalib. parif.* 66.
Campanula rotundifolia. *Ger. Em.* 452. *R. Syn.* 277.
Campanula minor sylvestris rotundifolia. *Park.* 651.
Campanula minor rotundifolia vulgaris. *Bauh. pin.* 93.
 β *Campanula minor rotundifolia alpina.* *Bauh. pin.* 93.
Anglis, lesser round-leaved Bell-flower.
 Habitat in ericetis et pascuis sterilibus frequens. 4. VIII.
patula. 2. CAMPANULA foliis strictis: radicalibus lanceolato-ovalibus, panicula patula. *Sp. pl.* 163.
Campanula esculentæ facie, ramis et floribus patulis.
Dill. elth. 68. t. 58. f. 68.
Anglis,

Anglis, Field Bell-flower.

Habitat in *sylvis et sepibus*. In Buddon wood, near Loughborough, in Leicestershire; Mr. Pultney. 4. VII. VIII.

3. CAMPANULA caule unifloro. *Fl. lapp.* 85. t. 9. *uniflora*.
f. 5. 6. *succ.* 185 *Sp. pl.* 163.

Anglis, Mountain Bell-flower.

Habitat in Wallia, D. Stillingfleet invenit. In montibus supra Grasmere, in Westmorlandia. 4. VII. VIII.

4. CAMPANULA foliis undulatis: radicalibus lanceo- *Rapuncu-*
lato-ovalibus panicula coarctata. *Hort. upf.* 40. *lus*.
Sp. pl. 164.

Rapunculus esculentus. *Bauh. pin.* 92. *R. Syn.* 276.

Rapunculus esculentus vulgaris. *Park.* 648.

Rapuntium parvum. *Ger. Em.* 453.

Anglis, Rampions.

Habitat in *aggeribus fossarum et arvis sed rarius*. Prope Croydon in agro Surriensi. 4. VIII.

** *Foliis scabris latioribus*.

5. CAMPANULA foliis ovato-lanceolatis, caule sim- *latifolia*.
plicissimo tereti, floribus solitariis pedunculatis,
fructibus cernuis. *Vir. Cliff.* 17. *Sp. pl.* 165. *Roy.*
lugdb. 245.

Campanula maxima foliis latissimis. *Bauh pin.* 94.
R. Syn. 276.

Trachelium majus Belgarum. *Park.* 643. five Gi-
ganteum. *Ger. Em.* 448.

Anglis, Giant Throatwort.

Habitat in *dumetis et sepibus*. In the mountainous part of Derbyshire, Staffordshire, Yorkshire, and elsewhere in the north of England plentifully. *R. Syn.* 4. VII. VIII.

6. CAMPANULA caule angulato, foliis petiolatis, ca- *Trache-*
lycibus ciliatis, pedunculis trifloris. *Fl. succ.* 189. *lium*.
Sp. pl. 166.

G

Cam-

PENTANDRIA MONOGYNIA.

Campanula vulgatio foliis *Urticæ*, vel major et asperior. *Baub. pin.* 94. *R. Syn.* 276.

Trachelium majus. *Ger. Em.* 448. flore purpureo. *Park.* 644.

Anglis, great Throatwort, or Canterbury Bells.

Habitat in sylvis et sepibus. 4. VII. VIII.

glomera-
ta.

7. *Campanula* caule angulato simplici, floribus sessilibus, capitulo terminali. *Fl. suec.* 190. *it. gotl.* 196. *it. scan.* 306. *Sp. pl.* 166.

Campanula pratensis flore conglomerato. *Baub. pin.* 94. *R. Syn.* 277.

Trachelium minus. *Park.* 644. *Ger. Em.* 449.

Anglis, lesser Throatwort, or Canterbury Bells.

Habitat in pascuis montosis et cretaceis frequens. 4. VII.

*** *Caule subdiviso.*

hybrida. 8. *CAMPANULA* caule basi subramoso stricto, foliis oblongis crenatis, calycibus aggregatis corolla longioribus, capsulis prismaticis. *Sp. pl.* 168.

Campanula arvensis erecta, vel *Speculum Veneris minus.* *Park.* 1331. *Ger. Em.* 439. *R. Syn.* 278.

Campanula arvensis minima erecta. *Hist. Ox.* II. 457. t. 2. f. 22.

Anglis, the lesser Venus Looking-glass, or coddled Corn-violet.

Habitat in arvis sed rarius. In arvis inter Northfleet et Stone, in Cantio, copiose. 6. VI. VII.

hederacea. 9. *CAMPANULA* foliis cordatis quinquelobis petiolatis glabris, caule laxo. *Dalib. parif.* 68. *Sp. pl.* 169.

Campanula Cymbalariæ foliis. *Ger. Em.* 452. *Park.* 652. *R. Syn.* 277.

Campanula Cymbalariæ foliis, vel folio hederaceo. *Baub. pin.* 93.

Anglis, tender Ivy-leaved Bell-flower.

Habitat in locis umbris humidis. On many moist and watery banks in the west of England, especially in Cornwall and Devonshire; I have found it also in the North, as about Sheffield in Yorkshire, Mr.

Lawson

Lawson. *In Bagley wood, near Oxford.* R. Syn.
In Wallia frequens. D. Stillingfleet. 4. V. VIII.

P H Y T E U M A. 203.

Cor. rotata laciniis linearibus, quinquepartita. *Stigma*
 trifidum. *Caps.* trilocularis.

1. PHYTEUMA capitulo subrotundo, foliis serratis; orbicularibus radicalibus cordatis. *Sp. pl.* 170. *ris.*

Rapunculus corniculatus montanus. Ger. Em. 455.
R. Syn. 278.

Rapunculus alopecuroides orbiculatus. Park. 648.

Rapunculus folio oblongo, spica orbiculari. Bauh.
pin. 92.

Anglis, horned Rampions, with a round head or spike
 of flowers.

Habitat in pascuis siccis. On the Downs of Suffex and
Hampshire. R. Syn. 4. VII.

Flores pentapetali inferi.

R H A M N U S. 235.

Cal. tubulosus: squamis stamina munientibus. *Cor.*
 nulla. *Bacca.*

1. RHAMNUS spinis terminalibus, floribus quadrifidis cathartico-
 dioicis. *Fl. juec.* 202. *Sp. pl.* 193. *cus.*

Rhamnus catharticus. Bauh. *pin.* 478. *hist.* I. 55.
R. Syn. 466.

Rhamnus solutivus. Ger. Em. 1337. f. *spina infec-*
toria vulgaris. Park. 243.

Anglis, Buckthorn, or purging Thorn.

Habitat in sylvis et sepibus. h. IV. V.

Pharmac. Spinæ Cervinæ *Baccæ.*

2. RHAMNUS inermis, floribus monogynis hermaproditis, foliis integerrimis. *Fl. juec.* 203. *Mat.*
med. 73. *Sp. pl.* 193. *Frangula.*

Frangula seu Alnus nigra baccifera. Park. 240. R.
Syn. 465.

Alnus nigra baccifera. Bauh. *pin.* 428. *hist.* I. 560.

Alnus nigra sive Frangula. Ger. Em. 1469.

Anglis, the black berry-bearing Alder.
Habitat in sylvis et sepibus. h. IV. V.
Pharmac. Frangulæ Cortex.

EUONYMUS. 240.

Cor. pentapetala. *Caps.* pentagona, quinquelocularis, quinquevalvis, colorata. *Sem.* calyptrata.

europæus. 1. EUONYMUS floribus plerisque quadrifidis. *Fl. suec.* 204. *Sp. pl.* 197.

Euonymus foliis oblongo-ovatis. *Hort. cliff.* 38.

Euonymus vulgaris. *Park.* 241. *R. Syn.* 468.

Euonymus vulgaris granis rubentibus. *Bauh. pin.* 428.

Euonymus Theophrasti. *Ger. Em.* 1468.

Anglis, Spindletree, or Prickwood.

Habitat in sepibus et sylvis. h. IV. V.

** *Flores pentapetali superi.*

RIBES. 247.

Petala quinque et *stamina* calyci inserta. *Stylus* bifidus. *Bacca* polysperma.

rubrum. * 1. RIBES inerme, racemis glabris pendulis, floribus planiusculis. *Hort. cliff.* 82. *Mat. med.* 99. *Sp. pl.* 200. *Fl. suec.* 205.

Ribes vulgaris fructu rubro. *Ger. Em.* 1593. *R. Syn.* 456.

Ribes fructu rubro. *Park.* 1561.

Grossularia sylvestris rubra. *Bauh. pin.* 455.

β *Grossularia vulgaris* fructu dulci. *Bauh. pin.* 455.

γ *Ribes* fructu parvo. *Merr. pin.*

Anglis, Currants.

Habitat in sylvis et sepibus. In sylvis in septentrionali comitatus Eboracensis parte, inque Dunelmensi Episcopatu et Westmorlandia, D. Johnson. *R. Syn.* h. V.

Pharmac. Ribium Baccæ.

alpinum. 2. RIBES inerme, racemis erectis: bracteis flore longioribus. *Sp. pl.* 200.

Ribes alpinus dulcis. *Bauh. hist.* II. 98.

Anglis,

Anglis, sweet Mountain-currants.

Habitat in sylvis et sepibus. In agro Eboracensi invenit, D. Dodsworth; in sepibus-juxta Bradford abundat, D. Richardson. R. Syn. 4. V.

3. RIBES inerme, racemis pilosis floribus oblongis. *Sp. nigrum.*
pl. 201. *Fl. suec.* 207.

Ribes nigrum vulgo dictum folio olente. *Baub. hist.* II.
98. R. Syn. 456.

Ribes fructu nigro. *Park.* 1562. *Ger. Em.* 1593.

Grossularia non spinosa fructu nigro. *Baub. pin.* 455.

Anglis, black Currants, or Squinancy Berries.

Habitat in sylvis humidis et ad ripas fluviorum. We observed it at Abington in Cambridgeshire, Blunham in Bedfordshire, in Warwickshire, Westmoreland and Cumberland. By the Hoppet-bridge near Braintree. R. Syn. 2. V.

H E D E R A. 249.

Petala quinque, oblonga. Bacca quinquesperma, calyce cincta.

1. HEDERA foliis ovatis lobatisque. *Fl. lapp.* 91. *Sp. Helix.*
pl. 75.

Hedera communis major et minor. *Baub. hist.* II.

III.

Hedera arborea. *B. pin.* item major sterilis. *Baub. pin.* 305.

Hedera arborea five scandens et corymbosa communis.
Park. 678.

Hedera helix. *Ger. Em.* 858.

Anglis, Ivy.

Habitat in sylvis et sepibus. 2. X.

Pharmac. Hederæ Folia, Baccæ, Gummi.

* Flores incompleti superi.

ILLECEBRUM. 256.

Cal. pentaphyllus, cartilagineus. Cor. o. Stigma simplex. *Caps.* quinquevalvis, monosperma.

1. ILLECEBRUM floribus verticillatis nudis, caulibus *verticillatum.*
procumbentibus. *Sp. pl.* 206.

PETNANDRIA MONOGYNIA.

Corrigiola. Cat. giff. app. suppl. 169. *R. Syn.* 160.
Polygonum serpillifolium verticillatum. R. Syn. ed. I.
 160.

Polygonum parvum flore albo verticillato. Bauh. hist.
III. 378.

Polygala repens. Park. 1333. *nivea. Bauh. pin.* 215.
Anglis, verticillate Knot-grass.

Habitat in palustribus udis et humidioribus pascuis circa
Pensans oppidum et alibi versus extremum Cornu-
biæ angulum. R. Syn. 24. VII.

G L A U X. 257.

Cal. monophyllus. Cor. o. Caps. unilocularis; quin-
quevalvis, quinquesperma.

maritima. I. GLAUX. Hort. cliff. 43. *Fl. suec.* 210. *Sp. pl.*
 207.

Glaux foliis elliptico-oblongis. Fl. lapp. 72.

Glaux maritima. Bauh. pin. 215; *R. Syn.* 285. *mi-*
nor. Park. 1283.

Glaux exigua maritima. Ger. Em. 562.

Anglis, Sea Milkwort, or black Saltwort.

Habitat in littoribus maritimis. 24. VII.

*** Flores incompleti inferi.*

T H E S I U M. 258.

Cal. monophyllus, cui stamina inserta. Sem. unicum.

Linophyl. 2. THESIUM panicula foliata; foliis linearibus. Fl. suec.
lon. 211. *Sp. pl.* 207.

Linaria adulterina. Ger. Em. 555; *R. Syn.* 202.

Linaria montana flosculis albicantibus. Bauh. pin.
 213.

Pseudo linaria montana alba. Park. 459.

Anglis, bastard Toad-flax.

Habitat in pascuis montosis præsertim solo cretaceo. As
upon Gogmagog hills, and Newmarket heath. R. Syn.
prope Basingstoke in Hampshire. 24. VI. VII.

D I G Y N I A.

* *Flores monopetali inferi.*

S W E R T I A. 284.

Cor. rotata. Nectariferi pori ad basin laciniarum corollæ. Caps. unilocularis, bivalvis.

1. SWERTIA corollis quinquefidis, foliis radicalibus *perennis.*
ovalibus. Sp. pl. 226.

Gentiana palustris latifolia. Bauh. pin. 188.

Gentiana XII. punctato flore. Clus. hist. I. 316.

Gentiana Pennei minor. Ger. Em. 433.

Anglis, Marsh Gentian.

Habitat in Wallia, D. Richardson invenit. 4. VIII.

G E N T I A N A. 285.

Cor. monopetala. Caps. bivalvis, unilocularis. Receptaculis duobus longitudinalibus.

* *Corollis quinquefidis subcampaniformibus.*

1. GENTIANA corollis quinquefidis campanulatis op- *Pneumopositis pedunculatis, foliis linearibus. Sp. pl. 228. nanthe.*
Fl. suec. 228.

Gentiana floribus terminalibus raris, corollis erectis plicatis, foliis linearibus. Hort. cliff. 80.

Gentiana palustris angustifolia. Bauh. pin. 188. R. Syn. 274.

Gentiana autumnalis Pneumonanthe dicta. Park. 406.

Pneumonanthe. Ger. Em. 438.

Anglis, Calathian Violet.

Habitat in pratis humidioribus. In pratis Strattoniensi, D. Stillingfleet invenit; prope Tunbridge, Dr. Wilmer; in palustribus prope Milthorp in comitatu Westmorlandico copiose. 4. VIII.

** *Corollis quinquefidis infundibuliformibus.*

2. GENTIANA corollis quinquefidis hypocrateriformi- *Amarella.*
bus fauce barbatis. Sp. pl. 230. Fl. suec. 229.

PENTANDRIA DIGYNIA.

Gentianella pratensis flore lanuginoso. *Bauh. pin.* 188.
R. Syn. 27c.

Gentianella fugax autumnalis elatior, centaurii minoris foliis. *R. Syn.* 275.

Gentianella fugax minor. *Ger. Em.* 437.

Anglis, Autumnal Gentian, or Fellwort.

Habitat in pascuis siccioribus. ☉. VII. VIII.

Centaureum. 3. *GENTIANA* corollis quinquefidis infundibuliformibus, caule dichotomo. *Sp. pl.* 229.

Centaureum minus. *Bauh. pin.* 278. *R. Syn.* 286.

Centaureum minus flore purpureo et albo. *Bauh. hist.*

III. 353.

Centaureum minus vulgare. *Park.* 272.

Centaureum parvum. *Ger. Em.* 547.

Anglis, lesser Centory.

Habitat in pascuis siccioribus et sterilioribus. ☉. VI. VIII.

Pharm. Centaurii minoris Herba, Flores.

*** *Corollis quadrifidis.*

campestris. 4. *GENTIANA* corollis quadrifidis fauce barbatis. *Sp. pl.* 231. *Fl. suec.* 230.

Gentiana corollo hypocrateriformi, tubo villis clauso, calycis foliis alternis majoribus. *Fl. lapp.* 94.

Gentianella fugax verna seu præcox. *R. Syn.* 275.

Gentianella alpina verna minor. *Bauh. pin.* 188.

Gentiana purpurea minima. *Col. ecph. I.* 232. t. 221.

Anglis, vernal dwarf Gentian.

Habitat in pascuis montosis. Found by Mr. Fitz-Roberts on the back side of Halsfellnab, near Kendal; as also in the parks on the other side of Kendal, and on the back of Birkhag. *R. Syn.* ☉. VII.

filiformis. 5. *GENTIANA* corollis quadrifidis imberbibus, caule dichotomo filiformi. *Sp. pl.* 231.

Gentiana caule dichotomo, foliis lineari-lanceolatis, floribus infundibuliformibus quadrifidis longissime pedunculatis. *Guett. stamp. I.* 305. *Dalib. parif.* 82.

Centau-

PENTANDRIA DIGYNIA.

39

Centaurium palustre luteum minimum. *R. Syn.* 286.

Vaill. parif. 32. t. 6. f. 3.

Anglis, Marth Centory.

Habitat in palustribus. I found it towards the end of

Cornwal in rotten marshy ground. R. Syn. ☉. VII.

C U S C U T A. 156.

Cal. quinquefidus. Cor. monopetala quinquefida.

Capf. bilocularis. Sem. duo.

1. CUSCUTA floribus sessilibus. *Sp. pl.* 124. *Fl. suec. europæa.* 144.

Cuscuta major. Baub. pin. 219. R. Syn. 281. f. Caf-

futha. Ger. Em. 577. Park. 10.

Anglis, Dodder, Hellweed, or Devil's Guts.

Habitat in arvis et ericetis passim. ☉. VII.

Pharm. Cuscutæ f. Epithymi Herba.

* *Flores incompleti.*

C H E N O P O D I U M. 273.

Cal. pentaphyllus, pentagonus. Cor. nulla. Sem.

unicum lenticulare, superum.

* *Foliis angulosis.*

1. CHENOPODIUM foliis triangulari-sagittatis integerrimis. *Fl. suec.* 214. *Sp. pl.* 218. *Bonus
Henricus.*

Blitum perenne Bonus Henricus dictum. Hist. Ox. II.

599. t. 30. f. 1. R. Syn. 156.

Bonus Henricus. Baub. hist. II. 965. Ger. Em. 329.

Lapathum unctuosum. Park. 1225.

Lapathum sylvestre latifolium l. seu unctuosum folio triangulo. Baub. pin. 115.

Anglis, common English Mercury, or Allgood.

Habitat in ruderatis, et ad vias, muros, agros passim.

4. VIII.

Pharmac. Boni Henrici Herba.

2. CHENOPODIUM foliis triangularibus subdentatis, *urbicum.* racemis confertis strictissimis caule approximatis longissimis. *Fl. suec.* 215. *Sp. pl.* 218.

Che-

PENTANDRIA DIGYNIA.

Chenopodium latifolium minus ramosum, petiolatis longissimis exfoliorum alis confertim nascentibus.
Buxb. ball. 69. t. 1.

Chenopodium erectum, foliis triangularibus dentatis, spicis e foliorum alis plurimis longis erectis tenuibus.
R. Syn. 155.

Anglis, upright Blite.

Habitat in ruderatis. ☉. VIII. IX.

rubrum. 3. *CHENOPODIUM* foliis cordato-triangularibus obtusiusculis dentatis, racemis erectis compositis subfoliosis caule brevioribus. *Syst. Nat. 948. Fl. suec. 217. Sp. pl. 218.*

Chenopodium erectum, foliis subtriangularibus antice sinuato-dentatis, racemis erectis foliosis lateralibus.
Hort. cliff. 85. Roy. lugdb. 219.

Blitum Pes anserinus acutiore folio. R. Syn. 154.

Atriplex sylvestris latifolia acutiore folio. Baub. pin. 119.

Atriplex sylvestris latifolia altera. Ger. Em. 328.

Anglis, sharp-leaved Goosefoot.

Habitat in fimetis et ruderatis frequens. ☉. VIII.

murale. 4. *CHENOPODIUM* foliis ovatis acutis dentatis, racemis ramosis nudis. *Fl. suec. 216. Sp. pl. 219.*

Blitum Pes anserinus dictum. R. Syn. 154.

Atriplex sylvestris latifolia. Baub. pin. 119.

Pes anserinus. Ger. Em. 328. Park. 749.

β *Blitum Morifono atriplex procumbens folio sinuato lucido crasso dictum. Hist. Blæs. R. Syn.*

Anglis, common Goosefoot, or Sowbane.

Habitat in locis cultis, ruderatis frequens. ☉. VIII.

hybridum. 5. *CHENOPODIUM* foliis cordatis angulato-acuminatis, racemis ramosis nudis. *Fl. suec. 220. Sp. pl. 219.*

Chenopodium foliis triangulari-sagittatis: infra medium sinuato-dentatis, racemis longissimis. Hort. cliff. 84. Roy. lugdb. 219.

Chenopodium Stramonii folio. R. Syn. 154. Vaill. parif. 36. t. 7. f. 2.

Cheno-

Chenopodio affinis, folio lato, laciniato, in longissimum mucronem procurrente, florum ramulis sparsis. *R. hist. III.* 123.

β Blito Pes anserinus dicto similis. *R. Syn.* 154.

Anglis, Maple-leaved Blite.

Habitat in ruderatis, fimetis passim. ☉. VIII.

6. CHENOPODIUM foliis rhomboideo-triangulis ero. *album*.
 sis postice integris: summis oblongis, racemis erectis. *Fl. suec.* 218. *Sp. pl.* 219.

Chenopodium foliis inferioribus ovatis acutis antrorsum dentatis: summis lanceolatis. *Hort. cliff.* 85.

Chenopodium sylvestre, opuli folio. *Vaill. paris.* 36.
t. 7. *f.* 1.

Blitum atriplex sylvestre dictum. *R. Syn.* 154.

Atriplex folio sinuato candicante. *Bauh. pin.* 119.

Atriplex sylvestris vulgarior sinuata major. *Park.* 748.

Atriplex vulgaris. *Ger. Em.* 326.

β Blitum folio subrotundo. *R. Syn.* 155.

Anglis, common Orache.

Habitat in fimetis, cultis, ruderatis. ☉. VIII.

7. CHENOPODIUM foliis rhomboideis dentato-sinu. *viride*.
 atis, racemis ramosis subnudis. *Fl. suec.* 219. *Sp.*
pl. 219.

Chenopodium folio oblongo integro. *Dill. cat. giff.*
 154.

Chenopodium foliis integris racemosum. *D. Dale. R.*
Syn. 155.

β Chenopodium crasso et obtuso oleæ folio. *R. Syn.* 156.

Anglis, green Blite.

Habitat in cultis passim. ☉.

8. CHENOPODIUM foliis deltoideis sinuato-dentatis *serotinum*.
 rugosis glabris uniformibus, racemis terminalibus.
Cent. pl. II. Syst. Nat. 948.

Blitum ficus folio. *Pet. H. Br. t.* 8. *f.* 3. *R. Syn.*
 155.

Atriplex vulgaris sylvestris folio oblongo angusto magno,
 utrinque sinu donato, fructu multo racemoso. *Buddl.*
H. S. IV. 38.

Anglis,

Anglis, late flowered Blite.

Habitat in *fimeticis ruderalis passim*. ☉. VIII. IX.

- glaucum*. 9. CHENOPODIUM foliis ovato-oblongis repandis, racemis nudis simplicibus glomeratis. *Fl. suec.* 221. *Sp. pl.* 220.
 Chenopodium angustifolium laciniatum minus. *Tourn.* 506. *R. Syn.* 155.
 Blitum procumbens folio botryoide subtus incano. *D. Rand.*
 Blitum quercus folio. *Pet. H. Br. t. 8. f. 1.*
Anglis, Oak-leaved Blite.
 Habitat in *ruderalis frequens*. ☉. VIII.

** *Foliis integris.*

- Vulvaria*. 10. CHENOPODIUM foliis rhombeo-ovatis integerrimis floribus conglomeratis. *Fl. suec.* 222. *Mat. med.* 108. *Sp. pl.* 220.
 Blitum foetidum Vulvaria dictum. *R. Syn.* 156.
 Atriplex olida. *Ger. Em.* 327. five sylvestris foetida. *Park.* 749.
 Atriplex foetida. *Baub. pin.* 119. *hist. II.* 974.
Anglis, stinking Orache.
 Habitat in *ruderalis et ad vias, et muros, frequens*. ☉. VIII.

- polypermum*. 11. CHENOPODIUM foliis ovatis integerrimis caule decumbente cymis dichotomis. *Fl. suec.* 223. *Sp. pl.* 220.
 Chenopodium Betæ folio. *Tourn.* 506. *R. Syn.* 157.
 Blitum polypermum. *Baub. pin.* 118. *Park.* 753.
Anglis, round-leaved Blite, or Allseed.
 Habitat in *cultis, fimeticis ruderalis passim*. ☉.

- maritimum*. 12. CHENOPODIUM foliis subulatis semicylindricis. *Sp. pl.* 221.
 Blitum Kali minus album dictum. *R. Syn.* 156.
 Kali minus. *Ger. Em.* 535. album *Park.* 749.
 Kali minus album semine splendens. *Baub. pin.* 289.
Anglis, Sea Blite, or white Glasswort.
 Habitat in *littoribus maritimis frequens*. ☉. VIII.

13. CHE-

13. CHENOPODIUM foliis linearibus teretibus car-*frutico-*
nos, caule fruticoso. *Hort. cliff.* 806. *Sp. pl.* 221. *sum.*
 Blitum fruticosum maritimum, Vermicularis frutex
 dictum. *R. Syn.* 156.
 Sedum minus fruticosum. *Baub. pin.* 284.
 Vermicularis frutex minor. *Ger. Em.* 523.
 Vermicularis fruticosa altera. *Park.* 731.
Anglis, Shrub Stonecrop, or Glasswort.
 Habitat in littoribus maritimis Norfolchiæ, Devonix, *Devonix*,
 Cornubiæ, &c. *h.* VIII.

S A L S O L A. 275.

- Cal.* pentaphyllus. *Cor.* o. *Caps.* monosperma. *Sem.*
 cochleatum.
 1. SALSOLA herbacea decumbens, foliis subulatis spi- *Kali.*
nos scabris, calycibus marginatis exillaribus. *Syst.*
Nat. 949.
 Salsola herbacea foliis subulatis mucronatis, calycibus
 ovatis exillaribus. *Sp. pl.* 222. *Fl. suec.* 225.
 Kali spinosum cochleatum. *Baub. pin.* 289. *R. Syn.*
 159.
Anglis, prickly Glasswort.
 Habitat in littoribus maritimis *passim.* ☉. VII. VIII.

B E T A. 274.

- Cal.* pentaphyllus. *Cor.* o. *Sem.* reniforme, intra
 substantiam baseos calycis.
 1. BETA. *Hort. cliff.* 83. *Mat. med.* 113. *Roy. lugdb.* *vulgaris.*
 220. *Sp. pl.* 222.
 Beta sylvestris maritima. *Baub. pin.* 157. *Park.*
 750.
 Beta alba et rubra. *Ger. Em.* 319.
Anglis, Sea Beet.
 Habitat in littoribus maritimis *ubique.* ☿. VIII.
Pharmac. Betæ rubræ et albæ Radix, Folia.

HER-

HERNIARIA. 272.

Cal. quinquepartitus. *Cor.* nulla. *Stamina* quinque
sterilia. *Caps.* monosperma.

glabra. 1. *HERNIARIA glabra.* *Bauh. hist. III.* 378. *Sp. pl.*
218.

Herniaria. *Ger. Em.* 569. *R. Syn.* 160.

Polygonum minus, seu Millegrana major. *Bauh. pin.*
281.

Millegrana major, seu Herniaria vulgaris. *Park.* 446.
Anglis, smooth Rupture-wort.

Habitat in glareosis. In promontorio Cornubiensi, the
Lizard-point dicto Herniariam glabram copiosam in-
veni. ☉. VII.

hirsuta. 2. *HERNIARIA hirsuta.* *Bauh. hist. III.* 379. *R. Syn.*
161.

Anglis, rough Rupture-wort.

Habitat in pratis glareosis sed rarius in pratis juxta
Colney-hatch prope Barnet. ☉. VII. VIII.

lenticu- 3. *HERNIARIA.*
lata.

Polygonum maritimum longius radiculatum nostras, ser-
pyllifolio circinato crasso nitente. *R. Syn.* 161.

Polygonum minus lentifolium. *Bauh. pin.* 282. *pr.*
131.

Polygonum minus monspeliense. *Park.* 446. *Pluk.*
ph. t. 53. *f.* 3.

Polygonum folio circinata. *Pet. h. br. t.* 10. *f.* 6.
Anglis, Sea Rupture-wort.

Habitat in littoribus maritimis. In littore Sabriniano,
non procul a Weston supra mare. *D. Pluk. R.*
Syn. 4.?

ULMUS. 281.

Cal. quinquefidus. *Cor.* o. *Bacca* exsucca com-
presso-membranacea.

campestris 1. *ULMUS foliis ovatis duplicato-ferratis basi inæqua-*
libus. *Sp. pl.* 225.

Ulmus vulgarissima folio lato scabro. *Ger. Em.* 1480.
R. Syn. 468.

Ulmus

Ulmus campestris et *Theophrasti*. *Bauh. pin.* 426.

Ulmus vulgaris. *Park.* 1404.

β *Ulmus minor folio angusto scabro*. *Ger. Em.* 1480.
R. Syn. 469.

Anglis, common Elm.

Habitat in sepibus. h. IV.

Pharm. *Ulmī Folia, cortex medius*.

2. *ULMUS* foliis oblongo-ovatis duplicato-ferratis basi *glabra*.
inæqualibus, cortice glabro.

Ulmus folio latissimo scabro. *Ger. Em.* 1481. *R.*
Syn. 469.

Ulmus latiore folio. *Park.* 1404.

β *Ulmus folio glabro*. *Ger. Em.* 1481. *Park.* 1404.
Anglis, broad-leaved Elm, or Wych-hasel.

Habitat in sepibus. h. IV.

* *Flores pentapetali superi dispermi. Umbellatæ.*

A. Involucro universali partialique.

ERYNGIUM. 287.

Flores capitali. Receptaculo paleaceo.

1. *ERYNGIUM* foliis radicalibus subrotundis plicatis *mariti-*
spinosis, capitulis pedunculatis. It. oel. 151. *Fl. mum.*
suæc. 233. *Sp. pl.* 233.

Eryngium maritimum. *Bauh. pin.* 386.

Eryngium marinum. *Ger. Em.* 1162. *Park.* 986.
R. Syn. 222.

Anglis, Sea holly, or Eryngo.

Habitat in littoribus maritimis. 4. VII. VIII.

Pharmac. *Eryngii Radix*.

2. *ERYNGIUM* foliis radicalibus pinnatis tripartitis. *campestre.*
Sp. pl. 233.

Eryngium vulgare. *Bauh. hist.* III. 85. *pin.* 386. *R.*
Syn. 222.

Eryngium mediterraneum. *Ger. Em.* 1162. f. *cam-*
pestre. *Park.* 986.

Anglis, common Eryngo.

Habitat in pascuis maritimis. On a rock which you de-
scend to the ferry from Plymouth over into Cornwall;
it

PENTANDRIA DIGYNIA.

it was also sent me by Mr. Thornton, who observed it not far from Daventry in Northamptonshire, beside the old Roman Way, called Watling street, near a Village called Brookhall; observed by Mr. Lawson on the shore called Friar goose, near Newcastle upon Tyne. R. Syn. Below Melling in Yorkshire plentifully. Blackf. spec. bot. 4. VII. VIII.

HYDROCOTYLE. 288.

Umbellæ simplex: Involucro tetraphyllo. Petala integra. Sem. semiorbiculato compressa.

vulgaris. I. HYDROCOTYLE foliis peltatis, umbellis quinquefloris. *Sp. pl.* 234.

Hydrocotyle foliis peltatis orbiculatis undique emarginatis. Hort. cliff. 88. *Fl. suec.* 234.

Hydrocotyle vulgaris. Tour. 328. *R. Syn.* 222.

Cotyledon palustris. Ger. Em. 529. *Park.* 1214.

Ranunculus aquaticus, Cotyledonis folio. Bauh. pin. 180.

Anglis, Marsh Pennywort, or White-rot.

Habitat in palustribus. 4. V.

SANICULA. 289.

Umbellæ confertæ, subcapitatæ. Fructus scaber. Flores disci abortientes.

europæa. I. SANICULA foliis radicalibus simplicibus, flosculis omnibus sessilibus. *Hort. cliff.* 90. *Fl. suec.* 235. *Sp. pl.* 235.

Sanicula officinarum. Bauh. pin. 319.

Sanicula f. Diapensia. Ger. Em. 948. *R. Syn.* 221.

Sanicula vulgaris five Diapensia. Park. 532.

Anglis, Sanicle.

Habitat in sylvis et sepibus. Sylva Charlton wood dicta in Cantio. 4. V. VI.

Pharmac. Saniculæ Herba.

B U P L E U R U M. 291.

Involucrum umbellula majora, pentaphylla. *Fructus* subrotundus, compressus, striatus.

1. BUPLEURUM involucris universalibus nullis, foliis *rotundifolium*. *Hort. upf.* 64. *Sp. pl.* 236.

Bupleurum perfoliatum rotundifolium annuum. *Tourn.* 310. *R. Syn.* 221.

Perfoliata vulgaris. *Ger. Em.* 536. *Park.* 580.

Perfoliata vulgarissima seu arvensis. *Baub. pin.* 272. *Anglis*, Thorow-wax.

Habitat in arvis. Circa Lewisham in Cantio, D. Alchorne reperit. ☉. VII. *in a field near Andover, bordering Pharmac.* Perfoliatæ Herba. *Upon long mead & Shepherd's spring.*

2. BUPLEURUM umbellis simplicibus alternis pentaphyllis subtrifloris. *Sp. pl.* 238. *tenuissimum.*

Bupleurum foliis linearibus acutis sessilibus. *Hort. cliff.* 104.

Bupleurum minimum. *Col. Ec. I.* 247. *Park.* 587. *R. Syn.* 221.

Bupleurum angustissimo folio. *Baub. pin.* 278. *Anglis*, the least Hares-ear.

Habitat in pascuis sed rarius. Near Ellesley, in the road from Cambridge to St. Neots; on a bank by the northern road, a little beyond Huntington; at Malden in Essex, in the marshes by the river side, plentifully; at Hastings in Suffex; and near the ferry in the isle of Thanet. *R. Syn.* ☉. VII. VIII.

E C H I N O P H O R A. 292.

Fructus pedunculis immerfi. *Flores* varii abortientes.

1. ECHINOPHORA foliolis subulato-spinosis integerimis. *Sp. pl.* 239. *spinosa.*

Echinophora maritima spinosa. *Tourn.* 656. *R. Syn.* 220.

Crithmum maritimum spinosum. *Baub. pin.* 288. seu *Pastinaca marina.* *Park.* 1286.

Crithmum spinosum. *Ger. Em.* 533.

Anglis, prickly Sampire, or Sea Parsnep.

Habitat in littoribus maritimis. Observed by Mr. Law-son

PENTANDRIA DIGYNIA.

son at Roofsbeck in Low-Fournies, Lancashire ;
on the sea beach, in the way from Feversham to Sea-
Salter. *Blackf. sp. bot.* 4. VII.

TORDYLIUM. 293.

Corolla radicatæ, omnes hermaphroditæ. *Fructus*
suborbiculatis, margine crenatus. *Involucra* longa
indivisa.

*latifoli-
um.*

1. TORDYLIUM umbella conferta nudiuscula, foliis
pinnatis; foliolis lanceolatis inciso-ferratis. *Sp. pl.*
240.

Caucalis umbella universali trifida, partialibus penta-
spermis, foliis pinnatis ferratis. *Hort. cliff.* 91.
Roy. lugdb. 95.

Caucalis arvensis echinata latifolia. *Baub. pin.* 152.
R. Syn. 219.

Caucalis apii foliis, flore rubro. *Ger. Em.* 1021.

Echinophora arvensis latifolia purpurea. *Park.* 920.

Anglis, purple flowered great bastard Parsley.

Habitat inter segetes in agro Cantabrigiensi et alibi.

R. Syn. ☉. VI. VII.

nodosum.

2. TORDYLIUM umbellis simplicibus sessilibus, fe-
minibus exterioribus hispidis. *Sp. pl.* 240.

Caucalis nodosa echinato semine. *Baub. pin.* 153.

Ger. Em. 1022. *Park.* 921. *R. Syn.* 220.

Anglis, knotted Parsley.

Habitat ad margines agrorum frequens. ☉ V.

CAUCALIS. 294.

Cor. radiatæ, disci masculæ. *Fructus* fetis hispidus.
Involucra integra.

arvensis.

1. CAUCALIS umbella conferta, involucro universali
monophyllo, feminibus oblongis, foliolis ovato lan-
ceolatis pinnatifidis.

Caucalis segetum minor, Anthriscio hispido similis. *R.*
hist. I. 468. *syn.* 220.

Caucalis pumila segetum Goodyero. *Ger. Em.* 1022.
Anglis, small corn Parsley.

Habitat in arvis frequens. ☉. VIII.

2. CAU-

2. CAUCALIS umbella conferta, feminibus globosis, *Anthriscus*.
foliolis ovatis pinnatifidis.

Tordylium umbella conferta, foliolis ovato-lanceolatis
pinnatifidis. *Sp. pl.* 240. *Fl. suec.* 236.

Caucalis minor flosculis rubentibus. *Ger. Em.* 1022.
R. Syn. 219.

Caucalis minor flore rubentē. *Park.* 921.

Caucalis femine aspero, flosculis rubentibus. *Bauh.*
pin. 153.

Anglis, hedge Parsley.

Habitat in *sepibus*. ♂. VIII.

3. CAUCALIS feminum aculeis triglochidibus uncina-*leptophyll-*
tis: pilis verticillatis hispidis. *Guett. stamp.* 80. *la.*
Dalib. parif. 85. *Sp. pl.* 242.

Caucalis tenuifolia, flosculis subrubentibus. *R. hist.* 1.
697. *syn.* 219.

Caucalis arvensis echinata parvo flore et fructu. *Bauh.*
pin. 152.

Caucalis tenuifolia purpurea. *Park.* 920.

Anglis, fine-leaved bastard Parsley.

Habitat in *arvis circa Kingston in agro Cantabrigienfis*.
R. Syn. ☉. VII.

DAUCUS. 296.

Corollæ subradiatæ, omnes hermaphroditæ. *Fructus*
pilis hispidis.

1. DAUCUS feminibus hispidis. *Hort. cliff.* 88. *Sp. pl.* *Carota*.
242.

Daucus vulgaris. *Clus. hist.* 198. *R. Syn.* 218.

Pastinaca tenuifolia sylvestris. *Bauh. pin.* 155. *Ger.*
Em. 1028. *Park.* 902.

β *Daucus sativus* radice lutea et alba. *Tourn.* 307.

γ *Daucus sativus* radice atro-rubente. *Tourn.* 307.

♂ *Daucus maritimus* lucidus. *Tourn.* 307.

Anglis, wild Carrot, or Bird's-nest.

Habitat in *pratis et pascuis ubique*. ♂. VI. VII.

Pharmac. *Dauci sylvestris semina*.

PENTANDRIA DIGYNIA.

C O N I U M. 299.

Fruetus subglobosus, quinquestriatus, utrinque crenatus.

macula- 1. CONIUM feminibus striatis. *Hort. upf.* 92. *Roy.*
tum. *lugdb.* 107. *Sp. pl.* 243.

Cicuta major. *Bauh. pin.* 160.

Cicuta. *Lob. ic.* 732. *Ger Em.* 1061. *Dorfl.* 83.
Fuchf 406. *Dodon. pempt.* 461. *Gesn. Coll.* 68.

Cicuta vera. *Gesn. Hort.* 253.

Cicuta vulgaris major. *Park.* 933. *Clus. hist.* II. 200.
Anglis, Hemlock.

Habitat ad maceries et sepes frequens. ☉. VI. VII.

Pharmac. *Cicutæ herba.*

A T H A M A N T A. 301.

Fruetus ovato-oblongus striatus.

Libano- 1. ATHAMANTA foliis bipinnatis planis, umbella he-
tis. mispherica, feminibus hirsutis. *Sp. pl.* 244.

Apium petræum seu montanum album. *Bauh. hist.*
III. 105. *R. Syn.* 219.

Daucus montanus apii folio albicans. *Bauh. pin.* 157.

Daucus montanus pimpinellæ saxifragæ hircinæ folio
nostras Dubrienfis. *Pluk. ph. t.* 173. *f. 1.*

Anglis, mountain stone Parsley.

Habitat in pratis siccis. On Gogmagog hills in Cam-
bridgehire, inter St. Albans et Stoney-Stratford.
24. VIII. IX.

Meum. 2. ATHAMANTA foliolis capillaribus, feminibus stri-
atis. *Hort. cliff.* 93. *upf.* 60. *Mat. med.* 126. *Sauv.*
monsp. 182. *Sp. pl.* 245.

Meum foliis anethi. *Bauh. pin.* 148.

Meum. *Ger. Em.* 1052. *R. Syn.* 207. *vulgatius.*
Park. 888.

Anglis, common Spignel, Meu, Bald or Bawd-money.

Habitat in pratis montosis in comitatibus Westmor-
landico, Eboracensi, Lancastriensi, Cumberlandico,
frequens. It is found near Dolgehle in Merioneth-
shire. *R. Syn.* 24. V.

Pharmac. *Athamantæ Radix.*

P E U C E D A N U M. 302.

Fruſtus ovatus, utrinque ſtriatus, ala cinctus. *Involucra* breviffima.

1. PEUCEDANUM foliis quinquies tripartitis filiformi-*officinale.*
bus linearibus. *Sp. pl.* 245.

Peucedanum foliis quinquies tripartitis lineari-subulatis integerrimis. *Hort. cliff.* 93. *Mat. med.* 126.

Peucedanum. *Ger. Em.* 1054. *K. Syn.* 206. Germanicum. *Bauh. pin.* 146.

Peucedanum vulgare. *Park.* 880.

Anglis, Hogs-fennel, Sulphur-wort, Hareſtrong.

Habitat in paluſtribus maritimis. *Prope* Shoreham in Suffexia, Walton non procul ab Harvico in Eſſexia; about a quarter of a mile below Feversham, by the river ſide, Mr. J. Sherard. *R. Syn.* 4. VI.

Pharmac. Peucedani Radix.

2. PEUCEDANUM foliis pinnatis, foliolis pinnatifidis: laciniis linearibus oppoſitis, caule ramiſſiſſimo patulo. *minus.*

Peucedanum pumilum. *Ger. Em.* 1054.

Peucedanum minus. *Park.* 880. *Bauh. pin.* 149. *R. Syn.* 217.

Selinum montanum pumilum Cluſii, foliis Foeniculi aut Peucedani, flore albo, ſemine Selini. *Bauh. hiſt.* III. 17.

Anglis, Rock Parſley.

Habitat in paſcuis montoſis. On St. Vincent's rock near Briſtol. *R. Syn.* 8. V.

C R I T H M U M. 303.

Fruſtus ovalis, compreſſus.

1. CRITHMUM foliolis lanceolatis carnoſis. *Sp. pl.* *maritimum.* 246.

Crithmum marinum. *Ger. Em.* 553. *Dod. p.* 705. *R. Syn.* 217.

Crithmum ſ. Foeniculum marinum minus. *Bauh. pin.* 288.

Crithmum marinum vulgare. *Park.* 1286.

Anglis, Sampire.

PENTANDRIA DIGYNIA.

Habitat in littoribus maritimis. 4. VIII.

Pharmac. Crithmi Herba.

HERACLEUM. 307.

Fruetus ellipticus, emarginatus, compressus, striatus,
Cor. difformis. *Involucrum* caducum.

Sphondylium. I. HERACLEUM foliolis pinnatifidis. *Hort. cliff.* 103.
Sp. pl. 249.

Sphondylium. *Ger. Em.* 1009. *R. Syn.* 205. vulgare
Park. 953.

Sphondylium vulgare hirsutum. *Baub. pin.* 157.

Sphondylium quibusdam vel Branca ursina Germanica,
Baub. hist. III. 2. 160.

β *Sphondylium* majus aliud laciniatis foliis. *Park.* 953.
Anglis, Cow Parsnep. *Cow-mumble* quibusdam
Habitat in sepibus, pratis et pascuis ubique. ♂. VII.

Hog-weed in Sussex-gather'd for Hogs.

LIGUSTICUM. 308.

Fruetus oblongus quinquefulcatus utrinque. *Corolla*
æquales.

scoticum. I. LIGUSTICUM foliis biternatis. *Sp. pl.* 250. *Fl.*
suec. 244.

Ligusticum scoticum apii folio. *Tourn.* 324. *R. Syn.*
 214.

Apium scoticum et *apium marinum* quibusdam. *R.*
Syn. 2d. ed. 109.

Anglis, Scottish Sea Parsley.

Habitat in clivis maritimis sed rarius. ♂. VII.

Cornubiense. 2. LIGUSTICUM foliis decompositis incis: radicali-
bus ternatis lanceolatis integerrimis. *Cent. pl.* 137.

Smyrnum tenuifolium nostras. *R. hist. III.* 254.
Syn. 209. t. 8.

Saxifraga Cornubiæ. *Pet. H. Br. t.* 26. f. 9.

Anglis, Cornwall Saxifrage.

Habitat in Cornubia. 4. VII.

ANGELICA. 309.

Fruetus subrotundus, angulatus, solidus, fulcis utrin-
 que 3.

I. AN.

PENTANDRIA DIGYNIA.

103

1. ANGELICA foliis æqualibus ovato-lanceolatis serratis. *Hort. cliff.* 97. *Fl. suec.* 246. *Sp. pl.* 251.
Angelica sylvestris. *Ger. Em.* 999. *Park.* 940. *R. Syn.* 208.
Angelica sylvestris major. *Bauh. pin.* 145.
Anglis, wild Angelica.
Habitat in sylvis, sepibus paludosis. 4. VI. VII.

S I U M. 510.

Fructus subovatus, striatus. *Involucrum* polyphyllum.
Petala cordata.

1. SIUM foliis pinnatis, umbella terminali. *Hort. cliff. latifolium.* 98. *Sp. pl.* 251.
Sium latifolium foliis variis. *Merr. pin.* *R. Syn.* 211.
Sium latifolium. *Bauh. pin.* 154.
Sium majus latifolium. *Ger. Em.* 256.
Sium Dioscoridis f. Pastinaca aquatica major. *Park.* 1240.
Anglis, great Water Parsnep.
Habitat in fluviis, paludibus; in pratis paludosis inter Rotherhithe et Deptford. *R. Syn.* 4. VII. VIII.
2. SIUM foliis pinnatis, umbellis axillaribus subsessili bus. *Sp. pl.* 251. *nodiflorum.*
Sium umbellatum repens. *Ger. Em.* 256. *R. Syn.* 211.
Sium aquaticum procumbens, ad alas floridum. *Hist. Ox. III.* 283. t. 5. f. 3.
Anglis, creeping Water Parsnep.
Habitat in fluviis et fossis ubique. 4. VII. VIII.
3. SIUM foliis pinnatis, umbellis axillaribus peduncula- tis, involucrio universali pinnatifido. *erectum.*
Sium five apium palustre foliis oblongis. *Bauh. pin.* 154. *R. Syn.* 211.
Sium majus angustifolium. *Ger. Em.* 256.
Sium minus alterum. *Park.* 1241.
Sium erectum foliis serratis, *D. Doody.* *R. Syn.* 211.
Anglis, upright Water Parsnep.
Habitat in fossis, rivulis et ad ripas fluviorum. 4. VII.

PENTANDRIA DIGYNIA.

S I S O N. 311.

Fruſtus ovatus, ſtriatuſ. *Involucrum* ſubtetraphyllum.

Amomum. 1. SISON foliis pinnatis, umbellis erectis. *Roy. lugdb.*

105. *Sp. pl.* 252.

Sison quod Amomum officinis noſtris. *Bauh. pin.* 154.

Sison vulgare, vel Amomum Germanicum. *Park.* 914.

Sium aromaticum Sison off. *Tourn.* 308 *R. Syn.* 211.

Petroſelinum Macedonicum Fuchſii. *Ger. Em.* 1016.

Anglis, baſtard Stone Parſley.

Habitat in ſylvis et ad ſepes ſubhumidis. 4. VIII. IX.

Pharmac. Amomi ſemina.

ſegetum. 2. SISON foliis pinnatis, umbellis cernuis. *Roy. lugdb.*

105. *Sp. pl.* 252.

Sium terreſtre, umbellis rarioribus. *Hiſt. Ox.* III.

283. t. 5. f. 6.

Sium arvenſe five ſegetum. *Tourn.* 308. *R. Syn.* 211.

Selinum Sii foliis. *Ger. Em.* 1018.

Selinum ſegetale. *Park.* 932.

Anglis, Corn Parſley, or Honewort.

Habitat in arvis et ad ſepes ſolo humido aut lutoſo.

8. VII.

inundatum. 3. SISON repens umbellis bifidis. *It. ſcan.* 397. *Sp. pl.*

252.

Sium foliis ſubmerſis capillaribus, emerſis pinnatis.

Roy. lugdb. 105.

Sium minimum, foliis imis ferulaceis. *Hiſt. Ox.* III.

223. t. 9. f. 5.

Sium puſillum foliis variis N. D. *Ph. Br. R. Syn.* 212.

Anglis, the leaſt Water Parſnep.

Habitat in ſoſſis, ſtagnis locis inundatis paſſim. In Eri-

ceto Putnejano et Hounſlejano verſus Hampton; in the ponds on Wandſworth common. *Blackſ. ſpec.*

Bot. inter Newington et Hornſey. 6. V. VI.

O E N A N T H E. 314.

Floſculi ſeffiles, difformes, in diſco ſteriles. *Fruſtus* calyce coronatus.

fiſtuloſa. 1 OENANTHE ſtolonifera, foliis caulinis pinnatis fi-
liſiformibus fiſtuloſis. *Sp. pl.* 254. Oenan-

Oenanthe aquatica. *Bauh. pin.* 162. *R. Syn.* 210.

Oenanthe palustris seu aquatica. *Park.* 895.

Filipendula aquatica. *Ger. Em.* 1060.

β Oenanthe aquatica triflora, caulibus fistulosis. *Hist.*

Ox. III. 289. *t.* 7. *f.* 8.

Anglis, Water Dropwort.

Habitat in fossis et paludibus frequens. 4. VII.

2. OENANTHE foliis radicalibus cuneatis fissis; *pimpinell-*
caulinis integris linearibus longissimis simplicioribus. *loides.*

Syst. Nat. 960. *Sp. pl.* 255.

Oenanthe Apii folio. *Bauh. pin.* 162.

Oenanthe staphylini folio aliquatenus accedens. *Bauh.*

hist. III. 191.

Oenanthe aquatica Pimpinellæ saxifragæ divisura no-
stras. *Pluk. ph. t.* 49. *f.* 4.

Anglis, Pimpernel Dropwort.

Habitat in fluentis pigrioribus et palustribus fossis in
parochia Quaplod agri Lincolnienfis non procul ab
oppido Spalding; *D. Pluk. R. Syn.* 4. VI.

3. OENANTHE foliis omnibus multifidis obtusis sub-crocata,
æqualibus. *Hort. cliff.* 99. *Sp. pl.* 254.

Oenanthe cicutæ facie Lobelii. *Park.* 894. *R. Syn.*
210.

Oenanthe Chærephylli foliis. *Bauh. pin.* 162.

Oenanthe succo viroso cicutæ facie Lobelio. *Bauh. hist.*
III. 193.

Filipendula cicutæ facie. *Ger. Em.* 1059.

Habitat ad ripas fluviorum. In salicetis circa Wind-
foriam, Bathoniam, ad ripas Thamefis alibique circa
Londinum nascitur. *R. Syn.* 4. VI.

B. Involucris partialibus; universali nullo.

B U N I U M. 298.

Corolla uniformis. Umbella conferta. Fructus ovati.

1. BUNIU M. *Hort. cliff.* 91. *Sp. pl.* 243.

Bulbocastanum. *Bauh. hist. III.* 30. *R. Syn.* 209.

Bulbocastanum majus folio Apii. *Bauh. pin.* 162.

Bulbocastanum majus et minus. *Ger. Em.* 1065.

Nucula

Bulbocast-
anum.

Nucula terrestris major. Park. 893.

Anglis, Earth-nut, Kipper-nut, Pig-nut, or Hawk-nut.

Habitat in pascuis et pratis. 4. V. VI.

SESELI. 322.

Umbellæ globosæ. Involucrum polyphyllum. Fructus ovatus, striatus.

Caruifolia. 1. SESELI foliis bipinnatis sublinearibus: petiolis basi membranaceis, seminibus ovalibus. *Sp. pl. 260.*

Seseli pratense nostras. Park. 905. R. Syn. 216.

Saxifraga anglica facie seseli pratensis. Ger. Em. 1047.

Saxifraga anglorum, foliis Foeniculi latoribus, radice nigra, flore candido, semine Foeniculi, similis Silao.

Bauh. hist. III. 171.

Caruifolia. Bauh. hist. III. 171. pin. 158. Vail. parif. t. 5. f. 2.

Anglis, Meadow Saxifrage.

Habitat in pratis et pascuis humidiusculis. 4. VIII.

CICUTA. 316.

Fructus subovatus, sulcatus.

virens. 1. CICUTA umbellis folio oppositis, petiolis marginatis obtusis. *Sp. pl. 255.*

Cicuta. Hort. cliff. 100. Mat. med. 129. Fl. suec. 253. Roy. lugdb. 109.

Cicuta aquatica. Gesn. Hort. 253. Coll. 68. Fl. lapp. 103.

Sium alterum. Dod. 589. olusatrici facie. Lob. ic. 208, Ger. Em. 256. R. Syn. 212.

Sium majus alterum angustifolium. Park. 1241.

Sium Erucæ folio. Bauh. pin. 154.

Anglis, long-leaved Water Hemlock.

Habitat in aquis. In a shallow pool of water on Hounslow heath, by the road-side near the town; and in some pools of water at Thistleworth. R. Syn. 4. VII.

PENTANDRIA DIGYNIA.

107

PHELLANDRIUM. 315.

Fruetus ovatus lævis, coronatus perianthio et pistillo.

1. PHELLANDRIUM foliorum ramificationibus diva- *aquati-*
ricatis. *Sp. pl.* 255. *cum.*

Phellandrium vel Cicutaria aquatica quorundam.

Bauh. hist. III. 183. *R. Syn.* 215.

Cicutaria palustris. *Lob. ic.* 735. *Ger. Em.* 1063.

tenuifolia. *Bauh. pin.* 161.

Anglis, Water Hemlock.

Habitat in fluviiis et fossis. ♂.

ÆTHUSA. 317.

Involucrum dimidiatum, triphyllum, pendulum. *Fructus* striatus.

1. ÆTHUSA. *Hort. cliff.* 100. *Sp. pl.* 256. *Cynapium.*

Cicutaria tenuifolia. *Ger. Em.* 1063. *R. Syn.* 215.

Cicuta minor seu fatua. *Park.* 933.

Cicuta minor Petroselino similis. *Bauh. pin.* 160.

Anglis, Fools Parsley.

Habitat in arvis et inter olera. ☉. VIII. IX.

CORIANDRUM. 318.

Cor. radiata. *Involucrum* universale monophyllum.

Partialia dimidiata. *Fruetus* sphaericus.

1. CORIANDRUM fructibus globosis. *Hort. cliff.* 100. *fativum.*
Sp. pl. 256.

Coriandrum. *Bauh. hist. III.* 98. *Ger. Em.* 1012.

R. Syn. 221.

Coriandrum majus. *Bauh. pin.* 158. vulgare. *Park.* 918.

Anglis, Coriander.

Habitat in fimetis et ad vias. ☉. VI.

Pharmac. Coriandri semina.

SCANDIX. 319.

Cor. difformis. *Fruetus* subulatus. *Petala* integra.

Flosculi disci sæpe masculi.

1. SCANDIX feminibus lævibus rostro longissimo. *Hort. Pecten.* ♀
cliff. 101. *Sp. pl.* 256.

Scandix

PENTANDRIA DIGYNIA.

Scandix femine rostrato vulgaris. *Baub. pin.* 152. *R. Syn.* 207.

Scandix vulgaris seu Pecten Veneris. *Park.* 916.

Pecten Veneris seu Scandix. *Ger. Em.* 1040.

Anglis, Shepherd's-needle, or Venus-comb.

Habitat in arvis. ☉. VI. VII.

Anthrif. 2. SCANDIX feminibus ovatis hispidis, corollis uniformibus, caule lævi. *Sp. pl.* 257.

Myrrhis sylvestris feminibus asperis. *Baub. pin.* 160.

R. Syn. 220.

Myrrhis sylvestris Neapolitana atque etiam Anglicana. *Park.* 935.

Cerefolium five Myrrhis nova Equicolorum Columnæ. *Ger. Em.* 1038.

Anglis, small Hemlock Chervil with rough Seeds.

Habitat in aggeribus ruderatis ubique. ☉. V. VI.

CHÆROPHYLLUM. 320.

Involucrum reflexum, concavum. *Petala* cordata.

Fruetus oblongus, lævis.

* *Floribus omnibus hermaphroditis.*

sylvestre. 1. CHÆROPHYLLUM flosculis omnibus fertilibus, caule æquali. *Sp. pl.* 258.

Chærophyllum feminibus lævibus nitidis, petiolis ramis æqualibus. *Hort. cliff.* 101. *Roy. lugdb.* 112.

Cicutaria vulgaris. *Baub. hist.* III. 181. *R. Syn.* 207.

Cicutaria alba Lugdunensis. *Ger. Em.* 1038.

Myrrhis sylvestris. *Park.* 935. feminibus lævibus. *Baub. pin.* 160.

Anglis, wild Cicely, or Cow-weed.

Habitat ad sepes. ☉. V. VI.

** *Floribus disci masculis.*

temulum. 2. CHÆROPHYLLUM caule scabro: geniculis tumidis, umbellis cernuis. *Sp. pl.* 258. *Fl. suec.* 258.

Chærophyllum caule maculato: geniculis tumidis. *Hort. cliff.* 102.

Cherophyllum sylvestre. *Baub. pin.* 152.

Cherophyllum

CherEOFOLIUM sylvestre. Ger. Em. 1038. Park. 915.

R. Syn. 207.

Anglis, wild Chervil.

Habitat ad sepes ubique. ☉. VII. VIII.

C. *Involucro nullo* ; nec universali, nec Partialibus.

P A S T I N A C A. 324.

Fruētus ellipticus, compresso-planus.

1. PASTINACA foliis simpliciter pinnatis. Hort. cliff. sativa.

105. Sp. pl. 263.

Pastinaca sylvestris latifolia. Bauh. pin. 155. R. Syn. 206.

Pastinaca latifolia sylvestris. Ger. Em. 1025. Park. 944.

Pastinaca Germanica sylvestris quibusdam Elaphoboscum. Bauh. hist. III. 149.

β Pastinaca latifolia sativa. Ger. Em. 1025. Park. 944.

Bauh. pin. 155.

Anglis, wild Parsnep. β Garden Parsnep.

Habitat in marginibus agrorum. ♂. VII.

S M Y R N I U M. 325.

Fruētus oblongus, striatus. Petala acuminata carinata.

1. SMYRNIUM foliis caulinis ternatis petiolatis serratis. Olusat.

tis. Hort. cliff. 105. Roy. lugdb. 114. Sauv. monsp. trum.

262. Sp. pl. 262.

Smyrnum. Matth. 773. R. Syn. 208.

Hippofelinum. Ger. Em. 1019. seu Smyrnum vulgare. Park. 930.

Hippofelinum Theophrasti f. Smyrnum Dioscoridis.

Bauh. pin. 154.

Anglis, Alexanders.

Habitat in fossis et rupibus maritimis. About Scarborough castle ; near Deptford, Battersea, and Vauxhall ; about Nottingham plentifully, Mr. J. Sherard.

R. Syn. ♂. V. VI.

Pharmac. Smyrni Radix, Herba, Semina.

ANETHUM. 326.

Fructus subovatus, compressus, striatus.

- Foeniculum.* 1. ANETHUM fructibus ovatis. *Hort. cliff.* 106.
Sp. pl. 263.
Foeniculum vulgare. *Park.* 884. *Ger. Em.* 1032.
R. Syn. 217.
Foeniculum vulgare Germanicum. *Bauh. pin.* 147.
 β *Foeniculum dulce.* *Bauh. pin.* 147.
Anglis, Finckle, β Fennel.
Habitat in clivis cretaceis passim. ♂. VIII.
Pharmac. *Foeniculi semina.*

CARUM. 327.

Fructus ovato-oblongus, striatus. *Involucrum* monophyllum. *Petala* carinata, emarginata.

- Carvi.* *1. CARUM. *Dod. pemp.* 299. *Fl. lapp.* 105. *suec.* 260.
Mat. med. 138. *Hall. helv.* 428. *Sp. pl.* 263. vulgare. *Park.* 910.
Carum seu Careum. *Ger. Em.* 1034. *R. Syn.* 213.
Cuminum pratense Carui officinarum. *Bauh. pin.* 158.
Anglis, Caraways.
Habitat in pratis et pascuis passim. ♂. V. VI.
Pharmac. *Carui semina.*

PIMPINELLA. 328.

Fructus ovato-oblongus. *Corolla* subradiata.

- major.* 1. PIMPINELLA foliis pinnatis: foliolis cordatis, impari trilobo.
Pimpinella saxifraga. *Ger. Em.* 1044. *R. Syn.* 213.
Pimpinella saxifraga hircina major. *Park.* 947.
Pimpinella saxifraga major umbella candida. *Bauh. pin.* 159.
 β *Pimpinella saxifraga major, degener, seu foliis dissectis.* *Hist. Ox.* III. 289.
Anglis, great Burnet Saxifrage.
Habitat in sylvis et sepibus. *In sylvis agro Cantabrigiensi, Bedfordiensi, Leicesteriensi, Cantiano, et alibi.* *R. Syn.* 4. VIII.

2. PIM-

2. PIMPINELLA foliis pinnatis: foliolis radicalibus *saxifraga*, subrotundis; summis linearibus. *Sp. pl.* 263.

Fl. suec. 261.

Pimpinella saxifraga minor, foliis Sanguisorbæ. *R. Syn.* 213.

Pimpinella major altera. *Bauh. pin.* 159.

Pimpinella saxifraga major nostras. *Park.* 946.

Pimpinella seu saxifraga minor. *Ger. Em.* 1044.

- β Pimpinella saxifraga minor. *Bauh. pin.* 160.

Pimpinella saxifraga hircina minor. *Park.* 947.

Anglis, small Burnet Saxifrage.

Habitat in pascuis siccioribus et glareosis. 4. VIII.

A P I U M. 329.

Fruetus ovatus-striatus. *Involucrum* monophyllum.

Petala inflexa.

1. APIUM foliis caulinis cuneiformibus. *Hort. cliff. graveolens.* 107. *Sp. pl.* 264.

Apium palustre et officinarum. *Bauh. pin.* 154. *R. Syn.* 214.

Apium vulgare seu palustre. *Park.* 926.

Eleoselinum seu Paludapium. *Ger. Em.* 1014.

Anglis, Smallage.

Habitat in palustribus et fossis. 3. VIII.

Æ G O P O D I U M. 330.

Fruetus ovato-oblongus, striatus.

1. ÆGOPODIUM foliis caulinis summis ternatis. *Sp. Podagraria.* pl. 265.

Podagraria. *Riv. pent.* 47. vulgaris. *Park.* 943.

Angelica sylvestris minor f. erratica. *Bauh. pin.* 155. *R. Syn.* 208.

Herba Gerardi. *Ger. Em.* 1001.

Anglis, Herb-gerard, Gout-weed, or Ash-weed.

Habitat ad sepes et in oleraceis, hortis, frequens. 4. V.

TRI-

T R I G Y N I A.

* Flores supèri.

V I B U R N U M. 332.

Cal. quinquepartitus, superus. *Cor.* quinquefida.
Bacca monosperma.

Lantana. 1. VIBURNUM foliis cordatis ferratis venosis subtus tomentosis. *Vir. cliff.* 25. *Hort. ups.* 86. *Sp. pl.* 268. *Viburnum.* *Park.* 1448. *Baub. pin.* 429. *R. Syn.* 460.

Lantana five *Viburnum.* *Ger. Em.* 1490.

Anglis, the pliant Mealy-tree. *Park.* The way-faring-tree. *Ger.*

Habitat in sepibus. h. V.

Opulus. 2. VIBURNUM foliis ovatis, petiolatis glandulosis. *Sp. pl.* 268.

Opulus Ruellii 281. *Tourn.* 607. *R. Syn.* 460.

Sambucus aquatica, flore simplici. *Baub. pin.* 456.

Sambucus palustris seu *aquatica.* *Park.* 208.

Sambucus aquatilis seu *palustris.* *Ger. Em.* 1424.

Anglis, Water Elder.

Habitat in sylvis et sepibus humidis. h. V. VI.

S A M B U C U S. 334.

Cal. quinquepartitus. *Cor.* quinquefida. *Bacca* trisperma.

nigra. 1. SAMBUCUS cymis quinquepartitis, caule arboreo. *Sp. pl.* 269.

Sambucus caule arboreo ramoso, floribus umbellatis. *Roy. lugdb.* 243.

Sambucus. *Ger. Em.* 1422. *vulgaris.* *Park.* 207. *R. Syn.* 461.

Sambucus fructu in umbella nigra. *Baub. pin.* 456.

β *Sambucus fructu* in umbella viridi. *Baub. pin.* 456.

γ *Sambucus laciniato folio.* *Baub. pin.* 456.

Anglis, common Elder.

Habitat in sylvis et sepibus humidis. h. IV.

Pharmac, Sambuci Cortex, Folia, Flores, Baccæ.

2. S A M-

PENTANDRIA TRIGYNIA.

113

2. *SAMBUCUS* cymis tripartitis, stipulis foliaceis. *Sp. Ebulus*.
pl. 269.
Sambucus humilis seu *Ebulus*. *Baub. pin.* 456. *R.*
Syn. 461.
Ebulus five *Sambucus humilis*. *Ger. Em.* 1426.
Park. 209.
Anglis, dwarf Elder, Walwort, or Danewort.
Habitat in sepibus et ad vias. *h.* VII.
Pharmac. *Ebuli Radix, Folia, Flores, Baccæ.*

** *Flores inferi.*

STAPHYLEA. 336.

- Cal.* quinquepartitus. *Petala* quinque. *Capf.* inflatæ,
 connatæ. *Sem.* duo globosa cum cicatrice.
 *1. *STAPHYLEA* foliis pinnatis. *Hort. cliff.* 112. *pinnatas*
Roy. lugdb. 436. *Sp. pl.* 270.
Staphylodendron. *Baub. hist. I.* 274. *R. Syn.* 468.
Nux vesicaria. *Park.* 1417. *Ger. Em.* 1437.
Pistacea sylvestris. *Baub. pin.* 401.
Anglis, Bladder Nut-tree.
Habitat in sepibus circa Pontemfractum comitatus Ebo-
racensis oppidum ; in Cantio circa Ashfordiam inve-
niri apud Parkinsonum legimus. *R. Syn.* *h.* VI.

ALSINE. 342.

- Cal.* pentaphyllus. *Petala* quinque, bipartita, æqua-
 lia. *Capf.* unilocularis, trivalvis.
 1. *ALSINE* petalis bipartitis, foliis ovato-cordatis. *Fl. media*.
lapp. 186. *suec.* 267. *Sp. pl.* 272.
Alfne media. *Baub. pin.* 250. seu minor. *Ger. Em.*
 611.
Alfne vulgaris seu morsus gallinæ. *Baub. hist. III.*
 363. *R. Syn.* 347.
Anglis, common Chickweed.
Habitat ubique in umbrosis, cultis. ☉. IV.—X.

I

TETRA-

T E T R A G Y N I A.

P A R N A S S I A. 345.

Cal. quinquepartitus. *Petala* quinque. *Nectaria* quinque cordata ciliata apicibus globosis. *Capf.* quadrivalvis.

palustris. 1. PARNASSIA. *Fl. lapp.* 108. *suéc.* 268. *Sp. pl.* 273. *Hall. helv.* 316.

Parnassia vulgaris et *palustris.* *Tourn.* 246. *R. Syn.* 355.

Gramen Parnassi minus. *Ger. Em.* 840.

Gramen Parnassi vulgare. *Park.* 429.

Gramen Parnassi flore albo simplici. *Bauh. pin.* 309.

Anglis, Grass of Parnassus.

Habitat in pratis humidis. 4. VIII.

P E N T A G Y N I A.

* *Flores inferi.*

S T A T I C E. 348.

Cal. monophyllus, integer, plicatus. *Petala* quinque. *Sem.* unicum.

Armeria. 1. STATICE caule nudo simplici capitato. *Sp. pl.* 274. *Fl. suéc.* 269.

Statice montana minor. *Tourn.* 341. *R. Syn.* 203.

Limonium majus flore globoso. *Hist. Ox.* III. 601. *f. 15. t. 1. f. 29.*

Caryophyllus montanus minor. *Bauh. pin.* 211.

Caryophyllus marinus minimus Lobelii. *Ger. Em.* 602.

Gramen marinum minus. *Park.* 1276.

Anglis, Thrift, Sea Gilly-flower.

Habitat in pratis maritimis. 4. VII. VIII.

Limoni- 2. STATICE caule nudo paniculato tereti foliis lævi-
um, bus. *Sp. pl.* 274. *Fl. suéc.* 270.

Limonium. *Ger. Em.* 411. *R. Syn.* 200. *majus vulgatus.* *Park.* 1234.

Limonium maritimum majus. *Bauh. pin.* 192.

β *Limo-*

β Limonium minus. *R. Syn.* 202

γ Limonium anglicum minus caulibus ramosioribus, floribus in spicis rarius fitis. *R. hist. III.* 247.

Anglis, Sea Lavender.

Habitat in littoribus maritimis. 4. VII. VIII.

Pharmac. Limonii Radix.

3. STATICE caule subnudo paniculato prostrato, ramis *reticulata*, sterilibus retroflexis nudis, foliis cuneiformibus. *Sp. pl.* 275.

Statice foliis obverfo-ovatis, ramis nudis humifusis, intricatis. *Sauv. monsp.* 15.

Limonium minus, virgulis retiformiter se invicem implicatis. *Pluk. ph. t.* 42. *f.* 4.

Limonium reticulatum. *R. hist.* 396. *Bocc. sic.* 82. *t.* 44.

Anglis, matted Sea Lavender.

Habitat in littoribus maritimis.

L I N U M. 349.

Cal. pentaphyllus. *Petala* quinque. *Caps.* quinque-valvis, decemlocularis. *Sem.* solitaria.

* *Foliis alternis.*

1. LINUM calycibus capsulisque mucronatis, petalis *usitatissimis* crenatis, foliis lanceolatis alternis, caule subsolitum. *Sp. pl.* 277.

Linum ramis foliisque alternis, lineari-lanceolatis, radice annua. *Hort. cliff.* 114. *Roy. lugdb.* 433.

Linum arvense. *Baub. pin.* 214. *R. hist.* 1073.

Linum sylvestre vulgatius. *Park.* 1334.

β Linum sativum. *Baub. pin.* 214. *Ger. Em.* 556. *R. Syn.* 362.

Anglis, Flax.

Habitat in arvis *passim*. ☉. VI.

Pharmac. Lini semina.

2. LINUM calycibus capsulisque obtusis, foliis alternis *perennans*, lanceolatis integerrimis. *Sp. pl.* 277. *Mill. fig. t.* 166. *f.* 2.

Linum foliis alternis lanceolatis integerrimis, calycibus apice obtusis, capsulis muticis. *Hort. upf.* 72.

PENTANDRIA PENTAGYNIA.

Linum sylvestre cæruleum perenne erectius, flore e capitulo majore. *R. Syn.* 362.

Linum perenne majus cæruleum, capitulo majore. *Hist. Ox.* II. 573.

β *Linum sylvestre cæruleum* perenne procumbens, flore et capitulo minore. *R. Syn.* 362.

Anglis, perennial blue Flax.

Habitat in pascuis et pratis sterilioribus. In agri Cantabrigienſis collibus Gogmagog dictis, ad margines ſartorum copioſe; alſo at Croſby Ravensworth, and between Shap and Threapland in Weſtmoreland, Mr. Lawſon; and on the balks of corn-fields about Bernack-heath in Northamptonſhire. *R. Syn.* in iſula Shepey copioſe. 4. VI. *Uſ'd in Siberia*. found in ſome fields near Whorwell woods in *Hants*.

tenuifolium.

3. *LINUM* calycibus acuminatis, foliis ſparſis linearifetaceis retrorſum ſcabris. *Sp. pl.* 278.

Linum perenne, foliis anguſtioribus, calyce ariſtato. *Hall. herb.* 371.

Linum sylvestre anguſtifolium, floribus dilute purpureſcentibus vel carneis. *Bauh. pin.* 214. *R. Syn.* 362.

Linum sylvestre VI. anguſtifolium. *Cluſ. hiſt.* I. 318. *Anglis*, narrow-leaved wild Flax.

Habitat in pratis maritimis aridis. In the paſtures by the ſea-ſide about St. Ives and Truro in Cornwall, plentifully. Found by Mr. J. Sherard plentifully in a field on the left hand of the lane that leads from the ſea to Pett, a ſmall village, four miles from Haſtings, and on Beacon hill in Kent. *R. Syn.* 4. VI. VII.

** *Foliis oppoſitis.*

× *catharticum.*

4. *LINUM* foliis oppoſitis ovato-lanceolatis, caule diſchotomo, corollis acutis. *Sp. pl.* 281. *Mat. med.* 157. *Fl. ſuec.* 271.

Linum sylvestre catharticum. *Ger. Em.* 560. *R. Syn.* 362.

Linum pratense floſculis exiguis. *Bauh. pin.* 214.

Chamælinum Cluſii flore albo ſ. *Linum sylvestre catharticum.* *Park.* 1336.

Anglis, purging Flax.

Habitat

Habitat in pratis et pascuis siccioribus. ☉. V. VI.

Pharm. Lini cathartici Herba.

5. LINUM foliis oppositis, caule dichotomo, floribus *Radiola*.
 tetrandris tetragynis. *Sp. pl.* 281. *Fl. suec.* 272.
Chamælinum vulgare. *Vaill. parif.* 33. t. 4. f. 6.
Radiola vulgaris serpyllifolia. *R. Syn.* 345. t. 15. f. 3.
Millegrana minima. *Ger. Em.* 569. f. *Herniaria* mi-
 nor. *Park.* 447.
Polygonum minimum f. *millegrana minima.* *Baub.*
pin. 282.
Anglis, the least Rupture-wort, or All-feed.
 Habitat in *Ericetis glareosis subhumidis.* On Wandf-
 worth common. *Blackf. spec. bot. in ericeto* Putne-
 jano copiose. ☉. VIII.

DROSER A. 352.

Cal. quinquefidus. *Petala* quinque. *Caps.* unilocu-
 lar: apice quinquevalvi. *Sem.* plurima.

1. DROSER A scapis radicatis, foliis orbiculatis. *Fl. rotundi-*
lapp. 109. *Zeyl.* 120. *suec.* 273. *Sp. pl.* 281. *folia.*
Ros folis folio rotundo. *Baub. pin.* 357. *R. Syn.* 357.
Ros folis. *Baub. hist. III.* 761. major. *Ger. Em.*
 1557. *Park.* 1052.
 2. Rorella rotundifolia perennis. *R. Syn.* 356.
Anglis, round-leaved Sun-dew, or Rosa-folis.
 Habitat in paludibus. 4. VII. VIII.
Pharmac. Ros folis Herba.
 2. DROSER A scapis radicatis, foliis oblongis. *Fl. lapp. longifolia.* x
 110. *suec.* 358.
Ros folis folio oblongo *Baub. pin.* 357. *R. Syn.* 356.
Ros folis sylvestris longifolius. *Park.* 1052.
Ros folis minor. *Ger. Em.* 1557.
 3. Rorella longifolia perennis. *R. Syn.*
Anglis, long-leaved Rosa-folis, or Sun-dew.
 Habitat in paludosis. Upon Hinton-moor near Cam-
 bridge: on Bagshot heath, Mr. Doody. *R. Syn.* at
 Battle's-well, near Harefield, plentifully. *Blackf. sp.*
bot. 4. VI, VII. Bagshot-Heath near y^e XXXI mile
 = stone. I 3 S I B-

PENTANDRIA POLYGYNIA.

SIBBALDIA. 354.

Cal. pentaphyllus decemfidus. *Petala* quinque, calyci inserta. *Styli* e latere germinis. *Sem.* quinque.

procumbens. 1. SIBBALDIA foliolis tridentatis. *Fl. lapp.* 111. *suec.* 216. *Sp. pl.* 284.

Pentaphylloides pumila foliis ternis ad extremitates trifidis. *R. Syn.* 256.

Pentaphylloides fruticosum minimum procumbens, flore luteo, foliis sericeis fragariæ ternis. *Pluk. ph.* t. 212. f. 3.

Fragariæ sylvestri affinis planta, flore luteo. *Sibb. pr. hist. nat. scot. p. II.* 25. t. 6. f. 1.

Anglis, bastard Cinquefoil.

Habitat in sylvis regionis Jernensis Scotiæ sponte provenit. *R. Syn.* 4. VII. VIII.

POLYGYNIA.

MYOSURUS. 355.

Cal. pentaphyllus, basi adnatus. *Nectaria* quinque subulata, petaliformia. *Sem.* numerosa.

minimus. 1. MYOSURUS foliis integris. *Hort. cliff.* 117.

Myosurus. *Bauh. hist. III.* 512. *R. Syn.* 251.

Holosteum affinis, Cauda muris. *Bauh. pin.* 190.

Holosteum Loniceri, Cauda muris vocatum. *Park.* 500.

Cauda muris. *Ger. Em.* 426.

Anglis, Mouse-tail.

Habitat in arvis pratis et pascuis solo glareoso. In a lane that goes from Copenhagen-house to Kentish-town. Dr. Wilmer. *Inter segetes circa Wimbleton, prope Londinum.* ☉. IV. V.

Classis

Classis VI.

HEXANDRIA
MONOGYNIA.

* *Flores Caliculati calyce corollaque instructi.*

B E R B E R I S. 399.

Cal. hexaphyllus. *Petala* sex; ad ungues glandulis 2.
Stylus o. *Bacca* disperma.

I. BERBERIS pedunculis racemosis. *Mat. med.* 290. *vulgaris.*
Fl. suec. 312. *Sp. pl.* 330.

Berberis spinis triplicibus. *Hort. cliff.* 122.

Berberis. *Park.* 1559. *dumetorum.* *Bauh. pin.* 454.
R. Syn. 465.

Spina acida sive *Oxyacantha.* *Ger. Em.* 1325.

Anglis, Barberry or Pipperidge-bush.

Habitat in sylvis et sepibus passim. h.

Pharmac. *Berberidis, Cortex, Baccæ.*

F R A N K E N I A. 401.

Cal. quinquefidus, infundibuliformis. *Petala* quinque.
Stigma sexpartitum. *Capsf.* unilocularis, trivalvis.

I. FRANKENIA foliis linearibus basi ciliatis. *Sp. pl. lævis.*
331.

Frankenia foliis aciformibus congestis. *Sauv. monsp.* 46.

Franca floribus solitariis sessilibus, foliis prismaticis triangularibus. *Guett. stamp. II* 459.

Franca maritima supina saxatilis glauca ericoides sem-pervirens, flore purpureo. *Mich. gen.* 23 t. 22 f. 1.

Lychnis supina maritima Ericæ facie. *R. Syn.* 338.

Erica supina maritima. *Ph br. anglica.* *Park.* 1484.

Polygonum maritimum minus, foliolis serpylli. *Bauh. pin.* 281.

HEXANDRIA MONOGYNIA.

Polygonum pusillo vermiculato serpylli folio. Ger. Em. 567.

Anglis, smooth Sea Heath.

Habitat in *palustribus maritimis*. In Lovingland, just over the water at Yarmouth; in the marshes about Thurrington in Essex; on the sandy ditch-banks by the way-side going from Malden to Goldhanger; in Selsey island plentifully. R. Syn. In *insula* Shepey copiose. 4. VIII.

pulverulenta.

2. FRANKENIA foliis obovatis retusis subtus pulveratis. Sp. pl. 332.

Frankenia foliis ovalibus. Sauv. monsp. 168.

Alfine maritima supina foliis Chamæscis. Tourn. 665. R. Syn. 352.

Anthyllis maritima Chamæscis similis. Baub. pin. 282. *Anglis*, broad-leaved Sea Heath.

Habitat in *littoribus maritimis*. Found on the coast of Suffex, and sent by Mr. Brewer. R. Syn. 3. VII.

* Flores *spathacei*.

A L L I U M. 370.

Cal. sexpartita, patens. *Spatha* multiflora. *Umbella* congesta. *Capf.* supera.

* *Foliis caulinis planis*.

Ampeloprasum.

1. ALLIUM caule planifolio umbellifero, umbella globosa, staminibus tricuspidatis, petalis carina scabris. Syst. nat. 977.

Allium caule planifolio umbellifero, umbella globosa, staminibus tricuspidatis, radice laterali. Sp. pl. 294.

Allium staminibus alterne trifidis, foliis gramineis, floribus sphaerice congestis, radice laterali solida. Hall. all. 5.

Allium holmense sphaerico capite. R. Syn. 370.

Allium sphaerico capite, folio latiore f. *Scorodo prasum* alterum. Baub. pin. 74. Rudb. elyf. 2. 151.

Scorodo prasum primum Clusii. Ger. Em. 180.

Anglis, great round-headed Garlick.

Habitat

Habitat in parva quadam insula Holms dicta in Sabri-
næ æstuario copiose proveniente observavit. D. New-
ton. R. Syn. 24.

2. ALLIUM caule planifolio bulbifero, vaginis tereti- *arenari-*
bus, spatha mutica, staminibus tricuspidatis. *It. um.*
scan. 227. Sp. pl. 297.

Allium sylvestre amphicarpon, foliis porraceis, flori-
bus et nucleis purpureis. R. Syn. 370.

Anglis, broad-leaved mountain Garlick.

Habitat in montibus Westmorlandicis observavit, D.
Lawson. In Troutbeck-holm by Great Strickland.
R. Syn. 24. VII.

** *Foliis caulinis teretibus.*

3. ALLIUM caule teretifolio bulbifero, staminibus tri- *vineale.*
cuspidatis. *Sp. pl. 299.*

Allium staminibus alterne trifidis, foliis fistulosis, capite
bulbifero, radice multiplici. *Hall. all. 4.*

Allium sylvestre. *Park. 870. Ger. Em. 179. R. Syn.*
369.

Allium campestre juncifolium capitatum purpurascens
majus. *Bauh. pin. 74.*

Allium sylvestre vinearum. *Bauh. pin. 72.*

Anglis, Crow-garlick.

Habitat in pratis et pascuis. 24. VI,

4. ALLIUM caule teretifolio bulbifero, foliis scabris se- *oleraceum.*
miteretibus subtus sulcatis. *Sp. pl. 299.*

Allium umbella bulbifera, vagina bicorni, foliis tere-
tibus. *Hall. all. 23. f. 4.*

Allium sylvestre bicornne flore exherbaceo albicante,
cum triplici in singulis petalis stria atro-purpurea.
R. Syn. 370.

Allium montanum bicornne, flore exalbido. *Bauh.*
pin. 75.

- β Allium montanum bicornne purpureum proliferum. *R.*
Syn. 396.

Anglis, wild Garlick with an herbaceous striated
flower.

Habitat

HEXANDRIA MONOGYNIA.

Habitat in pratis montosis, et inter segetes. In a Cornfield in Black-Notley in Essex, belonging to the hall called Westfield, adjoining to Leeze-lane, plentifully. R. Syn. in pratis prope Bristolium copiose. α et β in arvis in Westmorlandia frequens. 4. VII.

*** Foliis radicalibus, scapo nudo.

- ursum. 5. ALLIUM scapo nudo semicylindrico, foliis lanceolatis petiolatis umbella fastigiata. Sp. pl. 300.
Allium sylvestre latifolium. Bauh. pin. 74. R. Syn. 370.
Allium ursum. Ger. Em. 180. Park. 871.
Anglis, Ramfon.
Habitat in sylvis et sepibus frequens. 4. V.

BULBOCODIUM. 368.

Cor. infundibuliformis, hexapetala : unguibus angustis. Caps. supera.

- serotinum. 1. BULBOCODIUM foliis subulato-linearibus. Roy. lugdb. 41. Sp. pl. 294.
Bulbocodium alpinum juncifolium, flore unico : intus albo extus squalide rubente. R. Syn. 374. t. 17. f. 1.
Pseudo-Narcissus gramineo folio. Bauh. pin. 51. pr. 27.
Bulbosa alpina juncifolia, pericarpio unico erecto in summo cauliculo dodrantali. R. Syn. 374.
Anglis, mountain Saffron.
Habitat in excelsis rupibus montis Snowden ; v. g. Trigvylchou y Clogwyn du ymhen y Gluder, Clogwyn yr Ardhu Crib y Disth, &c. D. Lhwyd. R. Syn. 4. VIII.

NARCISSUS. 364.

Petala sex : Nectario infundibuliformi, monophyllo, integro. Stam. intra nectarium.

- poeticus. *1. NARCISSUS spatha subuniflora, nectarii limbo rotato brevissimo. Hort. ups. 74. Sp. pl. 289.
Narcissus medio luteus. Ger. Em. 124. vulgaris. Park. 74. R. Syn. 371.

Narcissus

Narcissus pallidus circulo luteo. *Bauh. pin.* 51.
Anglis, common pale Daffodil, or Primrose peerless.
Habitat in pratis et pascuis sed rarius. 4. IV.

2. NARCISSUS spatha uniflora, nectarii limbo campanulato erecto petalis æquali. *Sp. pl.* 289. *Pseudo-Narcissus.*
 Narcissus sylvestris pallidus calyce luteo. *Bauh. pin.* 52. *R. Syn.* 371.

Pseudo-Narcissus Anglicus. *Ger. Em.* 138. vulgaris.
Park. 74.

Anglis, wild English Daffodil.

Habitat in sylvis et ad sepes. In sylvis prope Erith in agro Cantiano copiose. 4. III. IV.

*** *Flore nudi.*

HYACINTHUS. 385.

Cor. campanulata: pori tres melliferi germinis.

1. HYACINTHUS corollis campanulatis sexpartitis a-non scripte revolutis. *Hort. cliff.* 125. *Roy. lugdb.* 27. *tus.*
Sauv. monsp. 17. *Sp. pl.* 316.

Hyacinthus oblongo flore cæruleus major. *Bauh. pin.* 43.

Hyacinthus anglicus. *Ger. Em.* 110. *R. Syn.* 373.

Hyacinthus Anglicus, Belgicus f. Hispanicus. *Park.* par. 122.

Anglis, English Hyacinth, or Harebells.

Habitat in sylvis et sepibus. 4. V.

SCILLA. 378.

Cor. hexapetala, patens, decidua. *Filamenta* filiformia.

1. SCILLA radice solida, floribus erectiusculis paucio-bifoliaribus. *Hort. cliff.* 123. *Sp. pl.* 309.

Hyacinthus stellaris bifolius Germanicus. *Bauh. pin.* 45.

Hyacinthus stellatus Fuchsii. *Ger. Em.* 106. *R. Syn.* 372.

Hyacinthus stellatus vulgaris f. bifolius Fuchsii. *Park.* par. 126.

Anglis, vernal star Hyacinth.

Habitat

Habitat in insulis quibusdam circa Cambriam copiose, ut v. g. Bradsea, necnon in clivis maritimis ipsius Cambriæ. R. Syn. 4. IV. V.

autumna- 2. SCILLA radice solida, foliis filiformibus linearibus, floribus corymbosis, pedunculis nudis adscendentibus longitudine floris. *Sp. pl.* 309.

Scilla radice solida, foliis setaceis, floribus fastigiatis, pedunculis arcuatis ex ala tuberculi mamillaris. *Guett. stamp.* 131.

Hyacinthus Autumnalis minor. *Ger. Em.* 110. *Park. par.* 132. *R. Syn* 373.

Hyacinthus stellaris autumnalis minor. *Bauh. pin.* 47. *Anglis*, the lesser autumnal star Hyacinth.

Habitat in pascuis siccis. In rupe S. Vincentii prope Bristolium inque promontorio, The Lizard Point dicto, copiose; eundem in Ericeto nigro in comitatu Cantii invenit, D. Plukenet. On Mouldsworth, over-against Hampton-court; near Ditton, and on Kingston-bridge, and going to Hounslow-heath. *Merr. pin. R. Syn.* 4. IX.

ORNITHOGALUM. 377.

Cor. hexapetala, erecta, persistens, supra medium patens. *Filamenta* alterna basi dilatata.

* *Seminibus omnibus subulatis.*

luteum. 1. ORNITHOGALUM scapo anguloso diphylo, pedunculis umbellatis simplicibus. *Fl. suec.* 285. *Sp. pl.* 306.

Ornithogalum luteum. *Bauh. pin.* 71. *Park. par.* 140. *R. Syn.* 372.

Ornithogalum luteum seu Cepe agraria. *Ger. Em.* 165. *Anglis*, yellow star of Bethlehem.

Habitat in comitatus Eboracensis Septentrionali parte prope pontem Greta et Bignal vicum in sylvis ad fluvium Tefam observavit. D. Johnson. *R. Syn. prope* Kendal in Westmorlandia. 4. IV.

pyrenai- 2. ORNITHOGALUM racemo longissimo, filamentis lanceolatis, pedunculis proliferis patentibus æqualibus; fructiferis scapo aproximatis. *Sp. pl.* 307. Ornithogalum cum.

Ornithogalum angustifolium majus, floribus ex albo virefcentibus. *Bauh. pin.* 70. *R. Syn.* 373.

Asphodelus bulbosus. *Ger. Em.* 97.

Asphodelus bulbosus Galeni f. *Ornithogalum majus* flore subvirente. *Park. par.* 136.

Anglis, spiked star of Bethlehem with a greenish flower.

Habitat in the way between Bath and Bradford, not far from Little Ashley, *Ph. br.* On the left hand of a farm, half a mile from Chichester, Southgate, in a meadow, plentifully, Mr. Manningham. *R. Syn.* 24. V.

- *3. *ORNITHOGALUM* floribus corymbosis, pedunculis scapo altioribus, filamentis emarginatis. *Sp. umbellatum.* *pl.* 307.

Ornithogalum umbellatum medium angustifolium. *Bauh. pin.* 70.

Ornithogalum vulgare et verius, majus et minus. *Bauh. hist. II.* 630. *R. Syn.* 372.

Ornithogalum vulgare. *Ger. Em.* 165. *Park. par.* 136.

Anglis, common star of Bethlehem.

Habitat in sylvis, pratis et pascuis, in Norfolk. D. Stillingfleet. 24.

FRITILLARIA. 372.

Cor. hexapetala, campanulata, supra ungues cavitate nectarifera. *Stam.* longitudine corollæ.

- *1. *FRITILLARIA* foliis omnibus alternis. *Hort. ups. Meleagris.* 81. *Sp. pl.* 305.

Fritillaria præcox purpurea variegata. *Bauh. pin.* 64.

Fritillaria vulgaris. *Park. par.* 40.

Fritillaria variegata. *Ger. Em.* 149.

Anglis, common chequered Daffodil, or Fritillary.

Habitat in pratis et pascuis. In Mawde Fields, near Rislip common, Middlesex, plentifully. *Blackf. spec. bot.* 23. in pratis inter Mortlake et Kew. 24. IV. V.

ASPARAGUS. 382.

Ger. sexpartita, erecta : petalis tribus interioribus apice reflexis. *Bacca* trilocularis : disperma.

officinalis. 1. ASPARAGUS caule herbaceo erecto, foliis setaceis, stipulis paribus. *Sp. pl.* 313. *Fl. suec.* 291.

Asparagus. *Park.* 454. *Ger. Em.* 1110. *R. Syn.* 267. *fativus.* *Bauh. pin.* 489.

Anglis, Sparagus, or Sperage.

Habitat in pratis maritimis. Ad promontorium dictum
The Lizard Point *invenimus ; said also to be found*
in the marshes near Bristol ; about Harwich in Essex ;
and by the Thames side near Gravesend. R. Syn. 2.
VII. VIII.

Pharmac. Asparagi Radix.

CONVALLARIA. 383.

Cor. sexfida. *Bacca* maculosa, trilocularis.

majalis. 1. CONVALLARIA scapo nudo. *Fl. lapp.* 112. *suec.* 293. *Sp. pl.* 314.

Lilium convallium. *Ger. Em.* 410. *R. Syn.* 265. *album.* *Bauh. pin.* 304.

Lilium convallium flore albo. *Park. par.* 349.

Anglis, Lily-convally, or May-lily.

Habitat in ericetis frequens. 2. V.

Pharm. Lili Convallii Flores.

x *multiflora.* 2. CONVALLARIA foliis alternis amplexicaulibus, caule tereti, pedunculis axillaribus multifloris.
Phil. bot. 218. *Sp. pl.* 315. *Fl. suec.* 295.

Polygonatum. *Ger. Em.* 903. *R. Syn.* 263. *vulgare.* *Park.* 696.

Polygonatum latifolium vulgare. *Bauh. pin.* 303.

β *Polygonatum humile anglicum.* *R. Syn.* 263.

Anglis, Solomon's Seal.

Habitat in sylvis. In the woods by Brambdean in
Hampshire, Mr. Doody ; and on Roehill in
Kent, Mr. J. Sherard. R. Syn. in the Beech-
woods about High Wickham, Buckinghamshire,
Mr.

Mr. Hill. *Blackf. spec. bot. in sepibus prope Stock-*
well in comitatu Surrienfi. 2. V. VI. in most of the woods
Pharmac. Polygonati Radix. {about andover.

3. CONVALLARIA foliis alternis amplexicaulibus, *Polygona-*
caule ancipiti, pedunculis axillaribus subunifloris. tum.

Mat. med. 168. Fl. suec. 294. Sp. pl. 315.

Polygonatum floribus ex singularibus pediculis. Bauh.
hist. III. 529. R. Syn. 263.

Polygonatum latifolium flore majore odore. Bauh.
pin. 303.

Polygonatum majus flore majore. Park. 696.

Polygonatum latifolium 2. Clusii. Ger. Em. 904.

- β *Polygonatum Hellebori albi folio caule purpurascente.*
R. Syn.

Anglis, sweet-smelling Solomon's Seal.

Habitat in fissuris rupium prope Settle, Skipton, &c.
in comitatu Eboracenfi. 2. V. VI.

ANTHERICUM. 180.

Corolla hexapetala, patens. Caps. ovata.

1. ANTHERICUM foliis ensiformibus, perianthiis tri- *calycula-*
lobis, filamentis glabris. Fl. suec. 282. Sp. pl. 311. tum.

Phalangium scoticum palustre minimum, iridis folio.

Tourn. 369. R. Syn. 375.

Pseudo-asphodelus alpinum. Bauh. pin. 29.

Pseudo-asphodelus palustris Scoticus minimus. R. hist.

1195.

Anglis, Scottish Asphodel.

Habitat in palustribus. Juxta rivulum non procul Ber-
vico in Scotia. R. Syn. 2. IX.

NARTHECIUM.

CAL. nullus.

COR. hexapetala, erecta. *Petalis lineari-lanceolatis.*

STAM. *Filamenta sex, pubescentia. Antheræ parvæ*
incumbentes.

PIST. *Germen trigonum. Stylus nullus. Stigma obtusum.*

PER. *Capsula ovata, acuta, trigona, trilocularis, tri-*
valvis.

SEM. *numerosa fusiformia.*

NAR-

Offifragum.

- I. NARTHECIUM foliis ensiformibus filamentis lanatis. *Narthecium. Moebing. E. N. C. 1742.*

389. t. 5 f. 1.

Anthericum foliis ensiformibus, filamentis lanatis.

Sp. pl. 311.

Phalangium Anglicum palustre Iridis folio. Tourn.

368. *R. Syn. 375.*

Asphodelus Lancastriæ verus. Ger. Em. 97.

Pseudo-Asphodelus palustris anglicus. Bauh. pin. 29.

Anglis Lancashire Asphodel, or Bastard Asphodel.

Habitat in paludibus cespitosis. 2. VIII. near the XXXI mile stone on Bagshot Heath. Putney common, near the bowling green. P E P L I S. 402.

Perianthium campanulatum: ore duodecimfido. Petala sex, calyci inserta. Caps. bilocularis.

Portulaca.

- I. PEPLIS. *Fl. lapp. 328. succ. 312. sp. pl. 352.*

Portulaca. Dill. cat. giff. 120, 239. app. 133. R. Syn. 368.

Alfine palustris minor serpyllifolia. Bauh. pin. 120.

Alfine rotundifolia f. Portulaca aquatica. Ger. Em. 614.

Alfine aquatica minor folio oblongo, f. Portulaca aquatica. Park 1260.

Anglis Water Purslane.

Habitat in palustribus et aquosis ubi per hyemem aquæ stagnarunt. 2. IX.

*** Flores apetalæ.

A C O R U S. 392.

Spadix cylindricus, tectus flosculis. Cor. hexapetalæ, nudæ. Stylus nullus. Caps. trilocularis.

- Calamus. I. ACORUS. Roy. lugdb. 6. Fl. succ. 297. Sp. pl. 324.*

Acorus verus f. Calamus aromaticus off. Bauh. pin. 34.

Acorus verus f. Calamus officinarum. Park, 140. R. Syn. 437.

Acorus verus, Officinis falso Calamus. Ger. Em. 62.

Anglis, sweet-smelling Flag, or Calamus.

Habitat in fluviis funda coenosis. In the River Yare near Norwich, about Hedley in Surry, and in Cheshire.

R. Syn. in ericeto Hounsflejo. 2. V.

Pharmac. Calami aromatici Radix.

J U N-

JUNCUS. 396.

Cal. hexaphyllus. *Cor.* o. *Caps* unilocularis.

* *Culmis nudis.*

1. JUNCUS culmo subnudo tereti mucronato, panicula terminali, involucrio diphylo spinoso. *Sp. pl.* 325. *acutus.*
Juncus panicula universalis culmo brevior infime contractata, secundariis paululum laxis, capsulis triquetris subovatis nitentibus. *Guett. stamp. II.* 412.
Juncus acutus capitulis Sorghi. *Baub. pin.* 10. *R. Syn.* 430.
Juncus maritimus capitulis Sorghi. *Park* 1192.
 β *Juncus* panicula universalis culmo longior secundariis laxis longe pedunculatis media præcipue. *Guett. stamp. II.* 412.
Juncus acutus maritimus anglicus. *R. hist.* 1303. *Hist. Ox. III.* 232. *f.* 8. *t.* 10. *f.* 15.
 γ *Juncus parvus* cum pericarpiis rotundis. *Baub. hist.* II. 522. *R. Syn.* 433.
Anglis, Sea hard Rush.
Habitat in maritimis paludosis. 4. VII. VIII.
2. JUNCUS culmo nuduo stricto, capitulo laterali. *conglomeratus.*
Roy. lugdb. 44. *Sp. pl.* 326.
Juncus lævis vulgaris panicula compactiore. *R. Syn.* 432.
Juncus lævis panicula non sparfa. *Baub. pin.* 12.
Juncus lævis glomerato flore. *Lob. ic.* 84. *Park.* 1192.
Anglis, round-headed Rush.
Habitat in pratis et pascuis paludosis. 4. VI. VIII.
3. JUNCUS culmo nudo stricto, panicula laterali. *Roy. effusus.*
lugdb. 44. *Fl. suec.* 299. *Sp. pl.* 326.
Juncus culmo nudo acuminato ad basin squamato, floribus pedunculatis. *Fl. lapp.* 177.
Juncus lævis vulgaris panicula sparfa nostras. *R. Syn.* 432.
Juncus lævis panicula sparfa major. *Baub. pin.* 12. *Park.* 1191.
Juncus lævis. *Ger. Em.* 39.

K

Anglis,

Anglis, common soft Rush.

Habitat in paludosis. 2, V. VIII.

inflexus. 4. JUNCUS culmo nudo: apice membranaco incurvo, panicula laterali. *Sp. pl.* 326.

Juncus culmo nudo, paniculam arcuatum tegente. *Sauv. monsp.* 9.

Juncus panicula universali laterali culmo longiore secundariis sparfis. *Guett. stamp.* 412.

Juncus acutus panicula sparfa. *Bauh. pin.* 11.

Juncus acutus. *Ger. Em.* 39. *R. Syn.* 432. vulgaris. *Park.* 1193.

Anglis, hard Rush.

Habitat in pascuis et ad vias. 2. VI.

filiformis. 5. JUNCUS culmo nudo filiformi nutante, panicula laterali. *Roy. lugdb.* 44. *Gron. virg.* 152. *Fl. succ.* 300. *Sp. pl.* 326.

Juncus parvus, calamo supra paniculam longius productio. *R. Syn.* 432. *Fl. lapp.* 118. *Pluk. ph.* t. 40. f. 8.

Juncus lævis, panicula sparfa, minor. *Bauh. pin.* 12. *Sch. agr.* 347. t. 7. f. 1.

Anglis, the least soft Rush.

Habitat in montibus cespitosiss. Found by Mr. Newton, not far from Ambleside in Westmoreland. 2. VIII.

squarrosus. 6. JUNCUS culmo nudo, foliis fetaceis, capitulis glomeratis aphyllis. *Sp. pl.* 327. *Fl. succ.* 302.

Juncus alpinus, foliis rigidis sessilibus, caule nudo. *Rupp. jen.* 146.

Juncus montanus palustris. *R. Syn.* 432.

Gramen junceum maritimum. *Lob. ic.* 18. *Ger. Em.* 35. majus. *Park.* 1270.

Gramen foliis et spica junci. *Bauh. pin.* 5. 10. tb. 78. *Anglis*, Moss-rush or Goose-corn.

Habitat in ericetis frequens. 2. VI.

** Culmis foliosis.

articulatus. 7 JUNCUS foliis nodoso-articulatis. *Fl. succ.* 305. *Sp. pl.* 327.

JUNCUS

- Juncus foliis articulosis, panicula inæquali, capitulis sessilibus capsulis subulatis triquetris.* *Roy. lugdb.* 43.
- Juncus foliis articulosis floribus umbellatis.* *Tourn.* 247. *R. Syn.* 433.
- Gramen junceum folio articulato aquaticum* *Baub.* *pin.* 5.
- Gramen aquaticum.* *Tab. ic.* 214. *Ger. Em.* 22.
- Gramen junceum aquaticum.* *Bauhini* 1270.
- β *Juncus nemorosus folio articulofo.* *Tourn.* 247. *R. Syn.* 433.
- Anglis, jointed leaved Rush.*
- Habitat in sylvis, pratis et pascuis humidis.* 4. VII. VIII.
8. *JUNCUS foliis linearibus canaliculatis capsulis ob-* *bulbosus.*
tufis. *Fl. suec.* 304. *Sp. pl.* 327.
- Juncus foliis angulatis, culmo subnudo, panicula terminali prolifera, capsulis ovatis.* *Roy. lugdb.* 43.
- Gramen junceum minimum capsulis triangularibus.*
R. Syn. ed. II. 275. *Hist. Ox. III.* 227. *J. 8. t. 9. f. 3.* *R. Syn.* 434.
- Graminis juncei varietas minor.* *Ger. Em.* 4.
- Anglis, bulbose Rush.*
- Habitat in ericetis humidis frequens.* 4. VIII.
9. *JUNCUS culmo dichotomo, foliis angulatis, flori-* *bufonius.*
bus solitaris sessilibus. *Roy. lugdb.* 43. *Fl. suec.* 305.
Sp. pl. 328.
- Juncus palustris humilior erectior.* *Tourn.* 246. *R. Syn.* 434.
- Gramen junceum.* *Ger. Em.* 4. *Parvum, f. Holosteum Matthioli, et Gramen bufonium Flandrorum.* *Park.* 1190.
- Gramen nemorosum calyculis paleaceis.* *Baub. pin.* 7.
- β *Gramen juncoides minimum Anglo-Britannicum, Holosteum Matthioli congener, aut bufoniæ gramine Flandrico.* *Lob. Ill.* 70.
- Anglis, Toad-rush.*
- Habitat in pratis et pascuis arenosis, præcipue ubi aquæ pluviales stagnant.* ①. VII. VIII.

pilosus.

10. JUNCUS foliis planis pilosis corymbo subramoso.
Fl. suec. 308.
 Gramen nemorosum hirsutum vulgare. *R. Syn.* 416.
 Gramen nemorosum hirsutum. *Ger. Em. Lob. ic.*
 16 majus. *Park.* 1184.
 Gramen nemorosum hirsutum majus latifolium. *Bauh.*
pin. 7. *th.* 101. *Hist. Ox.* 111. 225. *f.* 8. *t.* 9. *f.* 1.
 Cyparella nemorensis, capitulis in singulis petiolis sin-
 gulis latifolia. *Rupp.* 116.
Anglis, common hairy Wood-rush, or Grass.
 Habitat in sylvis densis frequens. In sylva Charlton-
 wood dicta in Cantio. 4. IV. V.

sylvaticus.

11. JUNCUS foliis planis pilosis, umbella supra decom-
 posita floribus subsessilibus.
 Juncus foliis planis latissimis, panicula sparsa, dupli-
 cato ramoso. *Hall. helv.* 257.
 Juncus foliis planis, panicula diffusa, pedunculis pro-
 liferis, flosculis intermediis sessilibus. *Roy. lugdb.*
 43.
 Gramen nemorosum hirsutum latifolium maximum.
R. Syn. 416.
 Gramen Luzulæ maximum. *Bauh. hist.* II. 493.
Anglis, great hairy Wood-rush.
 Habitat in sylvis et sepibus prope Hamstead. 4. V.

campestris.

12. JUNCUS foliis planis pilosis, spicis sessilibus pe-
 dunculatisque. *Fl. suec.* 309. *Sp. pl.* 329.
 Gramen hirsutum capitulis psyllii. *Bauh. pin.* 7. *pr.*
 15.
 Gramen nemorosum hirsutum minus angustifolium.
Park. 1185.
 Gramen exile hirsutum. *Ger. Em.* 16. *R. Syn.* 416.
 β Gramen hirsutum elatius, panicula juncea compacta.
R. Syn. 416.
Anglis, small hairy Wood-rush.
 Habitat in pratis et pascuis siccioribus, β in paludibus
 turfosis. 4. IV.

TRIGYNIA.

* *Flores inferi.*

TRIGLOCHIN. 409.

Cal. triphyllus. *Petala* tria calyciformia. *Stylus* o.
Capf. basi dehiscens.

1. TRIGLOCHIN capsulis trilocularibus sublinearibus. *palustre*.
Fl. suec. 321. *Sp. pl.* 338.

Juncajo palustris et vulgaris. *Tourn.* 266. *R. Syn.* 435.

Gramen trigloch. *Baub. hist.* II. 508.

Gramen junceum spicatum seu trigloch. *Baub. pin.* 6.

Gramen marinum spicatum. *Ger. Em.* 18. alterum,
Park. 1279.

Anglis, Arrow-headed Grass.

Habitat in pratis et pascuis udis passim. 4. VII. VIII.

2. TRIGLOCHIN capsulis sexlocularibus ovatis. *Fl. mariti-*
suec. 322. *Sp. pl.* 339. *mum.*

Gramen marinum spicatum. *Lob. ic.* 16. *Park.* 1279.

R. Syn. 435.

Gramen junceum et spicatum alterum. *Baub. pin.* 6.

Anglis, Sea spiked Grass.

Habitat ubique in pratis maritimis. 4. V. VI.

RUMEX 407.

Cal. triphyllus. *Petala* tria, conniventia. *Sem.* unicum triquetrum.

* *Hermaphroditi* : valvulis granulo notatis.

1. RUMEX floribus hermaphroditis : valvulis integer. *sanguini-*
rimis : unica granifera, foliis cordato lanceolatis. *neus.*
Sp. pl. 334.

Lapathum folio acuto rubente. *Baub. pin.* 115. *R. Syn.* 142.

Lapathum fativum sanguineum. *Ger. Em.* 390.

Anglis, Bloodwort.

K 3

Habitat

Habitat in sylvis prope Hamstedium. ♂. VII.
Pharmac Lapathi sanguinei *Folia, Semina.*

- acutus.* 2. RUMEX floribus hermaphroditis : valvulis dentatis graniferis, foliis cordato-oblongis. *Hort cliff.* 138. *Fl. suec.* 316. *Sp. pl.* 335. *Mat. med.* 176.
 Lapathum acutum. *Ger. Em.* 388. *R. Syn.* 142. majus. *Park.* 1224.
 Lapathum folio acuto, plano. *Bauh. pin.* 115.
Anglis, sharp-pointed Dock.
 Habitat in incultis, sylvis et aquosis. 4. VI.
Pharmac. Oxylapathi, *Radix, Semina.*
- crispus.* 3. RUMEX floribus hermaphroditis : valvulis integris graniferis, foliis lanceolatis undulatis acutis. *Sp. pl.* 335.
 Lapathum folio acuto crispo. *Bauh. pin.* 115. *R. Syn.* 141.
 Lapathum acutum minus. *Park.* 226.
 Lapathum acuti varietas folio crispo. *Ger. Em.* 387.
Anglis, curled Dock.
 Habitat in pratis pascuis et ad vias. 4. VI. VII.
- obtusifolium.* 4. RUMEX floribus hermaphroditis : valvulis dentatis, foliis cordato-oblongis obtusiusculis crenulatis. *Sp. pl.* 335.
 Lapathum vulgare folio obtuso. *Bauh. hist. II.* 984. *R. Syn.* 141.
 Lapathum sylvestre folio subrotundo. *Bauh. pin.* 115.
 Lapathum sylvestre vulgatus. *Park.* 1225.
 Lapathum sylvestre folio minus acuto. *Ger. Em.* 388.
Anglis, broad-leaved Dock.
 Habitat ad vias et in locis humidis. 4. VII. VIII.
- pulcher.* 5. RUMEX floribus hermaphroditis : valvulis dentatis graniferis, foliis radicalibus lyratis. *Syst. nat.* 990. *Sp. pl.* 336.
 Lapathum pulchrum Bononienae sinuatum. *Bauh. hist. II.* 288. *R. Syn.* 142.
 Lapathum ramis procumbentibus, feminis involucreto dentato, foliis inferioribus instar fidium. *Hist. Ox* II. 580.

Anglis,

Anglis, Fiddle-dock.

Habitat *ad vias et in pratis siccis*. 4. VI.

6. RUMEX floribus hermaphroditis : valvulis dentatis *maritigraniferis*, foliis linearibus. *Lech. scan.* 26. *Sp. pl. mus.*

335.

Lapathum folio acuto flore aureo. *Bauh. pin.* 115.

R. Syn. 142.

Lapathum aquaticum, angustissime acuminato folio.

Bocc. mus. II. 142. *t.* 104.

Lapathum aureum glomerulis densis. *Pet. h. br. t.*

2. *f.* 8.

Anthoxanthon. *Bauh. hist. II.* 988.

- β Lapathum aureum. *Pet. h. br. t.* 2. *f.* 7.

Anglis, Golden Dock.

Habitat *ad vias et in ruderatis passim*. 4. VII. VIII.

7. RUMEX floribus hermaphroditis : valvulis integer-*Britanirimis*, omnibus graniferis, foliis lanceolatis vaginis *ca.* obsoletis. *Mat. med.* 17. *Sp. pl.* 334.

Lapathum maximum aquaticum f. Hydrolapathum.

Bauh. hist. II. 986. *R. Syn.* 140.

Lapathum aquaticum folio cubitali. *Bauh. pin.* 116.

Lapathum longifolium nigrum palustre, f. *Britanica antiquorum vera*, vel Hydrolapathum nigrum. *Munting. br. t.* 1.

Hydrolapathum magnum. *Ger. Em.* 389.

Hydrolapathum majus. *Park.* 1225.

Anglis, great Water Dock.

Habitat *in fossis, palustribus frequens*. 4. VII. VIII.

Pharmac. Hydrolapathi Radix.

** *Hermaphroditi : valvulis granulo destitutis.*

8. RUMEX floribus hermaphroditis digynis. *Hort. digynus. cliff.* 138. *Fl. suec.* 317. *Sp. pl.* 337.

Rumex foliis orbiculatis emarginatis. *Fl. lapp.* 132.

Acetosa rotundifolia repens Eboracensis, folio in medio deliquium patiente. *Hist. Ox. II.* 583. *f.* 5. *t.* 36. *f. penult.*

Acetosa repens Westmorlandica, cochleariæ foliis apicibus nonnihil sinuatis. *Pluk. ph. t.* 252. *f.* 2.

Acetosa Cambro-britanica montana. *Park.* 745.

Anglis, round-leaved mountain Sorrel.
Habitat in montibus Wallicis, Eboracensi, Westmorlandico, Cumberlandico, frequens. 4. VI.

*** *Floribus declinis.*

- Acetosa.* 9. RUMEX floribus dioicis, foliis oblongis sagittatis.
Fl. lapp. 130. *succ.* 318. *Sp. pl.* 337.
 Lapathum acetosum vulgare. *R. Syn.* 143.
Acetosa vulgaris *Park.* 742.
Acetosa pratensis. *Bauh.* *pin.* 114.
Oxalis, seu acetosa. *Ger. Em.* 396.
 β *Acetosa montana maxima.* *Bauh.* *pin.* 114. *R. Syn.* 143.
Anglis, common Sorrel.
Habitat in pratis et pascuis, β in rupibus maritimis prope Harlech in comitatu Merionethensi collegit. D. Lhwyd. *R. Syn.*
Pharmac. *Acetosæ pratensis et oxalis Radix, Folia, Semina.*

- Acetofella.* 10. RUMEX floribus dioicis, foliis lanceolato-hastatis.
Fl. lapp. 131. *succ.* 319. *Sp. pl.* 338.
 Lapathum acetosum repens lanceolatum. *R. Syn.* 143.
Acetosa arvensis lanceolata. *Bauh.* *pin.* 114.
Acetosa minor lanceolata. *Park.* 744.
Oxalis tenuifolia. *Ger. Em.* 397.
Anglis, Sheep's Sorrel.
Habitat in arvis, pratis, pascuis, arenosis. 4. V. VI. VII.
Pharmac. *Acetosæ lanceolatæ Folia.*

COLCHICUM. 415.

Spatha. *Cor.* sexpartita; tubo radicato. *Caps.* tres connexæ, inflatæ.

- autum-nale.* 1. COLCHICUM foliis planis lanceolatis erectis. *Sp. pl.* 341.
Colchicum commune. *Bauh.* *pin.* 67. *R. Syn.* 373.
Colchicum Anglicum purpureum et Anglicum album.
Park. 153. *Ger. Em.* 157.
Anglis, Meadow Saffron.

Habitat

Habitat in pratis tam planis quam montosis, solo pingui,
in Occidentalibus ut et Septentrionalibus Angliæ.
R. Syn. 4. IX.

P O L Y G Y N I A.

A L I S M A. 418.

Cal. triphyllus. Petala tria. Sem. plura.

1. ALISMA foliis ovatis acutis, fructibus obtusis trigono-
Plantaginis. Sp. pl. 342. Fl. suec. go. Δ.
Plantago aquatica. Baub. hist. III. 787. R. Syn.
257. major. Ger. Em. 417. Park. 1245.
Plantago aquatica latifolia. Baub. pin. 190.
Anglis, great Water Plantain.
Habitat in aquosis et ad ripas fluviorum ubique. 4. VI.
VII.
2. ALISMA foliis lineari-lanceolatis, fructibus globofo-
squammosis. Sp. pl. 343. Fl. suec. 325. ranunculoides.
Alisma fructu globofo, undique echinato. Hort. cliff.
141.
Plantago aquatica minor. Park. 1245. R. Syn. 257.
Plantago aquatica humilis. Ger. Em. 417.
Plantago aquatica angustifolia. Baub. pin. 190.
Anglis, lesser Water Plantain.
Habitat in aquosis sed rarius. 4. VI.—VIII.
3. ALISMA foliis ovatis obtusis, pedunculis solitariis. natans.
Sp. pl. 343. Fl. suec. 324.
Alisma repens, foliis gramineis et subrotundis. Mart.
Tourn. 17.
Damaſonium repens, potamogetonis rotundifolii folio.
Vaill. Aët. 1719. p. 29. t. 4. f. 8.
Anglis, creeping Water Plantain.
Habitat in the great lake, below the old castle, at Llan-
berys. Mr. Brewer. Mart. Tourn. 4. VIII.

4. ALISMA

- Damafo-* 4. ALISMA foliis cordato-oblongis, floribus hexagynis.
nium. *Sp. pl.* 343.
Alisma fructu sexcorni. Hort. cliff. 142. *Roy. lugdb.*
 46.
Plantago aquatica minor stellata. Ger. Em. 417.
Plantago aquatica stellata. Baub. pin. 190.
Plantago aquatica minor muricata. Park. 1245.
Damaconium stellatum Dalechampii, Baub. hist. III.
 789. *K. Syn.* 272.
Anglis, Star-headed Water Plantain.
Habitat in fossis et aquis stagnantibus. Circa Londi-
num frequens. 2. VI.—VIII.

Classis

Classis VII.

HEPTANDRIA

MONOGYNIA.

TRIENTALIS. 419.

Cal. heptaphyllus. *Cor.* septem-partita, æqualis,
plana. *Bacca* exsucca.

1. TRIENTALIS foliis lanceolatis integerrimis. *Kalm. europæa.*
canad. II. 263. *Fl. suec.* 326. *Sp. pl.* 344.

Alfinanthemos. *Thal. sylv.* 15. *Moerb. hort.* 8. III.
R. Syn. 286.

Pyrola alfines flore europæa. *Bauh. pin.* 191. *Park.*
509.

Pyrola alfines flore major. *Bauh. prod.* 100.

Herba trientalis. *Bauh. hist.* III. 506.

Alfine alpina. *Schwenk. filsf.* 17. *descr.*

β *Pyrola alfines flore Brasiliana.* *Bauh. pin.* 191. *pr.*
100.

Anglis, Chickweed, Winter-green.

Habitat in sylvis, et ericetis in comitatibus Eboracensi,
Northumberlandico et Westmorlandico; prope
Bingley in comitatu Eboracensi copiose. 4. VI.

Classis

Classis VIII.

OCTANDRIA
MONOGYNIA.

* Flores completi.

EPILOBIUM. 426.

Cal. quadrifidus. *Petala* quatuor. *Capf.* oblonga,
infera. *Sem.* papposa.* *Staminibus declinatis.*X *angustifolium.* 1. EPILOBIUM foliis sparsis lineari-lanceolatis. *Sp. pl.*
347. *Fl. suec.* 327.*Epilobium* floribus difformibus, pistillo declinato. *Fl.*
suec. Ed. I. 304.*Lyfimachia Chamænerion* dicta angustifolia. *Bauh.*
pin. 245.*Lyfimachia speciosa*, quibusdam *Onagra* dicta, fili-
quosa. *Bauh. hist. II.* 906. *R. Syn.* 310.*Chamænerion flore Delphinii.* *Park.* 548.*Anglis*, Rosebay Willow-herb.*Habitat in sylvis et sepibus. In the meadows near*
Sheffield in Yorkshire, and in several other places
of the north plentifully. R. Syn. On Maize-hill be-
yond Greenwich. 4. VII. VIII. in a little Ash Grove
*by the Meadow near Charlton in Andover parish. Hants*** *Staminibus erectis regularibus, petalis bifidis. Plentifully**hirsutum.* 2. EPILOBIUM foliis lanceolatis undulato-ferratis de-
currentibus; inferioribus oppositis, caule subsim-
plici, racemo terminalibus.*Epilobium* foliis oppositis lanceolatis ferratis decur-
renti-amplexicaulibus. *Sp. pl.* 347. *Fl. suec.* 328.*Lyfimachia filiquosa hirsuta* parvo flore. *Bauh. pin.*
245. *pr.* 116. *R. Syn.* 311.*Lyfimachia sylvatica.* *Ger. Em.* 476.*Lyfimachia filiquosa sylvestris hirsuta.* *Park.* 549.
Anglis,

Anglis, small flowered hairy Willow-herb.

Habitat in *aquosis et ad rivulos*. 4. VII.

3. *EPILOBIUM* foliis lanceolatis ferratis subdecurren- *ramosum*.
tibus inferioribus oppositis, caule ramoso.

Lyfimachia filiquosa hirsuta magno flore. *Bauh. pin.*

245. *R. Syn.* 311.

Lyfimachia filiquosa. *Ger. Em.* 476.

Anglis, great flowered Willow-herb, or Codlings and Cream.

Habitat in *sepibus humidis et ad rivulos passim*. 4. VII.

4. *EPILOBIUM* foliis oppositis ovatis dentatis. *Fl. lapp. montanum*
147. *suec.* 329.

Lyfimachia campestris. *Ger. Em.* 478.

Lyfimachia filiquosa glabra major. *Bauh. pin.* 245.

Lyfimachia filiquosa major. *Park.* 548.

Anglis, smooth-leaved Willow-herb.

Habitat in *pratis humidis glareosis frequens*. 8. VI.
VII.

5. *EPILOBIUM* foliis lanceolato-linearibus denticu- *tetrago-*
latis oppositis, caule tetragono. *Sauv. monsp.* 75. *num.*

Lyfimachia filiquosa glabra media f. minor. *Ger. Em.*

479. *R. Syn.* 311.

Lyfimachia filiquosa glabra minor. *Bauh. pin.* 245.

Anglis, narrow-leaved Willow-herb.

Habitat *ad fossas et rivulos*. 4. VII.

6. *EPILOBIUM* foliis oppositis lanceolatis integer- *palustre*.
rimis, petalis emarginatis caule erecto. *Sp. pl.*

348. *Fl. suec.* 330.

Lyfimachia filiquosa glabra minor angustifolia. *Ger.*

Em. 479. *R. Syn.* 311.

Lyfimachia filiquosa glabra angustifolia. *Bauh. pin.*

245.

Anglis, marsh Willow-herb.

Habitat in *palustribus*. 4. VII.

- alpinum.* 7. **EPILOBIUM** foliis oppositis ovato-lanceolatis integerrimis, filiquis sessilibus, caule repente. *Sp. pl.* 348. *Fl. suec.* 331.
Chamænerion alpinum alpinæ foliis. *Sch. alp.* 132. 332.
Lyfimachia filiquosa glabra minor latifolia. *R. Syn.* 311.
Anglis, Mountain Willow-herb.
Habitat ad rivulos. In the rivulet on the sides of Cheviot-Hills in Northumberland. *R. Syn.* 4. VII.

VACCINIUM. 434.

Cal. superus. *Cor.* monopetala. *Filamenta* receptaculo inserta. *Bacca* quadrilocularis, polysperma.

* *Foliis* annotinis *f. deciduis.*

- Myrtillus.* 1. **VACCINIUM** pedunculis unifloris, foliis ferratis ovatis deciduis, caule angulato. *Sp. pl.* 350. *Fl. suec.* 333.
Vitis idæa, foliis oblongis crenatis, fructu nigricante. *Bauh. pin.* 470.
Vitis idæa angulosa. *Bauh. hist. I.* 520. *R. Syn.* 457.
Vaccinia nigra. *Ger. Em.* 1415. *vulgaria.* *Park.* 1455.
Anglis, Black Whorts, Whortle-Berries, or Bilberries.
Habitat in ericetis et sylvis frequens. *h.* IV.
- uliginosum.* 2. **VACCINIUM** pedunculis unifloris, foliis integerrimis ovalibus basi ciliatis. *Sp. pl.* 350. *Fl. suec.* 332.
Vitis idæa foliis subrotundis exalbidis. *Bauh. pin.* 470.
Vitis idæa magna quibusdam, *f. Myrtyllus* grandis. *Bauh. hist. I.* 518. *R. Syn.* 457.
Vitis idæa foliis subrotundis major. *Ger. Em.* 1416.
Vaccinia nigra fructu majore. *Park.* 1455.
Anglis, the great Bilberry Bush.
Habitat in montosis Cumberlandiæ medio itinere inter Hexham et Pereth oppida; sed copiosius prope vicum Gambleby, ad sextum ab oppido Pereth lapidem, via quæ Novum Castellum ducit: utrobique in passuis palustribus, &c. *R. Syn.* *h.* IV. V.

3. VAC-

3. VACCINIUM racemis nudis, foliis integerrimis re- *cantabri-*
volutis lanceolato-ovatis, subtus tomentosis. *cum.*

Erica cantabrica flore maximo, foliis myrti subtus incanis. *Tourn.* 603. *R. Syn.* 472.

Erica St. Dabeoci Hibernis. *D. Lhwyd. R. Hist.* III. *dendr.* 98.

Erica Hibernica foliis myrti pilosis, subtus incanis.

R. Hist. III. *App.* 244. *Pet. gaz. nat.* t. 27. f. 4. *Anglis*, Irish Whorts.

Habitat in montibus Mayo squalido et spongioso solo frequens, ut et per totum Hair-Connacht in Gallovidia.

R. Syn. h.

** *Foliis sempervirentibus.*

4. VACCINIUM racemis terminalibus nutantibus, foliis ob ovatis revolutis integerrimis subtus punctatis. *Vitis idæa.*
Sp. pl. 351.

Vaccinia rubra. *Ger. Em.* 1415. buxeis foliis. *Park.* 1458.

Vitis idæa foliis subrotundis non crenatis, baccis rubris. *Bauh. pin.* 470.

Vitis idæa semper-virens fructu rubro. *Bauh. hist.* I. 522. *R. Syn.* 457.

Anglis, red Whorts, or Whortle-berries.

Habitat in montibus Eboracensibus, Derbiansibus, Staffordiensibus inter ericas copiose. *R. Syn.* h.

IV. V.

5. VACCINIUM foliis integerrimis revolutis ovatis, *Oxycoccus.*
caulibus repentibus filiformibus nudis. *Sp. pl.* 351.
Fl. suec. 335.

Vaccinium palustre. *Park.* 1229.

Vaccinia palustris. *Ger. Em.* 1425.

Vitis idæa palustris. *Bauh. pin.* 471.

Oxycoccus seu *Vaccinia palustris.* *Bauh. hist.* I. 525.
R. Syn. 267.

Anglis, Cran-berries, Moss-berries, or Moor-berries.

Habitat in palustribus turfosis in comitatibus Westmorlandico, Cumberlandico, Lancastriensi, Eboracensi, &c. copiose. h. V. VI.

ERICA.

ERICA. 435.

Cal. tetraphyllus. *Cor.* quadrifida. *Filamento* receptaculo inserta. *Antheræ* bifidæ. *Capf.* quadrilobularis.

* *Antheris bicornibus.*

vulgaris. 1. ERICA antheris bicornibus inclusis corollis inæqualibus campanulatis mediocribus, foliis oppositis sagittatis. *Sp. pl.* 352. *Fl. suec.* 336.

Erica foliis quadrifariam imbricatis triquetris glabris erectis, corollis inæqualibus calyce brevioribus. *Hort. cliff.* 146.

Erica vulgaris. *Park.* 1480. *R. Syn.* 470. *f. pumila.* *Ger. Em.* 1380.

Erica vulgaris glabra. *Bauh. pin.* 485.

Anglis, common Heath, or Ling.

Habitat in ericetis. *h.* VI.—VIII.

cinerea. 2. ERICA antheris bicornibus inclusis, corollis ovatis racemosis, foliis ternis glabris linearibus. *Sp. pl.* 352.

Erica tenuifolia. *Ger. Em.* 1380. *R. Syn.* 471.

Erica humilis, cortice cinereo, Arbuti flore. *Bauh. pin.* 486.

Erica virgata *f.* VI. Clusii. *Park.* 1483.

Anglis, fine-leaved Heath.

Habitat in ericetis et sylvis siccissimis. *h.* VI. VIII.

Tetralix. 3. ERICA antheris bicornibus inclusis, corollis subglobofis aggregatis calyce longioribus foliis quaternis ciliatis patentibus. *Sp. pl.* 353. *Fl. suec.* 337.

Erica major flore purpureo. *Ger. Em.* 1382.

Erica ex rubro nigricans scoparia. *Bauh. pin.* 486.

Erica brabantica folio coridis hirsuto quaterno. *Bauh. hist.* I. 358. *R. Syn.* 471.

Anglis, cross-leaved Heath.

Habitat in ericetis humidis. *h.* VII. VIII.

** *Antheris simplicibus obtusis emarginatis.*

ciliaris. 4. ERICA antheris simplicibus inclusis, corollis ovatis irregularibus, floribus terno-racemosis, foliis ternis ciliatis. *Sp. pl.* 354.

Erica

Erica vulgaris hirsuta. Ger. Em. 1382. R. Syn. 471.

Erica hirsuta anglica. Bauh. pin. 602.

Erica vulgaris hirsutior. Park. 1480.

Anglis, rough-leaved Heath.

Habitat in ericetis passim. h. VI.

5. *ERICA* antheris bifidis simplicibus exsertis, corollis *multifloras* cylindricis longioribus, foliis quinis linearibus patentibus. Sp. pl. 355.

Erica foliis Corios multiflora. Bauh. hist. I. 356. R. Syn. 471.

Erica juniperifolia narbonensis dense fruticaris. Lob. hist. 620. Sauv. monsp. 46.

Anglis, Fir-leaved Heath.

Habitat in incultis. On Goon hilly downs going from Helston to the Lizard Point in Cornwall. R. Syn. h. VI. VII.

** Flores incompleti.

D A P H N E. 436.

Cal. o. Cor. quadrifida, marcescens, stamina includens. Bacca monosperma.

1. *DAPHNE* racemis axillaribus, foliis lanceolatis glâ- *Laureola.* bris. Hort. cliff. 147. Mat. med. 180. Sp. pl. 357.

Laureola. Ger. Em. 1405. Park. 205. R. Syn. 465.

Laureola semper-virens flore viridi, quibusdam Laureola mas. Bauh. pin. 462.

Anglis, Spurge Laurel.

Habitat in sylvis et sepibus. In sepibus inter Wimbles-ton et Martini. h. III. IV.

Pharmac. Laureolæ Folia, Bacca.

2. *DAPHNE* floribus sessilibus ternis caulinis, foliis lanceolatis deciduis. Sp. pl. 356. Fl. succ. 338. *Mezereum.*

Laureola folio deciduo, flore purpureo. Bauh. pin. 426.

Camælæa Germanica f. *Mezereon.* Ger. Em. 1403. vulgo. Park. 205.

Anglis, Spurge Olive.

Habitat in sylvis prope Andover copiose. D. Miller. h. III.

Pharmac. Mezerei Folia, Bacca, Cortex.

D I G Y N I A.

B L A C K S T O N I A.

CAL. *Perianthium* octofidum : *laciniis* lanceolatis persistentibus.

COR. monopetala. *Tubus* nullus. *Limbus* octofidus : *laciniis* lanceolatis longitudine calycis.

STAM. *Filamenta* octo, subulata. *Antheræ* simplices.

PIST. *Germen* ovato-oblongum. *Stylus* nullus. *Stigmata* duo.

PER. *Capfula* oblonga, acuminata, unilocularis, bivalvis.

SEM. numerosa, parva.

x *perfoliata*. I. BLACKSTONIA.

Gentiana corollis octofidis, foliis perfoliatis. *Sp. pl.* 232.

Centaureum luteum perfoliatum. *Bauh. pin.* 278. *hist.* III. 355. *R. Syn.* 287.

Centaureum parvum luteum Lobelii. *Ger. Em.* 547.

Centaureum minus luteum et perfoliatum non ramosum. *Park.* 271.

Anglis, yellow Centory.

Habitat in pratis et pascuis montosis. ☉. VII.

Plentifully about Doverdale in Worcestershire.

T R I G Y N I A.

P O L Y G O N U M. 445.

Cal. o. *Cor.* quinquepartita, calycina. *Sem.* unicum angulatum.

* *Bistortæ spica unica.*

Bistorta. I. POLYGONUM caule simplicissimo monostachyo, foliis ovatis in petiolum decurrentibus. *Mat. med.* 188. *Sp. pl.* 360.

Bistorta major. *Ger. Em.* 399. *R. Syn.* 147. vulgaris. *Park.* 391.

Bistorta radice minus intorta. *Bauh. pin.* 192.

Anglis,

Anglis, the greater Bistort, or Snakeweed.

Habitat in pratis humidis in comitatibus Cumberlandico, Westmorlandico, Eboracensi, frequens; prope Battersea. 24. V. VI.

Pharmac. Bistortæ Radix.

2. POLYGONUM caule simplicissimo monostachyo, vivipar. foliis lanceolatis. *Sp. pl.* 360. *Fl. suec.* 340. *rum.*

Bistorta minor. *Ger. Em.* 399. *R. Syn.* 147. nostras. *Park.* 392.

Bistorta alpina minor. *Bauh. pin.* 192.

- β Bistorta minima alpina, foliis imis subrotundis et minutissime ferratis. *R. Syn.* 147.

Anglis, small Bistort, or Snakeweed.

Habitat In pratis montosis. In several places in Westmoreland, and near Settle in Yorkshire. *R. Syn.* 24. VI.

** *Perficariæ pistillo bifido, aut stamina minus 8.*

3. POLYGONUM floribus pentandris semidigynis, amphibistaminibus corolla longioribus. *Sp. pl.* 361. *Fl. suec.* 341. *um.*

Perficaria falicis folio perennis, Potamogiton angustifolium dicta. R. Syn. 145.

Potamogiton angustifolium. Ger. Em. 821.

Potamogiton falicis folio. Bauh. pin. 131.

Fontalis major longifolia. Park. 1254.

Anglis, Perennial Arsmart.

Habitat in fossis et paludibus frequens. 24. VI. VII.

4. POLYGONUM floribus hexandris digynis, spicis *Perficaria* ovato-oblongis, foliis lanceolatis stipulis ciliatis. *Sp. pl.* 361.

Perficaria maculosa. Ger. Em. 445. *R. Syn.* 145.

Perficaria mitis maculosa et non maculosa. Bauh. pin. 101.

Perficaria vulgaris mitis feu maculosa. Park. 856.

- β *Perficaria angustifolia. Bauh. pin.* 101. *pr.* 43. *ex* singulis geniculis florens. *Merr. pin. R. Syn.* 145.

- γ *Perficaria folio subtus incano. Tourn.* 510. *R. Syn.* 145.

OCTANDRIA TRIGYNIA.

♂ *Perficaria foliis falicis albæ vulgaris.* *Buddl. hort. succ.*
Vol. IV. f. 20. R. Syn. 145.

ε *Perficaria latifolia geniculata, caulibus maculatis.* *R.*
Syn. 145.

ζ *Perficaria maculosa procumbens foliis subtus incanis.*
R. Syn. 146.

Anglis, dead or spotted Arsmart.

Habitat in pratis pascuis, arvis, ruderatis humidis ubique. γ. δ. in arenosis aridis. ☉. VIII. IX.

pensylvanicum. 5. **POLYGONUM** floribus octandris digynis, pedunculis hispidis, foliis lanceolatis, stipulis muticis. *Sp. pl. 362.*

Perficaria mitis major, foliis pallidioribus. D. Bobarti.
R. Syn. 145.

Anglis, pale Arsmart.

Habitat in arvis et ruderatis frequens. ☉. VIII.

Hydropiper. 6. **POLYGONUM** floribus hexandris semidigynis, foliis lanceolatis, stipulis submuticis. *Sp. pl. 361.*
Fl. succ. 343.

Perficaria urens seu Hydropiper. Baub. pin. 101.

Perficaria vulgaris acris f. Hydropiper. Baub. hist.
III. 780. R. Syn. 144.

Perficaria vulgaris acris seu minor. Park. 856.

Hydropiper. Ger. Em. 445.

Anglis, Water-pepper, or Arsmart.

Habitat in aquis et ad rivulos et fossas. ☉. VII. VIII.
Pharmac. Perficariæ non maculatæ Herba.

minus. 7. **POLYGONUM** floribus hexandris digynis, foliis lanceolatis, stipulis ciliatis, caule divaricato patulo.
Perficaria pusilla repens. Ger. Em. 446. Park. 857.
R. Syn. 145.

Perficaria minor. Baub. pin. 101. Hist. Ox. II. f. 5.
t. 29. f.

Anglis, small creeping Arsmart.

Habitat in aquis et ad fossas sed rarius. In the meadows beyond Peterborough-house, and in Tothill-fields, Westminster. Blackf. spec. bot. ☉. VII.

Poly-

Polygona foliis indivisis, floribus octandris.

1. POLYGONUM floribus octandris trigynis axillari- *aviculare*.
bus, foliis lanceolatis, caule procumbente herbaceo.
Sp. pl. 392. *Fl. suec.* 339.

Polygonum mas vulgare. *Ger. Em.* 565. *R. Syn.*
146. majus. *Park.* 443

Polygonum latifolium. *Bauh. pin.* 281.

Polygonum feu Centinodia. *Bauh. hist. III.* 374.

β Polygonum brevi angustoque folio. *Bauh. pin.* 281.

γ Polygonum oblongo angustoque folio. *Bauh. pin.*
281.

δ Polygonum folio rotundo. *Pet. h. br. t.* 10. f. 2.

Anglis, Knot-grafs.

Habitat in viis, plateis, agris ubique. ☉. VI. IX.

Pharmac. Polygoni f. Centinodiæ Herba.

9. POLYGONUM floribus octandris trigynis axillari- *mariti-*
bus, foliis ovati-lanceolatis, caule suffruticoso sem- *um*.
pervirente. *Sp. pl.* 361.

Polygonum marinum. *Bauh. hist. III.* 376. *R. Syn.*
147.

Polygonum marinum maximum. *Ger. Em.* 564.

Polygonum maritimum latifolium. *Bauh. pin.* 281.

Polygonum marinum majus. *Park.* 444.

Anglis, Sea Knot-grafs.

Habitat in littoribus maritimis. Prope oppidum Pen-
fans in Cornubia, between Abermeney and Lihan-
dwyn, Mr. Lhwyd; at Brackelsham in Suffex, Dr.
Dillenius. R. Syn. 4. VII.

*** *Helxine foliis subcordatis.*

10. POLYGONUM foliis cordatis, caule volubili, flo- *Convol-*
ribus planiusculis. *Sp. pl.* 364. *Fl. suec.* 344. *vulus.*

Convolvulus minor semine triangulo. *Bauh. pin.* 295.

Convolvulus minor atriplicis folio. *Park.* 171.

Helxine semine triangulo. *Bauh. hist. II.* 157.

Volubilis nigra. *Ger. Em.* 863.

Fegopyrum scandens sylvestre. *R. Syn.* 144.

Anglis, black Bindweed.

Habitat in arvis, hortis, cultis, frequens. ☉. VI.—IX.

OCTANDRIA TETRAGYNIA:

- Fagopyrum.* *II. POLYGONUM foliis cordato-sagittatis, caule erectiusculo inermi, feminum angulis æqualibus.
Sp. pl. 364. *Fl. suec.* 345.
Fagopyrum. *Dod. pempt.* 511. *R. Syn.* 144.
Fagopyrum. *Ger. Em.* 89. *Park.* 1141.
Erysimum Theophrasti folio hederaceo. *Bauh. pin.* 27.
Anglis, Buck wheat, or Brank.
Habitat in arvis, cultis, passim. ☉. VII. VIII.

TETRAGYNIA.

A D O X A. 450.

- Moschatellina.* *Cal. bifidus, inferus.* *Cor.* 4. f. 5. -fida supera. *Bacca* 4 f. 5 -locularis, calyce coalita.
 I. ADOXA. *Sp. pl.* 367. *Fl. suec.* 347.
Moschatellina foliis fumarizæ bulbosæ. *Bauh. hist.* III. 206. *R. Syn.* 267.
Ranunculus nemorosus Moschatella dictus. *Park.* 226.
Ranunculus nemorum moschatellina dictus. *Bauh. pin.* 178.
Radix cava minima viridi flore. *Ger. Em.* 1090.
Anglis, tuberous Moschatel.
Habitat in sylvis et umbrosis. 4. III. IV.

P A R I S. 449.

- quadrifolia.* *Cal. tetraphyllus. Petala quatuor angustiora. Bacca* quadrilocularis.
 I. PARIS. *Sp. pl.* 364. *Fl. suec.* 346.
Herba Paris. *Ger. Em.* 405. *Park.* 390. *Bauh. hist.* III. 316. *Tourn.* 233. *Till. ab.* 33. ic. 32. *R. Syn.* 264.
Solanum quadrifolium bacciferum. *Bauh. pin.* 167.
Anglis, Herb Paris, True-love, or One-berry.
Habitat in sylvis et umbrosis. In sylvula non longe ab *Hampstedio, C. Miller.* 4. V. VI.
Pharmac. *Herba Paris, Fructus, Radix.*

ELATINE. 451.

Cal. tetraphyllus. *Petala* quatuor. *Caps.* quadrilobularis, quadrivalvis, depressa.

1. ELATINE foliis verticillatis. *Roy. lugdb.* 452. *Sp. Alfinastrum.*
pl. 368. *Fl. suec.* 349.

Elatine foliis emerfis linearibus, immerfis capillaceis.

Sauv. monsp. 1743. p. 50.

Alfinastrum gallii folio. *Vaill. paris.* 6. t. 1. f. 6.

Equisetum palustre linariæ scopariæ folio. *Bauh. pin.*

15.

Ericoides facie pinastellæ. *Rupp. jen.* 99.

Anglis, Waterwort.

Habitat in fossis. Found by Mr. J. Sherard on the boggy ground, on the common, just by the road from Eltham to Chiffelhurst. *R. Syn.* 346. 4. VIII.

Classis IX.

ENNEANDRIA
HEXAGYNIA.

BUTOMUS. 455.

*Cal. o. Petata sex. Caps. sex. polyspermæ.**umbellatus.* 1. BUTOMUS. *Cæs.* 553. *R. Syn.* 273. *Fl. lapp.* 159.
suæ. 350. *Sp. pl.* 372.*Juncus floridus.* *Bauh. hist.* II. 524. *Park.* 1197.
major. *Bauh. pin.* 12.*Gladiolus palustris* *Cordi.* *Ger. Em.* 29.*Anglis,* flowering Rush, or Water Gladiole.*Habitat in fossis limosis, et ad ripas fluviorum et stagnorum.* 4. VI. *Near Blackwall.**Radix comeditur ab inferioris Lena
fluvii partis incolis, prope glaciale
oceanum sitis, Yakoutes dictis.*

Classis

*Classis X.*DECANDRIA
MONOGYNIA.

* Flores Polypetali æquales.

MONOTROPA. 477.

Cal. o. Petala decem, horum quinque exteriora basi excavato-mellifera. Caps. quinquevalvis. Quinta pars numeri quibusdam excluditur.

I. MONOTROPA floribus lateralibus octandris, terminali decandro. *Fl. suec. 351. Sp. pl. 387.* *Hypopitys.*

Hypopitys lutea. Cat. giff. 99. app. 134. t. 7. R. Syn. 317.

Orobanche hypopitys. Bauh. pin. 88. pr. 31.

Orobanche verbasculi odore. Plot. ox. 146. t. 9. f. 6. Pluk. ph. t. 209. f. 5.

Orobanche flore brevior dupli, verbasculi odore.

Hist. Ox. III. 504. f. 12. t. 16. f. 20.

Anglis, Bird's-nest smelling like Primrose-roots.

Habitat in sylvis sed rarius. At the roots of trees in the woods near Stoken-church; in fagetis Suffexiæ, Mr. Manningham; prope Tring in Hertfordshire, Mr. Doody. R. Syn. In a wood near Maidstone in Kent, Mr. Jarvis. Blackf. sp. bot. 39. 4. VII.

PYROLA. 490.

Cal. quinquepartitus. Petala quinque. Caps. quinquelocularis, angulis dehiscens.

I. PYROLA staminibus adscendentibus, pistillis declinatis. *Fl. suec. 360. Sp. pl. 396. Mat. med. 212. lia.* *rotundifolia.*

Pyrola foliis subrotundis, scapo racemoso. Hort. cliff.

162.

Pyrola.

Pyrola. *Ger. Em.* 408. *Bauh. hist.* III. 535. *R. Syn.* 363.

Pyrola nostras vulgaris. *Park.* 508.

Pyrola rotundifolia major. *Bauh. pin.* 191.

Anglis, common Winter-green.

Habitat in sylvis et nemoribus. In Septentrionalibus Angliæ locis reperitur; in Meridionalibus rarius occurrit; near Halifax plentifully, and in Stokenchurch woods, Oxfordshire. *R. Syn.* 4. VI. VII.

Pharmac. Pyrolæ Herba.

minor. 2. PYROLA floribus racemosis dispersis, staminibus pistillisque rectis. *It. gotl.* 4. *Fl. suec.* 361. *Sp. pl.* 369.

Pyrola tuba recta minor, folio frequentius serrato, spica brevior et densior. *Hall. helv.* 7.

Pyrola minor. *Riv. pent.* 149. *R. Syn.* 363.

Pyrola folio minore et durior. *Bauh. pin.* 191.

Anglis, lesser Winter-green.

Habitat in sylvis. In sylvula inter Clapham et Ingleton, in comitatu Eboracensi. 4. VIII.

secunda. 3. PYROLA racemo unilaterali. *Fl. suec.* 362. *Sp. pl.* 396.

Pyrola floribus uno versu sparsis. *Fl. lapp.* 168.

Pyrola folio mucronato serrato. *Bauh. pin.* 181. *R. Syn.* 363.

Pyrola tenerior. *Park.* 509.

Pyrola secunda tenerior. *Clusii hist.* 117. *Ger. Em.* 408.

Anglis, dented-leaved Winter-green.

Habitat in sylvis et nemorosis. In Haslewood, near Sir Walter Vavasor's park in Yorkshire. *R. Syn.* 4. VI.

** Flores monopetali æquales.

A N D R O M E D A. 485.

Cal. quinquepartitus. *Cor.* ovata: ore quinquefido.

Capf. quinquelocularis.

polifolia. 1. ANDROMEDA pedunculis aggregatis, corollis ovatis,

tis, foliis alternis lanceolatis revolutis. *Fl. suec.*

353. *Sp. pl.* 393.

Ledum palustre nostras, arbuti flore. R. Syn. 472.

Cistus Ledon palustris Rosmarini folio, Arbuti flore.

R. Syn. Ed. I. 142. *t.* I.

Rosmarinum sylvestre minus nostras. Park. 76.

Erica humilis Rosmarini foliis, unedonis flore, capsula cistoide. Pluk. ph. t. 175. *f.* I.

Anglis, marsh Cistus, or wild Rosemary.

Habitat in paludibus turfosis in comitatibus Eboracensi, Lancastriensi, Westmorlandico, Cumberlandico, frequens. 4. III. IV.

A R B U T U S. 488.

Cal. quinquepartitus. Cor. ovata : ore basi pellucida.

Bacca quinquelocularis.

1. ARBUTUS caule erecto, foliis glabris ferratis, bac- *Unedo.*
cis polyspermis. *Sp. pl.* 395. *Roy. lugdb.* 440.
Sauv. monsp. 56.

Arbutus. Ger. Em. 1496. *Park.* 1489. *Cam. epit.*
168. *R. Syn.* 464.

Arbutus folio ferrato. Bauh. pin. 460.

Anglis, common Strawberry-tree.

Habitat in occidentali Hiberniæ parte. R. Syn. 2.

2. ARBUTUS caulis procumbentibus, foliis rugosis fer- *alpina.*
ratis. *Fl. lapp.* 161. *suec.* 359. *Hort. cliff.* 163.
Roy. lugdb. 440. *Hall. helv.* 415. *Sp. pl.* 395.

Vaccinia rubra, foliis myrtinis crispis. D. Merret.
R. Syn. 457.

Vitis idæa foliis oblongis albertibus. Bauh. pin. 470.
Tourn. 608.

Vitis idæa repens, fructu racemoso magno nigerrimo.
Rudb. it. 9. lapp. 100.

Anglis, mountain Strawberry-tree.

*Habitat in montibus siccis. Four miles from Hepten-
stal, near Widdop, on a great stone by the river Gort
in Lancashire; found also by Mr. Lhwyd plentifully
for some miles together, in that end of Mull next to
Eycolumb-Kill, Scotland. R. Syn.* 2.

D I-

DIGYNIA.

CHRYSO SPLENIUM. 493.

Cal. quadrifidus f. quinquefidus, coloratas. *Cor.* o.

Capsf. birostris, unilocularis, polysperma.

oppositifolium. 1. CHRYSO SPLENIUM foliis oppositis. *Sauv. monsp.* 128. *Sp. pl.* 398.

Chryso splenium f. saxifraga aurea. *Faborn. hist.* 1224.

Chryso splenium foliis amplioribus auriculatis. *Tourn.* 146.

Saxifraga aurea. *Ger. Em.* 841. *Park.* 425. *R. Syn.* 158.

Saxifraga rotundifolia aurea. *Baub. pin.* 309.

Anglis, common golden Saxifrage.

Habitat ad scaturigines in sylvis umbrosis. In Charlton wood. *Blackf. spec. bot.* In ericeto Hampstediano prope Londinum. 4. IV.

alternifolium. 2. CHRYSO SPLENIUM foliis alternis. *Fl. suec.* 365. *Sp. pl.* 398.

Chryso splenium foliis pediculis oblongis insidentibus. *Tourn.* 146.

Saxifraga aurea foliis pediculis oblongis insidentibus. *R. Syn.* 158.

Sedum palustre luteum majus, foliis pediculis longis insidentibus. *Hist. Ox. III.* 477. f. 12. t. 8. f. 8.

Anglis, alternate-leaved golden Saxifrage.

Habitat cum priore sed rarius. Prope Bingley in comitatu Eboracensi copiose. 4. IV.

SAXIFRAGA. 494.

Cal. quinquepartitus. *Cor.* pentapetala. *Capsf.* birostris unilocularis.

* *Foliis indivisis, caule subnudo.*

stellaris. 1. SAXIFRAGA foliis ferratis, caule nudo ramofo, petalis lanceolatis. *Fl. suec.* 367. *Sp. pl.* 400.

Saxifraga caule nudo simplici, foliis lanceolatis dentatis petalis acutis, *Fl. lapp.* 175.

Geum

Geum palustre minus, foliis oblongis crenatis. *Tourn.* 252. *R. Syn.* 354.

Cotyledon aquatica hirsuta. *R. hist.* 1046.

Sanicula myosotis alpina, floribus albicantibus fere umbellatis. *Pluk. ph. t.* 58. *f.* 2. *et t.* 222. *f.* 4.

Sedum montanum hirsutum, mucronato et dentato folio, flore albo guttato. *Hist. Ox. III.* 478. *f.* 12. *t.* 9. *f.* 13.

Anglis, hairy Kidney-wort.

Habitat in petris et rupibus aqua humentibus, et ad rivulos. On the mountains of Wales, as Snowdon, Carnedd-Llewellyn, Caderidris, &c. Also on Hardknot and Wrenose, by Buckbarrow well in Longfledale, Westmoreland. *R. Syn.* 4. VI. VII.

2. *SAXIFRAGA* foliis obovatis crenatis subseffilibus, *nivalis*. caule nudo, floribus congestis. *Fl. suec.* 368. *Sp. pl.* 401.

Saxifraga caule nudo simplici, foliis elliptico-subrotundis crenatis, floribus capitatis. *Fl. lapp.* 176. *t.* 2. *f.* 5, 6.

Saxifraga foliis oblongo-rotundis dentatis, floribus compactis. *R. Syn.* 354. *t.* 16. *f.* 1.

Anglis, mountain Sengreen.

Habitat in summis alpibus arvonie, e rupibus Clogwyn y Garnedd Crib y Distilh, Glogwyn du ymhen y Glyder, &c. juxta Lhan-berys. *D. Lhwyd. R. Syn.* 4. VI.

** *Foliis indivisis, Caule folioso.*

3. *SAXIFRAGA* foliis ovatis oppositis imbricatis: sum- *oppisitifolia.* mis ciliatis. *Fl. suec.* 369. *Sp. pl.* 402.

Saxifraga foliis ovatis quadrangulo-imbricatis, ramis procumbentibus. *Fl. lapp.* 179.

Saxifraga alpina ericoides, flore cæruleo. *Tourn.* 253. *R. Syn.* 353.

Sedum alpinum ericoides cæruleum. *Bauh. pin.* 284. *pr.* 132. *hist. III.* 694. *Hist. Ox. III.* 480. *f.* 12. *t.* 10. *f.* 36.

Anglis, mountain heath-like Sengreen.

Habitat

Habitat in rupibus et petris. In rupibus montis Ingleborough non ad summitatum, sed ad latus occidentale, D. Richardson. On the high rocks of Snowdon, plentifully, D. Lhwyd. R. Syn. 4. III. IV.

aizoides. 4. SAXIFRAGA foliis caulinis lineari-subulatis sparsis nudis inermibus, caulibus decumbentibus. *Fl. suec.* 371. *Sp. pl.* 403.

Saxifraga alpina angusto folio, flore luteo guttato. R. Syn. 353.

Sedum minus alpinum luteum nostras. R. Syn. ed. II. 212.

Sedum alpinum flore pallido. *Baub. pin.* 284. *Hist. Ox.* III. 477. *J.* 12. *t.* 6. *f.* 3.

Sedum minus IV. seu alpinum I. *Clus. hist.* III. 60. *Anglis*, yellow mountain Sengreen.

Habitat in montibus. On the sides of Ingleborough hill plentifully, and in the like places in Westmoreland. R. Syn. 4. VIII.

autumnalis. 5. SAXIFRAGA foliis caulinis linearibus alternis ciliatis: radicalibus aggregatis. *Sp. pl.* 402.

Saxifraga foliorum margine ciliari, floribus luteis maculosis. *Hall. helv.* 399.

Saxifraga angustifolia autumnalis, flore luteo guttato, foliis florum magis acuminatis. *Breyn. cent.* 106. *t.* 48.

Geum angustifolium autumnale, flore luteo guttato. *Tourn.* 252. R. Syn. 355.

Anglis, autumnal Sengreen.

Habitat in palustribus turfosis. Found by Dr. Kingstone on Knotsford-moor, Cheshire. R. Syn. 4. VII. VIII.

*** *Foliis lobatis, caule erecto.*

cæspitosa. 6. SAXIFRAGA foliis radicatis aggregatis linearibus integris trifidisque, caule erecto subnudo subbifloro. *Sp. pl.* 404.

Saxifraga foliis simplicibus et trifidis, caule pæne aphylo paucifloro. *Hall. helv.* 405.

Saxifraga foliis partim integris partim trifidis. *Segu. ver.* 451. *t.* 9. *f.* 4.

Sedum

Sedum tridactylites alpinum minus. *Bauh. pin.* 284.
pr. 131.

Anglis, small mountain Sengreen.

Habitat in montibus supra Ambleside in comitatu Westmorlandico. 24. VIII.

7. *SAXIFRAGA* foliis cuneiformibus trifidis alternis, *trydactyli-*
caule erecto ramofo. *Fl. suec.* 375. *Sp. pl.* 404. *tes.*

Saxifraga foliis trifidis basi angustis caule erecto. *Roy.*
lugdb. 457. *Sauv. monsp.* 208.

Saxifraga verna annua humilior. *Tourn.* 252. *R. Syn.*
354.

Sedum tridactylites tectorum. *Bauh. pin.* 285.

Paronychia rutaceo folio. *Ger. Em.* 624.

Paronychia foliis incis. *Park.* 556.

Anglis, Rue-leaved Sengreen, or Whitlow-grass.

Habitat in muris, tectis, ruderatis passim. ①. IV.

Pharmacop. *Paronychiæ Herba.*

8. *SAXIFRAGA* foliis reniformibus lobatis, caule ra- *granulata.*
mofo, radice granulosa. *Hort. cliff.* 167. *Mat.*
med. 215. *Fl. suec.* 372. *Sp. pl.* 403.

Saxifraga rotundifolia alba. *Bauh. pin.* 309. *R.*
Syn. 354.

Saxifraga alba. *Ger. Em.* 841. *vulgaris.* *Park.* 424.

Saxifraga alba radice granulosa. *Bauh. hist.* III. 706.

Anglis, white Sengreen, or Saxifrage.

Habitat in pratis et pascuis siccioribus. Prope Wandsworth copiose. 24. V.

Pharmac. *Saxifraga albæ Radix, Foliæ, Flores, Semina.*

**** *Foliis lobatis, caulibus procumbentibus.*

9. *SAXIFRAGA* foliis caulinis linearibus integris tri- *hypnoides*
fidisve, stolonibus procumbentibus, caule erecto
nudiusculo. *Sp. pl.* 405.

Saxifraga foliis omnibus angustis petiolatis apice tri-
dentatis, caule folioso ramofo viscido. *Hall. helv.*
404.

Saxifraga muscosa trifido folio. *Tourn.* 252. *R. Syn.*
354.

Sedum alpinum trifido folio. *Bauh. pin.* 284.

Sedum

Sedum alpinum laciniatis Ajugæ foliis. Park. 739.
Anglis, trifid Sengreen.

Habitat in montibus Westmorlandicis, Eboracensibus,
Derbiansibus, Cambrobritannicis. passim. 4. IV. V.

SCLERANTHUS. 497.

Cal. monophyllus. Cor. o. Sem. duo calyce inclusa.

annuus. 1. SCLERANTHUS calycibus fructus patulis. Fl.
succ. 377. Sp. pl. 406.

Knawel. Trag. 393. R. Syn. 159.

Polygonum germanicum vel Knawel Germanorum.
Park. 747.

Polygonum selinoides five Knawel. Ger. 453.

Polygonum gramineo folio majus erectum. Bauh.
pin. 281.

Anglis, German Knot-grass, or annual Knawel.

Habitat in arvis et locis arenosis passim. ☉. VIII.

perennis. 2. SCLERANTHUS calycibus fructus clausis. Fl.
succ. 378. Sp. pl. 406.

Knawel incanum flore majore perenne. R. hist. 213.
Syn. 160. t. 5. f. 1.

Alchimilla gramineo folio majori flore. Tourn. 509.
Vaill. parif. 4. t. 1. f. 5.

Anglis, perennial Knawel.

Habitat in arvis glareosis. In the sandy grounds and
balks of corn fields about Elden in Suffolk, plenti-
fully. R. Syn. 4. VIII.

SAPONARIA. 499.

Cal. monophyllus nudus. Petala quinque unguicu-
lata. Caps. oblonga unilocularis.

officinalis. 1. SAPONARIA calycibus cylindricis, foliis ovato-
lanceolatis. Hort. cliff. 165. upf. 106. Mat. med.
214. Roy. lugdb. 444. Grøn. virg. 106. Gort.
gehr. 245. Sp. pl. 408.

Saponaria. Ger. Em. 444. vulgaris. Park. 641.
Bauh. hist. III. 346.

Saponaria major lævis. Bauh. pin. 206.

Lychnis Saponaria dicta. R. Syn. 339.

β *Lychnis*

DECANDRIA DIGYNIA.

161

- β *Lychnis Saponaria dicta folio convoluto. R. Syn. 339.*
Saponaria concava Anglica. Baub. pin. 206.
Anglis, Sopewort.
Habitat ad sepes, in Ericeto nigro prope Morden Col-
ledge in Cantio. 4. VII. VIII.
Pharmac. Saponaria. Folia Radix.

DIANTHUS, 500.

Cal. cylindricus monophyllus: basi squamis quatuor.
Petala quinque unguiculata. Caps. cylindrica uni-
locularis.

* *Flores aggregati.*

1. **DIANTHUS** floribus aggregatis fasciculatis, squamis calycinis lanceolatis villosis tubum æquantibus. *Armeria.*
Syst. Nat. 1029. Sp. pl. 410.

Caryophyllus latifolius barbatus minor annuus, flore minore. R. Syn. 337.

Caryophyllus pratensis. Ger. Em. 594. noster major et minor. Park. 1338.

Caryophyllus barbatus sylvestris. Baub. pin. 209.

Armeria sylvestris altera. Lob. ic. 448.

Anglis, Deptford Pink.

Habitat in pratis et pascuis glareosis in Sylva Charleton Wood dicta, in Cantio; prope Croydon copiose.
 ○. VII.

2. **DIANTHUS** floribus aggregatis capitatis squamis calycinis ovatis obtusis muticis tubum æquantibus. *prolifer.*
Sp. pl. 410.

Caryophyllus sylvestris prolifer. Baub. pin. 209. Park. 1338. R. Syn. 337.

Viscaria. Ger. Em. 601.

Anglis, Limewort.

Habitat in glareosis sed rarius. Found by the Rev. Mr. Manningham, in Selsey Island, Suffex. R. Syn.
 ○. VIII.

** *Flores solitarii, plures in eodem caule.*

3. **DIANTHUS** floribus solitariis squamis calycinis, lanceolatis binis, corollis crenatis. *deltoides.*
Fl. suec. 382. Sp. pl. 411.

M

Cary-

DECANDRIA DIGYNIA.

Caryophyllus minor repens nostras. R. Syn. 335. Dill. *elth.* 402.

Caryophyllus sylvestris vulgaris latifolius. Bauh. *pin.* 209.

Anglis, Maiden Pinks.

Habitat in ericetis pratis et pascuis frequens in ericeto Hamstediano prope Londinum. 4. VI. VII.

glaucus. 4 DIANTHUS floribus solitariis, squamis calycinis lanceolatis brevibus corollis crenatis. *Sp. pl.* 411. *Hort. upf.* 104.

Tunica ramosior, flore candido cum corolla purpurea. Dill. *elth.* 400. t. 298. f. 348.

Anglis, Mountain Pink.

Habitat in montosis. On Chedder-rocks, in Somersetshire. Mr. Brewer. 4. VII.

*** *Caule uniflora herbaceo.*

arenarius. 5. DIANTHUS caulibus subunifloris, squamis calycinis ovatis, obtusis, petalis multifidis, foliis linearibus. *Sp. pl.* 412. *Fl. suec.* 384.

Dianthus foliis brevibus, squamis muticis caule unifloro. *Sauv. meth.* 143.

Armeria species flore in summo caule singulari. R. *Syn.* 336.

Armeria sylvestris humilis flore unico. Bauh. *pin.* 209. rubello. *Merr. pin.*

Caryophyllus sylvestris, *Alfines holosteæ arvensis glabræ* foliis flore unico, calyce barbato. *Pluk. ph.* t. 8 p. f. 3.

Anglis, Stone Pink.

Habitat in pratis siccissimis et muris sed rarius. On Chedder-rocks. Mr. Brewer. In the North of England. Mr. Doody. R. *Syn.* 4. VII.

TRIGYNIA.

CUCUBALUS. 502.

Calyx. inflatus. *Petala* quinque unguiculata absque corona ad faucem. *Caps.* trilocularis.

1. CU-

1. CUCUBALUS calycibus campanulatis, petalis dibacciferus. stantibus, pericarpis coloratis, ramis divaricatis. *Sp. pl.* 414.
Cucubalus Plinii. *Lugd.* 1429. *Tourn.* 339. *R. Syn.* 267.
Alpine scandens baccifera. *Bauh. pin.* 250.
Anglis, berry-bearing Chickweed.
Habitat in sepibus in Insula Mona. D. Richardson.
R. Syn. 4. VII.
2. CUCUBALUS calycibus subglobosis glabris reticulato-venosis capsulis trilocularibus, corollis subaudis. *Fl. suec.* 385. *Sp. pl.* 414.
Lychnis sylvestris quæ Benalbum vulgo. *Bauh. pin.* 205. *R. Syn.* 337.
Behen album officinarum. *Ger. Em.* 678.
Papaver spumeum sive Benalbum vulgo. *Park.* 263.
Anglis, bladder Campion, or white corn Campion.
Habitat in arvis pratis et pascuis siccioribus frequens.
4. VII.
3. CUCUBALUS floribus lateralibus undique decumbentibus, caule indiviso, foliis basi reflexis. *Sp. pl.* 414. *Fl. suec.* 386.
Lychnis major noctiflora Dubrensis perennis. *R. hist.* 995. *Syn.* 340.
Lychnis montana noctiflora hirsuta latifolia alba, floribus absque appendicibus. *Till. pis.* 105.
Anglis, Dover Campion.
Habitat in montosis. Found on Dover Cliffs, by Mr. Newton. *R. Syn.* 4. VII.
4. CUCUBALUS floribus dioicis, petalis fetaceis indivisis. *Hort. cliff.* 272. *Roy. lugdb.* 445. *Sp. pl.* 415.
Lychnis viscosa flore muscoso. *Bauh. pin.* 266.
Sesamoides Salamanticum magnum. *Ger. Em.* 493.
Muscipula Salamantica major. *Park.* 636.
Anglis, Spanish Campion, or Catchfly.
Habitat in pascuis glareosis. In and about the gravel-pits on the north side of Newmarket town: also by the way-sides all along from Barton Mills to Thetford in Norfolk. *R. Syn.* 4. VII. VIII.

DECANDRIA TRIGYNIA.

acaulis. 5. CUCUBALUS *acaulis.* *Fl. lapp.* 104. *fuec.* 337.
Sp. pl. 415.

Lychnis alpina minima. *R. hist.* 1004. *Syn.* 341.
Caryophylleus IX Clusii caryophyllus pumilio alpi-
nus. *Ger. Em.* 593.

Lychnis alpina pumila folio gramineo f. muscus alpi-
nus Lychnidis flore. *Bauh. pin.* 206.

Anglis, mos Campion.

Habitat in montibus Arvonie, Devonie, frequens. 4.
VII.

S I L E N E. 503.

Cal. ventricosus. Petala quinque unguiculata: coro-
nata ad faucem. Caps. trilocularis.

* *Floribus solitariis lateralibus.*

anglica. 1. SILENE *hirsuta petalis emarginatis, floribus erec-*
tis, fructibus reflexis pedunculatis alternis. *Syst.*
Nat. 1031. *Sp. pl.* 416.

Viscago cerasii foliis, vasculis pendulis anglica. *Dill.*
elth. 417. *t.* 309. *f.* 398.

Lychnis sylvestris flore albo minimo. *R. hist.* 996.
Syn. 339.

Anglis, small corn Catchfly or Campion.

Habitat in arvis. In the corn fields near Colchester,
Essex. Mr. Newton. Blackf. Sp. Bot. among the
corn near the Devil's-ditch in Cambridgeshire. Mr.
Dent. R. Syn. ☉. VI. VII.

amoena. 2. SILENE *petalis bifidis: coronula sub-coadunata,*
floribus secundis, pedunculis oppositis trifloris, ra-
mis alternis. *Syst. Nat.* 1031. *Sp. pl.* 417.

Lychnis marina anglica. *Ger. Em.* 470. *R. Syn.* 337.

Lychnis marina repens alba. *Park.* 638.

β *Lychnis maritima anglica amplioribus foliis.* *D.*
Brewer. Rand. hort. chel.

Anglis, Sea Campion.

Habitat in littoribus maritimis frequens. 4. VIII.

3. SILENE

** *Floribus lateralibus confertis.*

3. **SILENE** petalis bifidis, floribus lateralibus secundis *nutans.*
cernuis caule incurvato. *Fl. suec.* 388. *Sp. pl.* 417.
Silene foliis lanceolatis, caule paniculato, floribus nu-
tantibus calyce striato corolla involuta. *Hort. cliff.*
171.
Lychnis sylvestris alba IX. clusii. *Ger. Em.* 470. *R.*
Syn. 340.
Lychnis montana viscosa alba latifolia. *Baub. pin.*
205.
Lychnis sylvestris alba f. *ocymoides minus album.*
Park. 631.
Polæmonium petræum Gesneri. *Baub. hist.* III. 351.
Anglis, Nottingham Catchfly.
Habitat in pratis montosis. On the walls of Notting-
ham-castle, and thereabout. *R. Syn.* 4. VI. VII.

*** *Floribus ex dichotomia caulis.*

4. **SILENE** calycibus fructus conicis, striis triginta, fo- *conoides.*
liis mollibus. *Hort. upf.* 110. *Sp. pl.* 418.
Lychnis sylvestris angustifolia caliculis turgidis striatis.
Baub. pin. 205. *R. Syn.* 341.
Lychnis sylvestris altera incana caliculis striatis. *Lob.*
ic. 338.
Anglis, greater corn Catchfly, or Champion.
Habitat in arvis glareosis. A little to the north of
Sandown-castle, plentifully. *R. Syn.*
5. **SILENE** calycibus decemangularibus: dentibus tu- *noctiflora.*
bum æquantibus, caule dichotomo. *Sp. pl.* 419.
Fl. suec. 389.
Lychnis noctiflora. *Baub. pin.* 205. *Park.* 632. *R.*
Syn. 340.
Anglis, night-flowering Catchfly.
Habitat in arvis. Found by Mr. Dent among corn, on
the left hand of the road leading from Newmarket
to Canvass-hall, in Wood-ditton. *R. Syn.* 4. VII.

6. **SILENE** floribus fasciculatis fastigiatis, foliis superi- *Armeria.*
oribus cordatis glabris. *Hort. upf.* 110. *Sp. pl.* 420.

DECANDRIA TRIGYNIA.

Lychnis viscosa purpurea latifolia lævis. *Bauh. pin.*
205 *R. Syn* 341.

Lychnis sylvestris. *I. Clus. hist.* 288.

Armerius flos quartus. *Dod pempt.* 176.

Anglis, broad-leaved Catchfly.

Habitat on the banks of the river half a mile below
Chester, Dr. Richardson. *R. Syn.* ☉. VIII.

S T E L L A R I A. 504.

Cal. pentaphyllus, patens. *Petala* quinque bipartita.
Capsf. unilocularis, polysperma.

Holostea. 1. *STELLARIA* foliis lanceolatis ferrulatis, petalis bifidis. *Fl. suec.* 392. *Sp. pl.* 422.

Caryophyllus holosteus arvensis glaber flore majore.
Bauh. pin. 210. *R. Syn.* 346.

Gramen leucanthemum. *Ger. Em.* 47. *Park.* 1325.

Anglis, greater Stichwort.

Habitat in sylvis et sepibus passim. 4. IV. V.

graminea. 2. *STELLARIA* foliis linearibus integerrimis, floribus paniculatis. *Fl. suec.* 393. *Sp. pl.* 422.

Caryophyllus holosteus arvensis glaber, flore minore.
Bauh. pin. 210. *R. Syn.* 346.

Gramen leucanthemum alterum. *Ger. Em.* 47.

Gramen leucanthemum minus. *Park.* 1325.

β *Caryophyllus holosteus arvensis medius.* *R. Syn.* 347.

γ *Alfina longifolia uliginosis proveniens locis.* *Bauh. hist.*
III. 365. *R. Syn.* 347.

Alfina aquatica media. *Bauh. pin.* 251.

Alfina fontana. *Tab. ic.* 712. *Ger. Em.* 613.

Anglis, lesser Stichwort.

Habitat in pratis pascuis, sepibus frequens. γ. in paludosis et pratis humidioribus passim. 4. VII.

memorum. 3. *STELLARIA* foliis cordatis petiolatis, paniculae pedunculis ramosis. *Sp. pl.* 421. *Fl. suec.* 291.

Alfina montana folio Smilacis instar, flore laciniato.

Hist. Ox. II. 550. f. 5. t. 23. f. 2. *R. Syn.* 347.

Alfina montana latifolia, flore laciniato. *Bauh. pin.*
251.

Alfina

Alfne hederacea montana maxima. Col. Ecphr. I. 290. t. 290. Park 761.

Anglis, broad-leaved Stichwort.

Habitat in sylvis et sepibus, et ad ripas fluviorum in comitatibus Eboracensi, Lancastriensi, Westmoriandico, Cumberlandico, passim. 4. VII. VIII.

A R E N A R I A. 505.

Cal. pentaphyllus, patens. Petala quinque integra.

Caps. unilocularis polyspermæ.

1. ARENARIA foliis ovatis acutis carnosis. *Fl. lapp. peploides. 149. suec. 396. Sp. pl. 423.*

Alfne marina foliis portulacæ. Fl. pruss. 12. t. 2. R. Syn. 351.

Alfne littoralis foliis portulacæ. Baub. pin. 251.

Anthyllis maritima lentifolia. Baub. pin. 282. Park. 282.

- Anthyllis lentifolia seu alfne cruciata marina. Ger. Em. 622.*

Anglis, Sea Chickweed.

Habitat in littoribus maritimis frequens. 4. VI. VII.

2. ARENARIA foliis ovatis acutis petiolatis nervosis. *trinervia. Fl. suec. 397. Sp. pl. 423.*

Alfne plantaginis folio. Baub. hist. III. 364. R. Syn. 349.

Anglis, Plantain-leaved Chickweed.

Habitat in sylvis et sepibus humidis frequens. 5. V. VI.

3. ARENARIA foliis subovatis acutis sessilibus, corollis *serpyllifolia. calyce brevioribus. Fl. suec. 398. Sp. pl. 423.*

Alfne minor multicaulis. Baub. pin. 250. R. Syn. 349.

Alfne minima. Ger. Em. 488. Baub. hist. III. 364.

Alfne aquatica minima. Park. 1259.

Anglis, the least Chickweed.

Habitat in tectis, muris, arenosis, aridis passim. 5. V. VI.

- saxatilis*. 4. ARENARIA foliis subulatis, caule paniculatis, calycem foliolis ovatis. *Sp. pl.* 424.
 Spergula foliis aciformibus densis, ramis ramosis subnudis. *Sauv. monsp.* 45.
 Alfine pusilla pulchro flore folio tenuissimo nostras f. faxifraga pusilla caryophylloides, flore albo pulchello. *R. Syn.* 351.
 Alfine alpina glabra tenuissimis foliis, floribus albis. *Herm. par.* 12. t. 12
 Alfine caryophylloides tenuifolia, flore albo punctato. *Pluk. ph. t.* 7. f. 3.
 Alfine saxatilis laricis folio minor et minori flore. *Tourn. magn. mosq.* 11. t. 11.
Angli., mountain Chickweed.
 Habitat in montosis Eboracensibus circa Settle, ut et Derbiansibus copiose. On the side of a watery bank, at about three miles from the Land's end. *R. Syn. in montosis* Westmorlandicis, circa Kendal copiose. 4. VIII.
- laricifolia*. 5. ARENARIA foliis setaceis, caule superne nudiusculo, calycibus subhirsutis. *Sp. pl.* 424.
 Alfine foliis fasciculatis, petiolis simplicibus, calyce hirsuto. *Hall herb.* 388.
 Alfine alpina junceo folio. *Bauh. pin.* 251. pr. 118. *R. Syll.* 60. *hist.* 1031.
 Alfine caryophylloides tenuifolia montana lini flore. *R. hist.* 1027.
 Auricula muris pulchro flore, folio tenuissimo. *Bauh. hist.* III. 360.
Anglis, Larch-leaved Chickweed.
 Habitat in montibus Westmorlandicis passim. 4. VIII.
- tenuifolia*. 6. ARENARIA foliis subulatis caule paniculato, pedunculis simplicibus, capsulis erectis petalis calyce brevioribus. *Sp. pl.* 424.
 Alfine tenuifolia. *Bauh. hist.* III. 364. *R. Syn.* 350. *Vaill. parif.* 7. t. 3. f. 1. *sequ. ver.* 418. t. 6. f. 2.
Anglis, fine-leaved Chickweed.
 Habitat in pratis siccis glareosis, sed rarius. In the corn-fields on the borders of Friplow-heath, Gogmagog

gog hills, and other places in Cambridgeshire. R. Syn. 350. In Cornbury quarry near Charlbury, Oxfordshire; Blackf. spec. bot. supra mure in oppido Battersea prope Londinum 24. VI. VII.

7. ARENARIA foliis linearibus stipulis membranaceis *rubra.*
vaginantibus. *Sp. pl. 423. Fl. suec. 399.*
Spergula purpurea. *Bauh. hist. III. 722. R. Syn. 351.*
Saginæ Spergula minima. *Park. 561.*
Alfine, Spergulæ facie minor f. Spergula minor sub-
cæruleo flore. *Bauh. pin. 251. pr. 119.*
- β Arenaria foliis linearibus longitudine internodiorum. *marina.*
Hort. cliff. 173. Gron. virg. 161. Sp. pl. 423.
Alfine Spergulæ facie media. *Bauh. pin. 251. R. Syn. 351.*
Spergula marina nostras. *Bauh. hist. III. 778.*
- γ Spergula maritima flore parvo cæruleo, semine vario.
C. giff. 30.
Anglis, purple-flowered Chickweed, or Spurrey. β Sea Spurrey.
Habitat in pratis et arvis glareosis, β in littoribus mari-
nis frequens. ☉. VI. VII.

P E N T A G Y N I A.

C O T Y L E D O N. 512.

- Cal. quinquefidus. Cor. monopetala, Squamæ necta-
riferæ quinque ad basin germinis Caps. quinque.*
1. COTYLEDON foliis cuculatis serrato-dentatis al- *umbilicus.*
ternis, caule ramoso floribus erectis. *Sp. pl. 429. ♀.*
Cotyledon foliis peltatis. *Vir. cliff. 39. Roy. lugdb. 454.*
Cotyledon vera radice tuberosa. *Bauh. hist. III. 683.*
R. Syn. 271.
Cotyledon major. *Bauh. pin. 289.*
Umbilicus veneris. *Ger. Em. 529. vulgaris. Park. 740.*
Anglis, Navel-wort, Kidney-wort, or Wall Penny-wort.

Habitat

Habitat in saxis et muris antiquis sed rarius; prope
Bristolium copiose; circa Troutbeck in comitatu
Westmorlandico. 4. V.—VII.

S E D U M. 513.

Cal. quinquefidus. *Cor.* pentapetala. Squamæ nec-
tariferæ quinque ad basin germanis. *Capf.* quin-
que.

* *Planifolia.*

- Telephium.* 1. SEDUM foliis planiusculis ferratis, corymbo folioso,
caule erecto. *Sp. pl.* 430.
Sedum foliis planiusculis patentibus ferratis, corymbo-
terminali. *Hort. cliff.* 176. *Roy. lugdb.* 456.
Telephium vulgare. *Baub.* pin. 287.
Telephium f. *Crassula* major vulgaris. *Park.* 726.
Crassula f. faba inversa. *Ger. Em.* 519.
Anacampteros, vulgo Faba crassa. *Baub. hist.* III.
681. *R. Syn.* 269.
β Telephium purpureum majus. *Baub.* pin. 287.
γ Telephium album. *Fuchf. hist.* 800.
Anglis, Orpine, or Live-long.
Habitat in tectis, muris, pascuis, et ad sepēs. 4. VIII.
Pharmac. Telephii *Herba.*

** *Teretifolia.*

- reflexum. 2. SEDUM foliis subulatis sparsis, basi membranacea
soluta: inferioribus recurvatis, floribus cymosis.
Fl. succ. 1296.
Sedum minus luteum ramulis reflexis. *Baub.* pin. 283.
R. Syn. 270.
Aizoon scorpioides. *Ger. Em.* 513.
Vermicularis scorpioides. *Park.* 733.
β Sedum minus hæmatoides. *Ger. Em.* 512. *R. Syn.*
269.
Sedum minus luteum folio acuto. *Baub.* pin. 283.
Anglis, yellow Stonecrop, or Prick-madam.
Habitat in muris, tectis, passim.
rupestre. 3. SEDUM foliis subulatis quinquefariam confertis basi
membranacea soluta, floribus cymosis. *Sp. pl.* 431.
Fl. succ. 401. Sedum

Sedum rupestre repens foliis compressis. *Dill. elth.*

343. t. 256. f. 333.

Sedum minus e rupe S. Vincentii. *R. Syn.* 271.

Sedum minus Vincentii. *Pet. h. br. t.* 41. f. 8.

Anglis, St. Vincent's Rock Stonecrop.

Habitat in rupibus St. Vincentii prope Bristolium : in
rupibus Hisvæ valliculæ Nantphracon imminentibus.

R. Syn. 24. VIII.

4. *SEDUM* foliis oblongis obtusis teretiusculis sessilibus *album*,
patentibus, cyma ramosa. *Fl. suec.* 402. *Sp. pl.*
432.

Sedum minus teretifolium album. *Bauh. pin.* 283.
R. Syn. 271.

Sedum minus officinarum. *Ger. Em.* 512.

Vermicularis flore albo. *Park.* 733.

Vermicularis f. *Crassula minor vulgaris.* *Park.* 734.

Anglis, white flowered Stonecrop.

Habitat in muris, tectis, passim. On a wall just beyond
the chapel at Kentish-town. *Blackf. spec. bot.* 24.

VI. VII.

Pharmac. *Vermicularis Herba.*

5. *SEDUM* foliis subovatis adnato-sessilibus gibbis erectiusculis alternis, cyma trifida. *Fl. suec.* 403. *Sp. pl.* 432. *Mat. med.* 218.

Sedum parvum acre flore luteo. *Bauh. hist.* III. 694.
R. Syn. 270.

Sempervivum minus vermiculatum acre. *Bauh. pin.*
283.

Vermicularis seu *Illecebra minor acris.* *Ger. Em.* 517.

Illecebra minor f. *Sedum tertium Dioscoridis.* *Park.*
735.

Anglis, Wall Stonecrop, or Pepper.

Habitat in muris, tectis, montibus, rupibus, frequens.
24. VI.

Pharmac. *Sedi minoris Herba.*

6. *SEDUM* foliis subovatis adnato-sessilibus gibbis erectiusculis sexfariam imbricatis, *Fl. suec.* 405. *Sp. pl.* *sexangulare.*
432.

Semper-

Sempervivum minus vermiculatum insipidum. Bauh. pin. 284.

Sempervivum minimum. Cam. epit. 856.

Anglis, insipid Stonecrop.

Habitat in pratis siccis. Prope Sheerness in insula Shepey copiose. §. VII.

annuum. 7. *SEDUM* caule erecto solitario annuo, foliis ovatis sessilibus gibbis alternis, cyma recurva. *Fl. suec. 405. Sp. pl. 432.*

Sedum minimum non acre flore albo. R. Syn. 270.

Sedum minimum, flore mixto ex albo et rubro, Merr. pin.

Anglis, Mountain Stonecrop.

Habitat in sterilioribus Suffolciæ, itinere a Yarmouth ad Donewich plurimum observavimus: item in rupibus Lancastriensibus et Westmorlandicis ad Winandermere: in tectis et muris Cambriæ præcipue montosis abunde; D. Richardson. In littore marino prope Brakelsham copiose. D. Dillenius. R. Syn. circa Rydall in Westmorlandia. 4. VIII.

dasyphyllum. 8. *SEDUM* foliis oppositis ovatis obtusis carnosiss, caule infirmo, floribus sparsis. *Sp. pl. 431.*

Sedum foliis cordato-ovatis compressis sæpius oppositis, floribus sparsis. Sauv. monsp. 8.

Sedum foliis semiglobosis subovatis sessilibus quadrifariam imbricatis. Wackend. ultr. 391.

Sedum minus, circinato folio. Bauh. pin. 283. Hist. Ox. III. 473. f. 12. t. 7. f. 35. R. Syn. 271.

Sedum parvum, folio circinato, flore albo. Bauh. hist. III. 961.

Aizoon dasyphyllon. Lugd. 1133.

Anglis, round-leaved Stonecrop.

Habitat in muris et tectis. Grows plentifully upon the house and all the walls at Market-eit, near Market-freet. Mr. Th. Knowlton. R. Syn. ☉. VII.

villosum. 9. *SEDUM* caule erecto, foliis planiusculis, pedunculisque subpilosis. *Sp. pl. 432.*

Sedum purpureum pratense. Bauh. hist. III. 692. R. Syn. 270.

Sedum

Sedum palustre subhirsutum purpureum. *Bauh. pin.* 285.

Sedum minus palustre. *Ger. Em.* 515.

Sedum arvense f. *palustre flore rubente.* *Park.* 734.
Anglis, Marsh Stonecrop.

Habitat in pratis montosis humidis. On the moist rocks about Ingleborough-hill; on Hartside-hill, near Gamblesby, in the way to Osten in Cumberland. In et juxta montem Hincklehaw prope Settle. Dr. Richardson. R. Syn. ☉. VI.

O X A L I S. 515.

Cal. pentaphyllus. Petala unguibus connexa. Caps. angulis dehiscens pentagona.

1. *OXALIS* scapo unifloro, foliis ternatis, radice squamoso-articulata. *Hort. cliff.* 175. *Fl. suec.* 406. *Mat. med.* 216. *Sp. pl.* 233. *Acetofella.*

Oxys alba. *Ger. Em.* 1201. *R. Syn.* 281.

Trifolium acetosum vulgare. *Bauh. pin.* 330. *Park.* 746.

β *Trifolium acetosum vulgare, flore purpureo.* *Merr. pin.*

Anglis, Wood Sorrel.

Habitat in sylvis, ericetis, umbrosis. 4. IV.

Pharmac. Lujulæ Folia.

A G R O S T E M M A. 516.

Cal. monophyllus coriaceus. Petala quinque unguiculata; Limbo obtuso, indiviso. Caps. unilocularis.

1. *AGROSTEMMA* hirsuta, calycibus corollam æquantibus, petalis integris nudis. *Fl. suec.* 407. *Sp. pl.* 435. *Githaco.*

Lychnis segetum major. *Bauh. pin.* 204. *R. Syn.* 338.

Lychnoides segetum five *Nigellastrum.* *Park.* 632.

Pseudo-melanthium. *Dalech. hist.* 438. *Bauh. hist.* III. 341.

Anglis, Cockle.

Habitat in arvis passim. ☉. VI.

LYCH-

LYCHNIS. 517.

Cal. monophyllus, oblongus, lævis. *Petala* quinque unguiculata. Limbo bifido. *Capf.* trilocularis.

Flos cuculi. 1. LYCHNIS petalis quadrifidis, fructu subrotundo. *Sp. pl.* 436.

Caryophyllus pratensis laciniato flore simplici, sive *Flos cuculi.* *Bauh. pin.* 210.

Lychnis plumaria sylvestris simplex. *Park. par.* 253. *R. Syn.* 338.

Armeria pratensis mas et foemina. *Ger. Em.* 600. *Anglis*, Meadow Pinks, Wild Williams, Cuckow Flower.

Habitat in pratis et pascuis humidioribus. 4. VI.

Viscaria. 2. LYCHNIS petalis integris. *Sp. pl.* 436. *Fl. succ.* 409.

Silene floribus pentagynis, capsulis quinquelocularibus. *Hort. cliff.* 172.

Lychnis sylvestris viscosa rubra angustifolia. *Bauh. pin.* 205. *Park.* 636. *R. Syn.* 340.

Muscipula angustifolia. *Ger. Em.* 601.

Anglis, red German Catchfly.

Habitat in pratis montosis. Found by Mr. Lhwyd on the sides of Craig Wreidhin-hill, in Montgomeryshire. *R. Syn.* V. VI.

dioica. 3. LYCHNIS floribus dioicis. *Sp. pl.* 437. *Fl. succ.* 411.

foemina. *Cucubalus* caule composito, calycibus oblongo-ovatis. *Fl. lapp.* 182.

Lychnis sylvestris albo flore. *Ger. Em.* 468. *Park.* 630. *R. Syn.* 339.

Lychnis sylvestris alba simplex. *Bauh. pin.* 204.

mas. β *Lychnis sylvestris rubello flore.* *Ger. Em.* 469. *R. Syn.* 339.

Lychnis sylvestris flore rubro. *Park.* 631.

Lychnis sylvestris f. aquatica purpurea simplex. *Bauh. pin.* 204.

Anglis, α white Campion : β red Campion.

Habitat in sylvis et sepibus ubique. 4. V.—VII.

CERASTIUM. 518.

Cal. pentaphyllus. Petala bifida. Caps. unilocularis, apice dehiscens.

* *Capsulis oblongis.*

1. CERASTIUM foliis ovatis, petalis calyci æqualibus, *vulgatum.*
caulibus diffusis. *Fl. suec. 415. Loeft. hisp. 143.*

Myosotis foliis ex ovatis acuminatis, flore calyci æquali. *Hall. helv. 384.*

Myosotis arvensis hirsuta, parvo flore albo. *Vaill. parif. 142. t. 30. f. 1.*

Auricula muris quorundam flore parvo, vasculo tenui longo. *Baub. hist. III. 359.*

Alfina hirsuta magno (parvo potius) flore. *Baub. pin. 251.*

Alfina hirsuta myosotis. *Adv. 193. R. Syn. 349.*

Anglis, narrow-leaved Mouse-ear Chickweed.

Habitat in pratis et pascuis passim. ☉. VI.

2. CERASTIUM erectum villoso-viscosum. *Hort. cliff. viscosum.*
173. upf. 115. Roy. lugdb. 450. Fl. suec. 414. Sp. pl. 437.

Alfina hirsuta myosotis latifolia precocior. *Cat. angl. R. Syn. 348.*

Alfina viscosa. *Park. 768.*

Alfina hirsuta altera viscosa. *Baub. pin. 251.*

Myosotis hirsuta altera viscosa. *Vaill. parif. 142. t. 30. f. 1.*

Anglis, broad-leaved Mouse-ear Chickweed.

Habitat in pratis et pascuis siccioribus. ☉. IV. V.

3. CERASTIUM floribus pentandris, petalis emarginatis. *Hort. cliff. 173. Fl. suec. 416. Sp. pl. 438. candrium.*

Cerastium hirsutum minus, parvo flore. *Cat. giff. 80.*

R. Syn. 348. t. 15. f. 1.

Myosotis arvensis hirsuta minor. *Vaill. parif. 142. t. 30. f. 1.*

Alfina hirsuta minor. *Baub. pin. 251.*

Anglis,

Anglis, the least Moufe-ear Chickweed.
Habitat in pratis et pascuis circa Londinum frequens.
 ☉. IV.

- arvense.* 4. CERASTIUM foliis lineari-lanceolatis obtusis glabris, corollis calyce majoribus. *Sp. pl.* 438. *Fl. suec.* 417.
Myosotis arvensis hirsuta, flore majore. *Vaill. parif.* 141. *t.* 30. *f.* 4.
Caryophyllus arvensis hirsutus flore majore. *Bauh. pin.* 210. *R. Syn.* 348.
Caryophyllus holosteus. *Ger. Em.* 477. *arvensis hirsutus.* *Park.* 1339.
Holosteum caryophyllæum. *Tabern. ic.* 233.
Anglis, Corn Moufe-ear Chickweed.
Habitat in ericetis, pratis, pascuis, arvis, solo glareoso passim, supra Dupper's-hill prope Croydon copiose.
 4. V.

- alpinum.* 5. CERASTIUM foliis ovato-lanceolatis, caule diviso, capsulis oblongis. *Sp. pl.* 438. *Fl. suec.* 418.
Cerastium foliis ovato lanceolatis, corollis calyce majoribus. *Fl. suec.* 1. 380.
Alfina myosotis facie, Lychnis alpina, flore amplo niveo repens. *R. Syn.* 349. *t.* 15. *f.* 2.
Caryophylleus holosteus alpinus latifolius. *Bauh. pin.* 210. *pr.* 104.
Anglis, Mountain Moufe-ear Chickweed.
Habitat in montibus Arvonix. Nullibi frequentius quam juxta scaturagines aquarum montis excelsi Widnah boream versus, Dr. Richardson. *R. Syn.* 4. VI.

** *Capsulis subrotundis.*

- tomentosum.* 6. CERASTIUM foliis oblongis tomentosis, pedunculis ramosis, capsulis globosis. *Sp. pl.* 440.
Cerastium foliis lanceolato-linearibus subhirsutis, corolla calycem superante. *Sauv. monsp.* 142.
Caryophyllus holosteus tomentosus latifolius. *Bauh. pin.* 210. *pr.* 104.

Alfina

DECANDRIA PENTAGYNIA.

177.

Alfine myofotis lanuginosa alpina grandiflora, seu Auricula muris villosa flore amplo membranaceo. R.

Syn. 349.

Anglis, Woolly Moufe-ear Chickweed.

Habitat in rupe prope Clogwyn y Garnedh totius Cambriæ altissima juxta Llanberrys in Arvoniam copiose.

R. Syn. 4. V.

7. CERASTIUM foliis cordatis sessilibus, floribus solitariis fructibus pendulis. Sp. pl. 438. aquaticum.

Alfine aquatica major. Baub. pin. 251.

Alfine major repens perennis. Baub. hist. III. 362.

R. Syn. 347.

Alfine major. Ger. Em. 611. et maxima. Park. 759.

Anglis, Marsh Moufe-ear Chickweed.

Habitat in aquosis et ad ripas fluviorum. 4. VII.

S P E R G U L A. 519.

Cal. pentaphyllus. Petala quinque integra. Caps. ovata, unilocularis, quinquevalvis.

1. SPERGULA foliis verticillatis, floribus decandris. arvensis.

Sp. pl. 440. Roy. lugdb. 451.

Alfine Spergula dicta major. Baub. pin. 251. R. Syn.

351.

Saginæ Spergula. Ger. Em. 1125. major. Park.

562.

Anglis, Corn Spurrey.

Habitat in arvis solo glareoso. ☉. VIII.

2. SPERGULA foliis verticillatis, floribus pentandris. pentandria.

Loeff. hisp. 143. Sp. pl. 440.

Spergula foliis filiformibus verticillatis raris, feminibus nigris. Sauv. monsp. 167

Spergula annua femine foliaceo nigro, circulo membranaceo albo cincto. Cat. giff. 46. Ep. nat. cur. cent. V. 75. t. 4. R. Syn. 351. Hist. Ox. II. 551.

Alfine Spergulæ facie minima, feminibus emarginatis.

Tourn. 244. Vaill. paris. 8.

Anglis, small Spurrey.

Habitat in Hibernia locis arenosis observavit D. G. Sheppard. R. Syn. ☉. VII.

N

3. SPER-

nodosa.

3. SPERGULA foliis oppositis subulatis lævibus, caulis simplicibus. *Sp. pl.* 440. *Fl. suec.* 420.

Alfne palustris foliis tenuissimis, seu Saxifraga palustris anglica. *Ger Em.* 567. *R. Syn.* 350.

Alfne palustris, ericæ folio, polygonoides, articulis crebrioribus, flore albo pulchello. *Pluk. ph. t.* 7.

f. 4.

Alfne nodosa Germanica. *Baub. pin.* 251.

Saxifraga palustris anglica. *Park.* 427.

Anglis, knotted Spurrey, or English marsh Saxifrage.

Habitat in pratis humidis. *Ad ripas Thamefis prope*

Nine-elms; in ericeto Hampstediano. 24. VII.

VIII.

Classis

Classis XI.

DODECANDRIA
MONOGYNIA.

ASARUM. 522.

Cal. quadrifidus, germi-
ni infidens. *Cor.* o. *Caps.*
coriacea, coronata.

1. ASARUM foliis reniformibus obtusis binis. *Fl. succ. europæum.*
421. *Sp. pl.* 442.

Afarum foliis subcordatis petiolatis. *Hort. cliff.* 178.
Mat. med. 219.

Afarum. *Bauh. pin.* 197. *hist. III.* 548. *R. Syn.* 158.

Afarum vulgare. *Park.* 266. *Ger. Em.* 836.

Anglis, Afarabacca.

Habitat in nemorosis. In a hedge on the left hand as
soon as you are in the gate, the first field from Cher-
nell's green, going to Sir Tho. Seabright's beach-
wood, Th. Knowlton; in several woods in Lanca-
shire. *R. Syn.* 4. V.

Pharmac. Afari Radix, Folia.

LITHRUM. 532.

Cal. duodecim-fidus. *Petala* sex, calyci inserta. *Caps.*
bilocularis polysperma.

1. LITHRUM foliis oppositis cordato-lanceolatis, flo-
ribus spicatis dodecandris. *Sp. pl.* 446. *Fl. succ. salicaria.*
422.

Lithrum foliis oblongis acutis, floribus verticillatis. *Fl.*
lapp. 197.

Salicaria vulgaris purpurea, foliis oblongis. *Tourn.*
253. *R. Syn.* 367.

Lyfimachia purpurea. *Park.* 546. *Ger. Em.* 476.

DODECANDRIA DIGYNIA.

Lyfimachia spicata purpurea. Baub. pin. 246.

β *Lyfimachia purpurea trifolia caule hexagono. Spigel. Ifag.*

Anglis, purple-spiked Loofestriſe, or Willow-herb.

Habitat in ad ripas fluviorum, et in paludibus. 4. VII.

*Hyſſopi-
folia.*

2. LITHRUM foliis alternis linearibus, floribus hexandris. *Hort. upſ. 118. Sp. pl. 447.*

Salicaria hyſſopifolio latiore et anguſtiore. Tourn. 253.

R. Syn. 367.

Hyſſopifolia. Baub. pin. 218.

Gratiola anguſtifolia. Ger. Em. 581. f. minor. Park. 220.

Anglis, Graſs-poly, ſmall Hedge Hyſſop.

Habitat in aquoſis et ubi aquæ per hyemem ſtagnarunt, ſed rarius, ☉. VIII. IX.

D I G Y N I A.

AGRIMONIA. 534.

Cal. quinquedentatus, altero obvallatus. Petala quinque. Sem. duo infundo calycis.

*Eupato-
ria.*

1. AGRIMONIA foliis caulinis pinnatis; foliolis undique ferratis; omnibus minutis interſtinctis, fructibus hiſpidis. *Syſt. nat. 1047.*

Agrimonia foliis caulinis pinnatis, fructibus hiſpidis. Hort. cliff. 179. Fl. ſuec. 423. Sp. pl. 448.

Agrimonia. Ger. Em. 712. R. Syn. 202. vulgaris. Park. 594.

Eupatorium veterum ſ. Agrimonia. Baub. pin. 321. Anglis, Agrimony.

Habitat in marginibus agrorum et ad ſepes. 4. VI.

Pharmac. Agrimonix Herba.

T R I-

TRIGYNIA.

RESEDA. 535.

*Cal. monophyllus, partitus. Petala laciniata. Caps.
ore dehiscens unilocularis.*

1. RESEDA foliis omnibus trifidis, inferioribus pinna-*lutea.*
tis. *Sp. pl.* 449.

Reseda vulgaris. Bauh. pin. 100. *R. Syn.* 366.

Reseda minor seu vulgaris. Park. 823. *Ger. Em.*
277.

- β *Reseda crispa gallica. Bocc. pl. rar.* 76.

Anglis, base Rocket.

*Habitat in arvis, pratis et pascuis præsertim solo creta-
ceo. ☉. VII.*

2. RESEDA foliis lanceolatis integris, calycibus quadri-*Luteola,*
fidis. *Sp. pl.* 448.

Luteola. Ger. Em. 494. *R. Syn.* 366. *vulgaris. Park.*
602.

Luteola herba salicis folio. Bauh. pin. 100.

Anglis, wild Woad, Yellow-weed, or Dyers-weed.

Habitat in muris, locis sterilioribus et incultis frequens.

☉. VI.

EUPHORBIA. 536.

*Cor. tetra seu pentapetala Cal. monophyllus. Caps.
trilocca.*

* *Dichotomæ (umbella trifida aut nulla.)*

1. EUPHORBIA dichotoma, foliis integerrimis semi-*Peplis.*
cordatis, floribus solitariis axillaribus. *Diff. euph.*

21.

*Euphorbia foliis subrotundis obtusis, floribus solitariis
in foliorum alis. Guett. stamp. II.* 420

Tithymalus maritimus supinus annuus, Peplis dictus.

R. Syn. 313.

Peplis. Bauh. hist. III. 668. *Ger. Em.* 503. *Park.*
194.

DODECANDRIA TRIGYNIA.

Peplis maritima folio obtuso. *Bauh. pin.* 293.

Anglis, small purple Sea Spurge.

Habitat in littore maris arenoso inter Pensantiam et Marketjeu oppida Cornubiæ copiose. ☉. VI.

** *Umbella trifida*.

- Peplus*. 2. *EUPHORBIA* umbella trifida; dichotoma, involu-
cellis ovatis, foliis integerrimis obovatis petiolatis.
Diff. euph. 27. *Sp. pl.* 456.
Euphorbia foliis obverse ovatis integerrimis, umbella
universali trifida triphylla, partialibus dichotomis
diphyllis. *Hort. cliff.* 199. *Roy. lugdb.* 197. *Da-
lib. parif.* 156.
Tithymalus parvus annuus, foliis subrotundis non
crenatis, *Peplus* dictus. *R. Syn.* 313.
Peplus five *Efula* rotunda. *Bauh. pin.* 292. *Ger. Em.*
503.
Anglis, petty Spurge.
Habitat in cultis oleraceis. ☉. VII.

- exigua*. 3. *EUPHORBIA* umbella trifida: dichotoma, involu-
cellis lanceolatis foliis linearibus. *Diff. euph.* 29.
Sp. pl. 456.
Tithymalus leptophyllus. *Park.* 193. *R. Syn.* 313.
Tithymalus s. *Efula* exigua. *Bauh. pin.* 291.
Efula exigua Tragi. *Ger. Em.* 501.
Anglis, dwarf Spurge.
Habitat in arvis inter Camberwel et Dulwich; circa
Shooter's-hill in Cantio copiose. ☉. VII.

*** *Umbella quinquefida*.

- segetalis*. 4. *EUPHORBIA* umbella quinquefida: dichotoma, in-
volucellis cordatis acutis, foliis lineari-lanceolatis,
ramis floriferis. *Diff. euph.* 34. *Sp. pl.* 458.
Euphorbia inermis, foliis alternis linearibus acutis,
partialibus umbellulæ ovato-rhombeis, petalis bi-
cornibus. *Hort. ups.* 142.
Tithymalus segetum longifolius. *Cat. cant. app. R.*
Syn. 312.
Tithymalus annuus, lunato flore, linariæ folio longi-
ore. *Hist. Ox.* III. 339. *f. 10. t. 2. f. 3.*
Tithymalus

Tithymalus maritimo affinis, linariæ folio. Bauh. pin.
291.

Anglis, Corn Spurge.

Habitat in arvis, circa Triplow prope Cantabrigiam,
D. Lyons invenit. ☉. VII. VIII.

5. EUPHORBIA umbella quinquefida : trifida : dichotoma involucellis obovatis, foliis cuneiformibus ser. *Helioscopia.*
ratis. Diff. euph. 42. Sp. pl. 459.

Tithymalus helioscopius. Ger. Em. 498. Park. 189.
Bauh. pin. 291. R. Syn. 313. f. solifolius. Bauh.
hist. III. 669.

Anglis, Sun-spurge, or Wart-wort.

Habitat in cultis oleraceis. ☉. VII.

6. EUPHORBIA umbella quinquefida : dichotoma, involucellis cordatis, foliis lineari-lanceolatis obtusiusculis acuminatis reflexis. *Sp. pl. 458.*

Tithymalus maritimus minor. Portlandicus. R. Syn.
313. t. 24. f. 6.

Anglis, Portland Spurge.

Habitat in arenosis maris littoribus. Found by the Rev.
Mr. Stonefreet in the narrow neck of land which
joins Portland to Devonshire. R. Syn. ☉. VIII.

7. EUPHORBIA umbella subquinquefida : bifida, involucellis cordato-reniformibus, foliis sursum imbricatis. *Diff. euph. 53. Sp. pl. 458.*

Tithymalus maritimus. Bauh. pin. 291.

Tithymalus paralius. Ger. Em. 499. Bauh. hist. III.
674. R. Syn. 312.

Tithymalus paralius f. maritimus. Park. 184.

Anglis, Sea Spurge.

Habitat in littoribus maritimis frequens. 24. VII.
VIII.

8. EUPHORBIA umbella quinquefida : dichotoma, involucellis cordatis, foliis lanceolatis serrulatis pubescentibus, capsulis verrucosis. *Syst. nat. 1049.*

Tithymalus myrsinites fructu verrucæ simili. Bauh.
pin. 291.

DODECANDRIA TRIGYNIA.

Tithymalus verrucosus. *Bauh. hist. III. 673. Park. 181. R. Syn. 312.*

Anglis, rough fruited Spurge.

Habitat in arvis. Founa by Mr. Dale in Essex, and by Dr. Robinson near York. R. Syn. 4. VIII.

platyphyl- los. 9. *EUPHORBIA umbella quinquefida: trifida: dichotoma, involucellis carina pilosis, foliis ferratis lanceolatis sessilibus. Syst. nat. 1049.*

Euphorbia umbella quinquefida: subquadrifida: dichotoma, involucellis primariis tetraphyllis, foliis ferratis lanceolatis sessilibus. Diff. euph. 41. Sp. pl. 461.

Euphorbia inermis, foliis lanceolatis, umbella universaliumque prima quinquefidis: secunda trifida: reliquis bifidis. Roy. lugdb. 198. Dalib. paris. 156.

Tithymalus platyphyllos. Fuchf. hist. 813. Bauh. hist. III. 670. R. Syn. 312.

Tithymalus arvensis latifolius germanicus. Bauh. pin. 291.

Anglis, broad-leaved Spurge.

Habitat in arvis sed rarius. At Black Notley in Essex. R. Syn. In the corn fields between Harefield-Common and Battle's-Well, plentifully. Blackf. sp. bot. 4. VII.

**** *Umbella multifida.*

Amygdaloides. 10. *EUPHORBIA umbella multifida: dichotoma, involucellis perfoliatis orbiculatis, foliis obtusis. Diff. euph. 46. Sp. pl. 463.*

Tithymalus characias amygdaloïdes. Ger. Em. 500. Bauh. pin. 290. R. Syn. 312.

Tithymalus characias vulgaris. Park. 186.

Tithymalus sylvaticus toto anno folia retinens. Bauh. hist. III. 671.

Anglis, Wood Spurge.

Habitat in sylvis et sepibus passim. 4. IV.

Characias. 11. *EUPHORBIA umbella multifida: bifida, involucellis perfoliatis emarginatis foliis integerrimis, caule frutescente. Diff. euph. 44. Sp. pl. 463.*

En.

Euphorbia inermis, foliis lanceolatis, umbella universalis multifida, partialibus dichotomis, involucrellis, femibifidis perfoliatis. *Hort. cliff.* 199. *ups.* 142. *Roy. lugdb.* 197.

Tithymalus characias rubens peregrinus. *Bauh. pin.* 290.

Tithymalus characias monspéliensium. *Ger. Em.* 499. *Park.* 186. *R. Syn.* 312.

Anglis, red Spurge.

Habitat in sylvis et sepibus sed rarius. On the papermill pool-dam in Heywood-park, in Staffordshire, Dr. Plot. *R. Syn.* 2. VI.

12. *EUPHORBIA* umbella sexfida: dichotoma, involucrellis ovalibus, foliis integerrimis, ramis nullis, capsulis verrucosis. *Diff. euph.* 50. *Sp. pl.* 462. *Tithymalus hybernicus*, vasculis muricatis erectis. *Dill. elth.* 387. t. 290. f. 374.

Tithymalus hybernicus. *Merr. pin.* *R. Syn.* 312.

Tithymalus latifolius hispanicus. *Bauh. pin.* 291. *Tabern. hist.* 987.

Anglis, knotty-rooted Spurge.

Habitat in arvis. Amongst the corn near Twittenham-park; against Richmond, and near Otterspool; Mr. Doody. *R. Syn.* 4. VII.

SEMPERVIVUM. 538. *Dodecagynia.*

Cal. duodecim-partitus. *Petala* duodecim. *Caps.* duodecim polyspermæ.

1. *SEMPERVIVUM* foliis ciliatis, propaginibus patentibus. *Sp. pl.* 464. *Fl. suec.* 428.

Sempervivum foliis radicalibus carnosiss, caulinis imbricatis membranaceis, corymbo racemoso reflexo. *Hort. cliff.* 179.

Sempervivum majus. *Ger. Em.* 508. *R. Syn.* 269.

Sedum majus vulgare. *Bauh. pin.* 283. *hist.* III. 687. *Park.* 730.

Anglis, Houfeleek.

Habitat in testis passim. 4. VII.

Pharmac. *Sempervivi majoris Herba.*

Classis

Classis XII.

ICOSANDRIA

MONOGYNIA.

P R U N U S. 546.

Cal. quinquefida, inferus. *Petala* quinque. *Drupæ* nux futuris prominatis.

- insinitia.* 1. PRUNUS spinosa, foliis ovatis subtus villosis, pedunculis geminis. *Cent. pl.* 34.
 Prunus sylvestris major. *Bauh. hist.* I. 196. *R. Syn.* 462.
 Pruna sylvestria præcocia. *Bauh. pin.* 444.
 Prunus sylvestris fructu majore nigro. *R. Syn. ed. II.* 301.
 β Prunus sylvestris fructu rubro, acerbo et ingrato. *R. Syn. ed. II.* 302.
 γ Prunus sylvestris fructu majore albo. *R. Syn. ed. II.* 302.
Anglis, α. β. black Bullace-tree. γ. white Bullace-tree.
 Habitat in *sepibus.* h. IV.
- spinosa.* 2. PRUNUS spinosa, foliis lanceolatis. pedunculis solitariis. *Syst. nat.* 1057. *Fl. suec.* 432. *Sp. pl.* 275.
 Prunus sylvestris. *Ger. Em.* 1497. *Park.* 1033. *Bauh. pin.* 444. *hist.* I. 193. *R. Syn.* 462.
Anglis, the Black-thorn, or Sloe-tree.
 Habitat in *sepibus.* h. III. IV.
Pharmac. Pruni sylvestris cortex, Flores, Fructus.
- Padus.* 3. PRUNUS floribus racemosis, foliis deciduis basi subtus biglandulosis. *Sp. pl.* 473. *Fl. suec.* 431.
 Cerasus avium nigra et racemosa. *Ger. Em.* 1504. *R. Syn.* 463.

Cerasus

Cerasus racemosa sylvestris fructu non eduli. Bauh. pin. 451.

Cerasus avium racemosa. Park. 1517.

Anglis, wild Cluster-cherry, or Birds-cherry.

Habitat in sylvis et sepibus. In Westmoreland, and the mountainous part of the west-riding of Yorkshire, and in the woods of Derbyshire. R. Syn. In the lane going from Temple-mills to Epping-forest, Dr. Wilmer. h. V.

*4. PRUNUS umbellis sessilibus; foliis ovato-lanceolatis *Avium*. subtus pubescentibus. *Fl. suec. 429.*

Cerasus sylvestris, fructu rubro. Bauh. hist. I. 1220. R. Syn. 463.

Anglis, common wild Cherry-tree.

Habitat in sylvis et sepibus passim. h. IV.

*5. PRUNUS umbellis subpedunculatis, foliis ovato-lanceolatis glabris. *Syst. nat. 1057. Sp. pl. 474.*

Cerasus sylvestris fructu nigro. Bauh. hist. I. 220. R. Syn. 463.

Cerasus sylvestris ac major, fructu subdulci nigro colore inficiente. Bauh. pin. 450.

Anglis, Black Cherry.

Habitat in sylvis et sepibus. h. IV.

Pharmac. Cerasorum nigrorum Fructus.

D I G Y N I A.

C R A T Æ G U S. 547.

Cal. quinquefidus. Petala quinque. Bacca infera, disperma.

1. CRATÆGUS foliis ovatis inæqualiter serratis subtus *Aria*. tomentosis. *Fl. suec. 433. Sp. pl. 475.*

Cratægus inermis, foliis ellipticis serratis transversaliter sinuatis subtus villosis. Fl. lapp. 199.

Mespilus alni folio subtus incano, Aria Theophrasti dicta R. Syn. 453.

Aria Theophrasti. Ger. Em. 1327.

Sorbus.

- Sorbus sylvestris Aria Theophrasti dicta. *Park.* 1421.
 Sorbus alpina. *Bauh. hist.* I. 65.
 Alni effige, lanato folio, major. *Bauh. pin.* 452.
Anglis, the white Bean-tree.
 Habitat in sylvis et sepibus præsertim montosis. *h.* V.

- torminalis.* 2. CRATÆGUS foliis cordatis septangulis: lobis infimis divaricatis. *Sp. pl.* 476.
 Cratægus foliis cordatis acutis: lacinulis acutis ferratis. *Hort. cliff.* 187. *Roy. lugdb.* 271.
 Mespilus apii folio sylvestris non spinosa, seu Sorbus torminalis. *Bauh. pin.* 454. *R. Syn.* 453.
 Sorbus torminalis. *Cam. epit.* 162. *Ger. Em.* 1471.
 seu vulgaris *Park.* 1420.
Anglis, common wild Service-tree, or Sorb.
 Habitat in sylvis et sepibus. *h.* IV.
Pharmac. Sorbi torminalis *Fruetus.*

- Oxyacantha.* 3. CRATÆGUS foliis obtusis subtrifidis ferratis. *Syst. nat.* 1057.
 Cratægus foliis obtusis bitrifidis. *Fl. suec.* 434. *Sp. pl.* 477.
 Mespilus apii folio sylvestris spinosa, f. Oxyacantha. *Bauh. pin.* 454. *R. Syn.* 453.
 Oxyacanthus. *Ger. Em.* 1325.
 Spina appendix vulgaris. *Park.* 1025.
 β Mespilus apii folio, sylvestris spinosa, folio et fructu majore. *R. Syn.* 454.
Anglis, the White-thorn, or Haw-thorn.
 Habitat in sylvis et sepibus ubique. *h.* V.
Pharmac. Spinæ albæ f. Oxyacanthæ *Flores, Fruetus.*

T R I G Y N I A.

S O R B U S. 548.

Calix quinquefidus. *Petala* quinque. *Bacca* infera, trisperma.

- domestica.* *1. SORBUS foliis pinnatis subtus tomentosis. *Hall. helv.* 351. *Sp. pl.* 477.

Sorbis

Sorbus. *Bauh. hist. I. 59. Ger. Em. 1471. R. Syn. 452. fativa. Bauh. pin. 415.*

Sorbus legitima. *Park. 1420.*

Anglis, the true Service, or Sorb.

Habitat in montibus nemorosis. In the mountainous parts of Cornwall, Walter Moyle, Esq; in the Morelands in Staffordshire. *D. Plot. R. Syn. 4. IV. V.*

Pharmac. Sorbi Fructus.

2. SORBUS foliis pinnatis utrinque glabris. *Hall. herb. aucuparia. 250. Sp. pl. 477. Fl. suec. 435.*

Sorbus sylvestris foliis domesticæ similis. *Bauh. pin. 415. R. Syn. 452.*

Sorbus sylvestris five Fraxinus bubula. *Ger. Em. 1473.*

Sorbus aucuparia. *Bauh. hist. I. 62.*

Ornus, five Fraxinus sylvestris. *Park. 1419.*

Anglis, the Quicken-tree, or Mountain Ash.

Habitat in sylvis et sepibus subhumidis. *h. V.*

P E N T A G Y N I A.

P Y R U S. 550.

Calix quinquefidus. Petala quinque. Pomum inferum, quinqueloculare, polyspermum.

1. PYRUS foliis ferratis, floribus corymbosis. *Fl. suec. communis. 436. Sp. pl. 1200.*

Pyrus foliis ferratis, promissasi productis. *Hort. cliff.*

190. *Sp. pl. 479.*

Pyrus sylvestris. *Ger. Em. 1457. Park. 1500. Bauh. pin. 439.*

Pyraster seu Pyrus sylvestris. *Bauh. hist. I. 57. R. Syn. 452.*

β Pyrus fativa. *R. Syn. 452.*

Anglis, the wild Pear-tree.

Habitat in sylvis et sepibus. *h. IV.*

2. PYRUS foliis ferratis umbellis sessilibus. *Fl. suec. Malus. 437. Sp. pl. 1200.*

Pyrus foliis ferratis, promissasi concavis. *Hort. cliff.*

189. *Sp. pl. 479. Mat. med. 237.*

Malus

Malus sylvestris. Ger. Em. 1461. Park. 1502. R. Syn. 452.

Malus sylvestria. Baub. pin. 435.

β *Malus fativa.* R. Syn. 451.

Anglis, α the Crab-tree, or Wilding, β the Apple-tree.

Habitat in sylvis et sepibus. h. V.

Pharmac. mali sylvestris. Fructus.

M E S P I L U S. 549.

Cal. quinquefidus. Petala quinque. Bacca infera, 5-sperm.

Germanica. *1. MESPILUS inermis, foliis lanceolatis integerrimis subtus tomentosis, floribus terminalibus solitariis sessilibus. *Syst. nat.* 1058.

Mespilus inermis, foliis lanceolatis integerrimis subtus tomentosis, calycibus acuminatis. *Hort. cliff* 189. *ups.* 129. *Mat. med.* 236. *Roy. lugdb.* 270. *Boehm. lipf.* 174. *Sp. pl.* 478.

Mespilus. Ger. Em. 1452. R. Syn. 453. *vulgaris* Park. 1422.

Mespilus Germanica folio laurino non ferrato, f. *Mespilus sylvestris.* Baub. pin. 453.

Anglis, Medlar.

Habitat in sepibus. In all the hedges about Minehiville. Mr. Du Bois. R. Syn. h. V.

Pharmac. Mespili Fructus.

S P I R Æ A. 554.

Cal. quinquefidus. Petala quinque. Caps. polyspermæ.

Filipendula. 1. SPIRÆA foliis pinnatis: foliolis uniformibus serratis caule herbaceo, floribus cymosis. *Fl. suec.* 439. *Sp. pl.* 490.

Filipendula. Baub. hist. III. 2. 189. Ger. Em. 1058. Park. 434. R. Syn. 259.

Filipendula vulgaris. Baub. pin. 163.

Anglis, Drop-wort.

Habitat in pratis et pascuis montosis. Supra Dupper's-hill, prope Croydon. D. Caute. Juxta Clapham prope Londinum. u. VII. Whorwell woods, Hants.

Pharmac. Filipendulæ Radix Folia.

2. SPIRÆA

2. SPIRÆA foliis pinnatis: impari majore lobato, caule *Ulmaria*. herbaceo, floribus cymosis. *Fl. suec.* 490. *Sp. pl.* 440.
Ulmaria. Clus. hist. II. 198. *Bauh. hist. III.* 488.
R. Syn. 259.
Ulmaria vulgaris. Park. 592.
Regina prati. Ger. Em. 1043.
Barba capri floribus compactis. Bauh. pin. 164.
Anglis, Meadow-sweet.
Habitat in pratis humidis et ad ripas fluviorum passim.
 4. VI. VII.
Pharmac. Ulmariae Herba.

P O L Y G Y N I A.

R O S A. 556.

Petala quinque. Cal. urceolatus, quinquefidus, carnosus collo coarctatus. Sem. plurima, hispida, calycis interiori lateri affixa.

* *Germinibus subglobosis.*

1. ROSA germinibus globosis pedunculisque glabris, *Eglante-* caule aculeis sparsis rectis, petiolis scabris, foliolis *ria.* acutis. *Syst. nat.* 1062.
Rosa aculeata, foliis odoratis subtus rubiginosis. Hall.
helv. 350. *Sp. pl.* 491.
Rosa sylvestris odora. Ger. Em. 1272. *R. Syn.* 454.
seu Eglenteria flore simplici. Park. par. 418.
Rosa sylvestris foliis odoratis. Bauh. pin. 483.
Anglis, Sweet-briar, or Eglantine.
Habitat in sepibus. h. V. VI.
2. ROSA germinibus globosis glabris pedunculis caule *spinoseffi-* petiolisque aculeatissimis. *Syst. nat.* 1062. *ma.*
Rosa caule petiolisque aculeatis, calycis foliolis indivis. *Fl. suec.* 442. *Sp. pl.* 491.
Rosa pumila, spinosissima, foliis pimpinellæ glabris, flore albo. Bauh. hist. II. 40. *R. Syn.* 455.
Rosa sylvestris pomifera minor. Bauh. pin. 488.
Rosa pimpinella s. pomifera minor. Park. 1018.
 Rosa

Rosa pimpinellæ folio. *Ger. Em.* 1270.

- β Rosa Ciphiana f. Rosa pimpinellæ foliis minor nostras,
flore eleganter variegato. *Scot. ill.* 46.

Anglis, Burnet Rose.

Habitat in ericetis et locis arenosis. *h.* VI.

- arvensis.* 3. ROSA germinibus globosis pedunculisque glabris,
caule petiolisque aculeatis floribus subcymosis.

Rosa sylvestris altera minor, flore albo nostras. *R.*
Syn. 455.

Rosa arvensis candida. *Bauh. pin.* 484.

Rosa canina humilior fructu rotundiore. *D. Plot.*

- β Rosa canina sylvestris, unico flore et fructu. *R. Syn.*
Anglis, white-flowered Dogs-rose.

Habitat in sepibus. *h.* VII.

- villosa.* 4. ROSA germinibus globosis aculeatis, pedunculis hisp-
pidis caule aculeis sparsis, petiolis aculeatis, foliis
tomentosis. *Syst. nat.* 1062.

Rosa foliis utrinque villosis, fructu spinoso. *Hall. herb.*
350. *Sp. pl.* 491.

Rosa sylvestris pomifera major nostras. *R. Syn.* 455.

Rosa sylvestris pomifera major. *Bauh. pin.* 484.

Rosa pomifera major. *Park. par.* 418.

- β Rosa sylvestris fructu majore hispido. *R. Syn. ed. II.*
296.

Anglis, Apple-rose.

Habitat in montosis Cambrobritannicis, septentrionali-
bus Eboracensis, et Westmorlandici agri copiose.

h. VI.

** *Germinibus ovatis.*

- canina.* 5. ROSA germinibus ovatis pedunculisque glabris, caule
petiolisque aculeatis. *Syst. nat.* 1062.

Rosa caule aculeato, petiolis inermibus, calycibus fe-
mipinnatis. *Fl. suec.* 441. *Sp. pl.* 491.

Rosa sylvestris inodora seu canina. *Park.* 1017. *R.*
Syn. 454.

Rosa canina inodora. *Ger. Em.* 1270.

Rosa sylvestris vulgaris flore odorato incarnato. *Bauh.*
pin. 483.

Anglis, red-flowered Dogs-rose, or Hep-tree.

Habitat

Habitat in sepibus. h. V. VI.

Pharmac. Rosæ sylvestris Flores, Spongiolæ (insectorum niduli) f. Bedeguar, Cynosbati.

R U B U S. 557.

Cal. quinquefidus. Petala quinque. Bacca composita acinis monospermis.

* Frutescentes.

1. RUBUS foliis quinato-pinnatis ternatisque, caule aculeato, petiolis canaliculatis. Fl. suec. 446. Mat. med. 242. Sp. pl. 493.

Rubus caule erecto hispido, foliis ternatis. Fl. lapp. 204.

Rubus idæus spinosus. Bauh. pin. 479.

Rubus idæus. Ger. Em. 1272. Park. 557.

Rubus idæus spinosus fructu rubro. Bauh. hist. II. 59. R. Syn. 467.

Anglis, Raspberry-bush, Framboise, or Hindberry.

Habitat in sylvis, sepibus, montibus, saxosis et udis Walliæ, et Septentrionalium Angliæ. In Stokenchurch wood plentifully; in Westmorlandia copiosissime. h. V. VI.

2. RUBUS foliis ternatis subnudis: lateralibus bilobis, cæsius, caule aculeato tereti. Syst. nat. 1063. Fl. suec. 445. Sp. pl. 493.

Rubus caule aculeato reflexo perenni, foliis ternatis, Fl. lapp. 205.

Rubus repens fructu cæsius. Bauh. pin. 479. Ger. Em. 1271.

Rubus minor fructu cæruleo. Bauh. hist. II. 59. R. Syn. 467.

Rubus minor, chamærubus five Humirubus. Park. 1013.

Anglis, small Bramble, or Dewberry-bush.

Habitat in sylvis et sepibus passim. h. V. VII.

3. RUBUS foliis quinato-digitatis ternatisque, caule petiolisque aculeatis. Fl. suec. 444. Sp. pl. 493.

O

Rubus

ICOSANDRIA POLYGYNIA.

Rubus vulgaris major. *Park.* 1013. fructu nigro. *B. hist.* II. 57. *R. Syn.* 467.

Rubus vulgaris f. *Rubus* fructu nigro. *Bauh. pin.* 479.

β *Rubus vulgaris major*, fructu albo. *R. Syn.* 467.
Anglis, common Bramble.

Habitat in sylvis et sepibus ubique. v. V.—IX.

** *Herbacei.*

saxatilis. 4. *RUBUS* foliis ternatis nudis flagellis reptantibus herbaceis. *Sp. pl.* 494. *Fl. suec.* 447.

Rubus saxatilis. *Ger. Em.* 1273.

Rubus alpinus saxatilis. *Park.* 1014.

Chamærubus saxatilis. *Bauh. pin.* 479. *R. Syn.* 261.
Anglis, Stone Bramble.

Habitat inter lapides ad latera montium in comitatibus Eboracensi, Westmorlandico, et Cumberlandico.
4. VI.

Chamæ-
-morus.

5. *RUBUS* foliis simplicibus lobatis, caule unifloro dioico. *Fl. suec.* 449. *Sp. pl.* 494.

Rubus caule bifolio unifloro, foliis simplicibus. *Fl. lapp.* 208. t 5. f. 1.

Chamæmorus. *Clus. hist.* I. 118. *Ger. Em.* 173.
R. Syn. 260.

Chamæmorus anglica. *Park.* 1014.

Chamæmorus foliis ribes *anglica.* *Bauh. pin.* 480.

Anglis, Cloud-berries, Knot-berries, or Knout-berries.

Habitat in paludibus turfosis supra montes Eboracenses, Westmorlandicos, Cumberlandicos, Arvonicos. 4. V. VI.

FRAGARIA, 558.

Cal. decem-fidus. *Petala* quinque. *Receptaculum* femininum ovatum, baccatum deciduum.

vesta. 1. *FRAGARIA* flagellis reptans. *Sp. pl.* 494. *Fl. suec.* 450.

Fragaria. *Ger. Em.* 998. *R. Syn.* 254. vulgaris.
Park. 758. *Bauh. pin.* 326. *Fl. lapp.* 209.

Fragaria minor, dulci fructu. *Radb. cat.* 17. *hort.* 43.
β *Fra*

- β *Fragraria fructu hispido.* *Ger. Em.* 998.
Anglis, Strawberry.
Habitat in sylvis et ad sepes frequens. 4. IV. V.
2. *FRAGARIA* caule decumbente repente. *Roy. lugdb. sterilis.*
 274. *Dalib. parif.* 147. *Sauv. monsp.* 177. *Sp. pl.*
 495.
Fragaria sterilis. *Baub. pin.* 327. *R. Syn.* 254.
Fragaria minime vesca. *Park.* 758. f. *steriles Lob. ic.*
 696. *Ger. Em.* 998.
Anglis, barren Strawberry.
Habitat in pascuis sterilioribus passim. 4. IV.

P O T E N T I L L A.

Cal. decemfidus. Petala quinque. Sem. subrotunda,
nuda, receptaculo parvo exsucco affixa.

* *Foliis pinnatis.*

1. [*POTENTILLA* foliis pinnatis, caule fruticoso. *fruticosa.*
Hort. cliff. 193. *Fl. suec.* 451. *Sp. pl.* 495.
Pentaphylloides rectum fruticosum Eboracense. Hist.
Ox. II. 193. f. 2. t. 23. f. 5. *Hortul. angl.* 54.
 t. 55.
Pentaphylloides fruticosa. R. Syn. 256. *Cat. ang.*
 228. *ic.*
Anglis, Shrub Cinquefoil.
Habitat ad ripam meredionalem Tesæ fluvii infra vicum
Thorp dictum variis in locis, nec non infra cænobium
Athelstani, vulgo Eggleston-Abbey, in agro Eborac-
ensi. Mr. Lawson observed thousands of these plants
by Mickle-force, in Teafdale. R. Syn. 2. VI.
2. *POTENTILLA* foliis pinnatis ferratis, caule repente. *Argentina*
Fl. lapp. 210. *Fl. suec.* 452. *Sp. pl.* 495.
Potentilla. Baub. pin. 321. *Park.* 593. *Cam. epit.*
 758.
Pentaphylloides argentina dicta. R. Syn. 256.
Argentina. Dod. pempt. 600. *Ger. Em.* 973.
Anglis, Silver-weed, or wild Tansey.
Habitat ad semitas, vias, inque pascuis depressis, ubi aquæ
hyemales præsertim stagnant frequens. 4. VI. VII.
Pharmac. Argentinæ Herba.

- rupestris*. 3. **POTENTILLA** foliis pinnatis alternis, foliolis quinatis ovatis crenatis, caule erecto. *Sp. pl.* 496. *Fl. suec.* 453.
 Pentaphylloides erectum *Bauh. hist. II.* 398. *Hist. Ox. II.* 193 *f. 2. t. 20. f. 1. R. Syn.* 255.
 Pentaphyllum fragiferum. *Ger. Em.* 991. *Park.* 397. *Clus. hist. II.* 107.
 Quinquefolium fragiferum. *Bauh. pin.* 326.
Anglis, upright bastard Cinquefoil.
 Habitat ad latera montis cujusdam Craig Wreidhin dicti in comitatu montis Gomerici Walliæ, D. Lhwyd. *R. Syn.* 4. VII.

* *Foliis quinatis.*

- argentea*. 4. **POTENTILLA** foliis quinatis cuneiformibus incisissimis subtus tomentosis, caule erecto. *Sp. pl.* 497. *Fl. suec.* 454.
 Potentilla foliis digitatis, caule erecto corymboso. *Fl. suec. ed. I.* 417.
 Pentaphyllum erectum, foliis profunde factis subtus argenteis, flore luteo. *Bauh. hist.* 598. *R. Syn.* 255.
 Pentaphyllum rectum minus. *Park.* 400.
 Quinquefolium folio argenteo. *Bauh. pin.* 235.
 Quinquefolium tormentillæ facie. *Ger. Em.* 987.
Anglis, Tormentil Cinquefoil.
 Habitat in pratis et pascuis solo glareoso passim. 4. VI.
- opaca*. 5. **POTENTILLA** foliis radicalibus quinatis cuneiformibus ferratis caulinis suboppositis, ramis filiformibus decumbentibus. *Cent. pl.* 35. *Syst. nat.* 1064.
 Pentaphyllum parvum hirsutum. *Bauh. hist. II.* 598. *R. Syn.* 255.
 Pentaphyllum incanum minus repens. *Ger. Em.* 989.
 Quinquefolium minus repens lanuginosum. *Park.* 399.
 Quinquefolium minus repens lanuginosum luteum. *Bauh. pin.* 225.
Anglis, small rough Cinquefoil.
 Habitat circa Kippax in agro Eboracensi. 4. VI.

6. **POTENTILLA** foliis radicalibus quinatis acute serratis retusis, caulinis ternatis, caule declinato. *Sp. pl.* 498. *Fl. suec.* 455.
 Pentaphyllum seu Quinefolium minus. *Tabern. ic.* 123.
 Pentaphyllum tormentillæ facie. *Rudb. cat.* 32. *hort.* 85. *Till. ab.* 52.
 Quinefolium minus repens luteum. *Bauh. pin.* 325. *R. Syll.* 213.
Anglis, Spring Cinquefoil.
 Habitat in pascuis sterilioribus prope Giggleswick in comitatu Eboracensi. 4. V. VI.
7. **POTENTILLA** foliis digitatis, caule repente, perunculis unifloris. *Fl. suec.* 456 *Sp. pl.* 499.
 Potentilla foliis digitatis longitudinaliter patentiserratis, caule repente. *Hort. cliff.* 194. *Gron. virg.* 163.
 Pentaphyllum vulgatissimum. *Park.* 398. *R. Syn.* 255.
 Quinefolium majus. *Ger. Em.* 987. repens. *Bauh. pin.* 325.
Anglis, common Cinquefoil, or five-leaved Grass.
 Habitat ad semitas, vias, &c. ubique. 4. VI.
Pharmac. Pentaphylli Radix.
8. **POTENTILLA** foliis quinatis apice conniventiserratis, caulibus filiformibus procumbentibus, receptaculis hirsutis. *Sp. pl.* 498.
 Quinefolium album majus. *Bauh. pin.* 325.
 Quinefolium sylvaticum majus flore albo. *Ger. Em.* 987.
Anglis, white Cinquefoil.
 Habitat in Wallia, D. Haviland. 4. VIII.

TORMENTILLA. 560.

Cal. octofidus. *Petala* quatuor. *Sem.* subrotunda, nuda receptaculo parvo exsucco affixa.

1. **TORMENTILLA** caule erecto, foliis sessilibus erecta.
 Tormentilla caule erecto. *Sp. pl.* 500. *Fl. suec.* 459.

*An infusion
of this plant is
the usual tea of
the Tartars of
Tomsk, a great
astringent.*

Tormentilla. *Ger. Em.* 992. *Bauh. hist. II.* 598. *Fl. lapp.* 213. *Hort. cliff.* 194. *Mat. med.* 249. *R. Syn.* 257. *vulgaris. Park.* 394.

Tormentilla sylvestris. *Bauh. pin.* 326.

Anglis, Tormentil, or Septfoil.

Habitat in pascuis sterilioribus ubique. 4. VI.

Pharmac. Tormentillæ Radix, Folia.

reptans.

2. **TORMENTILLA** caule repente, foliis petiolatis
Tormentilla caule repente. *Sp. pl.* 500.

Tormentilla reptans. *Pet. h. br. t.* 41. *f.* 10. *R. Syn.* 257.

Pentaphyllum reptans alatum foliis profundius ferratis.
Hist. nat. ox. c. 6. *p.* 7. *t.* 9. *f.* 5.

Pentaphyllum minus viride flore aureo tetrapetalo, rad-
iculas in terram e geniculis demittens. *Hist. ox.*
190. 12.

β Pentaphyllum aureum minus, sylvaticum, nostras, fo-
liis tripartito divisus ex cauliculorum geniculis radi-
cescens. *Pluk. alm.* 285.

Anglis, creeping Tormentil.

Habitat in sylvis et pascuis sterilioribus passim. 4. VII.

GEUM. 561.

Cal. decemfidus. *Petala* quinque. *Sem.* cauda pi-
lofa.

urbanum. 1. **GEUM** floribus erectis, fructu globofo : aristis un-
cinatis nudis foliis lyratis. *Syst. nat.* 1065. *Fl. suec.*
460. *Sp. pl.* 501.

Caryophyllata. *Ger. Em.* 995. *R. Syn.* 253. *vulga-*
ris. Bauh. pin. 321. *Park.* 136. flore parvo luteo.
Bauh. hist. II. 398.

β Caryophyllata vulgaris majore flore. *Bauh. pin.* 321.
Anglis, common Avens, or Herb-bennet.

Habitat in sylvis et sepibus frequens. 4. VI. VIII.

Pharmac. Caryophyllatæ Radix.

virale.

2. **GEUM** floribus nutantibus, fructu oblongo : aristis
plumosis. *Fl. suec.* 461. *Sp. pl.* 501.

Caryophyllata montana purpurea. *Ger. Em.* 995. *R.*
Syn. 253.

Ca-

Caryophyllata montana f. *palustris purpurea*. *Park.*
136.

Caryophyllata aquatica nutante flore. *Bauh. pin.* 321.
Anglis, Water Avens.

Habitat in montosis Septentrionalibus Angliæ circa Settle,
Ingleton, &c. also in Wales, about Snowdon-hill,
&c. *R. Syn.* 4. VII.

DRYAS. 562.

Cal. octofidus. Petala octo. Sem. caudata, pilosa.

1. DRYAS pentapetala, foliis pinnatis. *Amoen. acad. pentape-*
II. 353. Sp. pl. 501. tala.

Caryophyllata pentaphyllæa. Bauh. hist. II. 398.
Park. 137. R. Syn. 254.

Caryophyllata alpina pentaphyllæa. Ger. Em. 995.

Caryophyllata alpina quinquefolia. Bauh. pin. 322.
Anglis, Cinquefoil Avens.

Habitat in the den of Petaick in Scotland, D. Sibbald.
R. Syn. 4.

2. DRYAS octopetala : foliis simplicibus. *Fl. suec. 462. octopetala.*
Sp. pl. 401.

Caryophyllata alpina chamædryos folio. Hist. ox. II.
432. R. Syn. 252.

Chamædryos spuria montana cisti flore. Park. 106.

Chamædryos alpina, cisti flore. Bauh. pin. 248.

Chamædryos alpina, flore fragariæ albo. Bauh. hist. III.
290.

Teucrium alpinum, cisti flore. Ger. Em. 659.

Anglis, Mountain Avens.

Habitat in montosis. Found by Mr. Heaton in Ireland,
on the mountains between Gort and Galloway. Hujus
plantæ speciminibus meditavit, D. Lhwyd, in monti-
bus Hibernicis juxta Sligo collectis, quæ et in omni-
bus altissimis Scotiæ occidentalis montibus abundat, ut
mihi retulit idem. *R. Syn. 4. VII.*

COMARUM. 563.

Cal. decemfidus. *Petala* quinque, calyce minora.
Receptaculum spongiosum, persistens.

palustre. I. COMARUM. *Fl. lapp.* 214. *suec.* 463. *Sp. pl.* 502.
 Pentaphylloides palustre rubrum. *Tourn.* 298. *R. Syn.*
 256.

Pentaphylloides rubrum palustre. *Ger. Em.* 987.

Quinquefolium palustre rubrum. *Bauh. pin.* 326.

β Pentaphylloides palustre rubrum crassis et villosis foliis
 Suecicum et Hybernicum. *Pluk. alm.* 284. t. 212.

f. 2.

Anglis, purple marsh Cinquefoil.

Habitat in palustribus putridis et coenosis non infrequens.
R. Syn. 4. VI.

Classis

*Classis XIII.*POLYANDRIA
MONOGYNIA.* *Flores tetrapetali.*

ACTÆA. 568.

Cal. tetraphyllus. *Cor.* tetrapetala. *Bacca* unilocularis. *Sem.* semiorbiculata.1. ACTÆA racemo ovato, fructibus baccatis. *Sp. pl. spicata.*509. *Fl. suec.* 464.*Actæa* caule inermi. *Fl. lapp.* 217.*Christophoriana.* *Clus. hist. II.* 86. *Ger. Em.* 969.*Till. ic.* 148. *R. Syn.* 262.*Christophoriana vulgaris.* *Park.* 379.*Aconitum racemosum.* *Bauh. pin.* 182.*Anglis,* Herb Christopher, or Bane-berries.*Habitat in dumetis infra, Malhamcove, juxta murum aquilonem versus, prope rupes ubi Fraxini juniores crescunt, reperiuntur hujus plantæ nonnullæ, D. Richardson. R. Syn. in sylvula prope Clapham in agro Eboracenfi. 4. V. VI.*

CHELIDONIUM. 572.

Cor. tetrapetala. *Cal.* diphyllus. *Siliqua* unilocularis, linearis.1. CHELIDONIUM pedunculis umbellatis. *Sp. pl. majus.*505. *Mat. med.* 252.*Chelidonium majus.* *Fuchf. hist.* 865. *Ger. Em.* 1069.*Chelidonium majus vulgare.* *Park.* 616. *Bauh. pin.*

144.

*Papaver corniculatum luteum, Chelidonia dictum.**R. Syn.* 309.β *Chelidonium majus foliis quernis.* *Bauh. pin.* 144.*Anglis,*

Anglis, Celandine.

Habitat *in ruderatis*. 4. V. VI.

Pharmac, Chelidonii majoris *Radix*, *Herba*.

- Glaucium*. 2. CHELIDONIUM pedunculis unifloris, foliis amplexicaulibus sinuatis, caule glabro. *Sp. pl.* 506.
 Papaver corniculatum luteum. *Park.* 261. *Bauh. pin.* 171. *hist. III.* 398. *R. Syn.* 309.
 Papaver cornutum flore luteo. *Ger. Em.* 367.
Anglis, yellow horned Poppy.
 Habitat *in littoribus maritimis*; *in insula* Shepey.
 ☉. VII.

- hybridum*. 3. CHELIDONIUM pedunculis unifloris, foliis pinnatifidis linearibus, caule lævi. *Sp. pl.* 506.
 Papaver caule ramoso, foliis linearibus pinnatifidis, capsulis sulcatis. *Hort. upf.* 136.
 Papaver corniculatum violaceum. *Bauh. hist. III.* 393. *pin.* 172. *R. Syn.* 309.
 Papaver cornutum flore violaceo. *Ger. Em.* 367.
 Papaver corniculatum flore violaceo. *Park.* 262.
Anglis, violet-coloured horned Poppy.
 Habitat *in agro Cantabrigiensi inter segetes inter Swafham et Burwel vicos*. *R. Syn.* ☉. VIII.

P A P A V E R. 573.

Cor. tetrapetala. *Cal.* diphyllus. *Capsf.* unilocularis, substigmatate persistente poris dehiscens.

* *Capsulis hispidis*.

- hybridum*. 1. PAPAVER capsulis globosis torosis hispidis, caule folioso multifloro. *Sp. pl.* 506.
 Papaver foliis ternato-pinnatifidis, capsulis sulcatis hispidis. *Hort. cliff.* 201.
 Papaver laciniato folio, capitulo hispido rotundiores. *R. Syn.* 308.
 Argemone capitulo rotundiores. *Park.* 369.
 Argemone capitulo breviores. *Bauh. pin.* 172.
 Argemone capitulo torulo. *Ger. Em.* 373.
Anglis, round rough-headed Poppy.
 Habitat *in arvis sed rarius*. ☉. VI. VII.

2. PAPAVER capsulis clavatis hispidis, caule folioso *Argemone*. multifloro. *Sp. pl.* 506. *Fl. suec.* 466.
 Papaver foliis ternato-pinnatis, fructu angulato. *Hort. cliff.* 201. *Ray. lugdb.* 419. *Sauv. monsp.* 252.
 Papaver laciniato folio capitulo hispido longiore. *R. Syn.* 308.
Argemone capitulo longiore. *Ger. Em.* 373. *Bauh. pin.* 172. *Park.* 370.
Anglis, long rough-headed Poppy.
 Habitat in arvis. ☉. VI.

** *Capsulis glabris.*

3. PAPAVER capsulis glabris globosis, caule piloso *Rhoeas*. multifloro, foliis pinnatifidis incis. *Sp. pl.* 507. *Fl. suec.* 467.
 Papaver laciniato folio, capitulo brevior glabro, annum *Rhoeas* dictum. *R. Syn.* 308.
 Papaver *rhoeas*. *Ger. Em.* 401.
 Papaver erraticum *rhoeas* five sylvestre. *Park.* 367.
 Papaver erraticum majus. *Bauh. pin.* 171.
Anglis, red Poppy, or Corn Poppy, or round smooth-headed Poppy.
 Habitat inter segetes. ☉. VI. VII.
Pharmac. Papaveris erratici Flores.
4. PAPAVER capsulis oblongis glabris, caule multiflo- *dubium*. ro setis adpressis, foliis pinnatifidis incis. *Sp. pl.* 1196. *Fl. suec.* 467.
 Papaver laciniato folio capitulo longiore glabro. *R. Syn.* 309.
Argemone capitulo longiore glabro. *Hist. ox. II.* 279. *f. 3. t. 14. f. 11.*
Anglis, long smooth-headed Poppy.
 Habitat in arvis circa Stockwell prope Londinum.
 ☉. VI.
5. PAPAVER capsulis glabris oblongis, caule multiflo- *cambri-* ro lævi, foliis pinnatis incis. *Sp. pl.* 508. *cum.*
 Papaver cambricum perenne, flore sulphureo. *Dill. elth.* 300. *t. 223. f. 290.*

Pa-

POLYANDRIA MONOGYNIA.

Papaver luteum perenne, laciniato folio, Cambrobri-
tanicum. *R. Syn.* 309.

Argemone lutea Cambrobritannica. *Park.* 369.

Anglis, yellow Poppy.

Habitat in montosis cambrobritannicis frequens; circa

Kendal in Westmorelandia copiose. *4. VI.—VIII.*

*somnife-
rum.*

*6. PAPAVER calycibus capsulisque glabris, foliis am-
plexicaulibus incis. *Sp. pl.* 508.

Papaver caule multifloro, foliis simplicibus glabris in-
cis caulinis. *Hort. ups.* 135. *Mat. med.* 250.

Papaver sylvestre. *Ger. Em.* 370. *R. Syn.* 308.

Papaver hortense semine albo. *Bauh. pin.* 170.

Anglis, wild Poppy.

Habitat in incultis passim. *○. VII.*

Pharmac. *Papaveris albi Capita, Semina.*

** Flores pentapetali.

T I L I A. 587.

Cor. pentapetala. *Cal.* quinquepartitus. *Bacca* ficca,
globosa, quinquelocularis, quinquevalvis, basi de-
hiscens.

europæa. *1. TILIA floribus nectario destitutis. *Sp. pl.* 514.

Tilia vulgaris platyphyllos. *Bauh. hist. I.* 133. *R.*
Syn. 473.

Tilia foemina. *Ger. Em.* 1483. *major.* *Park.* 1407.

Tilia foemina folio majore. *Bauh. pin.* 426.

β *Tilia foemina* folio minore. *Bauh. pin.* 426.

γ *Tilia* foliis molliter hirsutis, viminibus rubris, fructu
tetragono. *R. Syn.* 473.

δ *Tilia ulmifolia*, semine hexagono. *Merr. pin.*

Anglis, the Lime, or Linden-tree.

Habitat in sylvis sepibus et areis. *h. VII.*

C I S T U S. 598.

Cor. pentapetala. *Cal.* pentaphyllus, foliolis duo-
bus minoribus. *Capsula* triquetra trivalvis.

* *Exstipulatæ herbacæ.*

guttatus. 1. CISTUS herbaceus exstipulatus, foliis oppositis tri-
nerviis, racemis ebracteatis. *Sp. pl.* 526.

Cistus

Cistus annuus, foliis radicalibus oppositis lanceolatis trinervis, ramiferis ternatis linearibus. *Sauv. monsp.* 149.

Cistus flore pallido, punicante macula insignito. *Baub. pin.* 465. *R. Syn.* 341. 342

Cistus annuus flore maculato. *Ger. Em.* 1281.

Cistus annuus flore guttato. *Park.* 661.

Anglis, annual Cistus.

Habitat in pratis arenosis. In the isle of Jersey, on the west side, near Grosnez-castle, Dr. Sherard, *R. Syn.* in Somersetshire, Mr. Brewer. *○. VI.*

**** Stipulatæ suffruticosæ.**

2. CISTUS suffruticosus procumbens, stipulis lanceolatis, foliis oblongis revolutis subpilosis. *Sp. pl.* 528. *Helianthemum.*

Cistus stipulis quaternis foliis oblongis utrinque nudis caule procumbente. *Hort. cliff.* 206. *Roy. lugdb.* 476.

Helianthemum vulgare. *Park.* 656. *R. Syn.* 341.

Helianthemum anglicum luteum. *Ger. Em.* 1282.

Chamæcistus vulgaris flore luteo. *Baub. pin.* 465.

β *Helianthemum* foliis majoribus: flore albo. *Baub. hist.* II. 16.

γ *Helianthemum* ampliore folio; flore roseo. *D. Sherard.*

Anglis, dwarf Cistus, or little Sun-flower.

Habitat in montosis siccioribus, præsertim cretaceis. 24. VII.

3. CISTUS suffruticosus procumbens stipulatus, foliis surrei-ovato-oblongis subpilosis, petalis lanceolatis. *Sp. nus. pl.* 527.

Helianthemum vulgare petalis florum perangustis. *R. Syn.* 341. *Dill. elth.* 177. t. 145. f. 174.

Anglis, narrow-leaved Cistus.

Habitat in pratis prope Croydon invenit D. Du Bois. *R. Syn.* 24. VII.

4. CISTUS suffruticosus stipulatus procumbens, foliis poliiifolius. oblongo-ovatis incanis, petalis crenatis.

Cha-

Chamæcistus montanus polii folio. *D. Pluk. R. Syn.* 342.

Cistus humilis alpinus durior, polii nostratis folio can-
dicante. *Pluk. ph. 23. f. 6.*

Anglis, Mountain dwarf Cistus.

Habitat upon Brent-downs in Somersetshire, near the
Severn Sea, D. Plukenet. R. Syn. 4. VII.

*** *Exstipulatæ suffruticosæ.*

hirsutus. 5. *CISTUS suffruticosus procumbens exstipulatus, foliis*
oppositis oblongis revolutis pilosis, floribus racemo-
sis.

Helianthemum alpinum folio Pilosellæ minoris Fuchsi.
Bauh. hist. II. 18. R. Syn. 342.

Anglis, hoary dwarf Cistus.

Habitat uno circiter miliari a Kendalia, loco saxoso Be-
tram-Beuke dicto occidentem versus copiose, D. Rich-
ardson. At Buckbarrow-Bank-Scar, betwixt Brig-
steer and Conswick; also on the rocks about Cartmel-
Wells in Lancashire, plentifully, observed by Mr.
Fitz-Roberts. 4. VII.

*** *Flores polypetali.*

N Y M P H Æ A. 579.

Cor. polypetala. Cal. tetra f. pentaphyllus. Bacca mul-
tilocularis, truncata.

lutea. 1. *NYMPHÆA calyce magno pentaphyllo. Fl. lapp.*
218. suec. 469. Sp. pl. 510.

Nymphæa lutea. Ger. Em. 819. R. Syn. 368.

Nymphæa major lutea. Bauh. pin. 131. Park. 1252.
Anglis, yellow Water Lilly.

Habitat in fluviis stagnis, fossis; inter Rotherhithe et
Deptford. 4. VIII.

alba. 2. *NYMPHÆA foliis cordatis integerrimis, calyce qua-*
drifido. Sp. pl. 510. Fl. suec. 470.

Nymphæa calyce tetraphyllo, corolla multiplici. Fl.
lapp. 219.

Nymphæa alba. Ger. Em. 819. R. Syn. 368. major.
Bauh. pin. 193.

Nymphæa

Nymphæa alba major vulgaris. Park. 1251.

Anglis, white Water Lilly.

Habitat in fluviis et stagnis sed rarius. 4. VII.

TRI G Y N I A.

DELPHINIUM. 602.

Cal. nullus. Petala quinque. Nectarium bifidum, pone cornutum.

1. DELPHINIUM nectariis monophyllis caule subdivi- *Consolida.*
fo. *Syst. nat. Fl. suec. 475. Sp. pl. 530.*

Delphinium fegetum, flore cœruleo. Tourn. 426. R. Syn. 273.

Consolida regalis arvensis, flore cœruleo. Bauh. pin. 142.

Consolida regalis sylvestris. Ger. Em. 1083.

Anglis, wild Larkspur.

*Habitat in arvis. Found in great plenty by Mr. J. She-
rard, amongst the corn, in Swafham-field, in Cam-
bridgehire. R. Syn. 6. VI.*

P E N T A G Y N I A.

A Q U I L E G I A. 605.

*Cal. 0. Petala quinque. Nectaria quinque cornicu-
lata, inter petala. Caps. quinque distinctæ.*

1. AQUILEGIA nectariis incurvis. *Mat. med. 272. vulgaris.*
Sp. pl. 533. Fl. suec. 478.

*Aquilegia. Cæsalp. syst. 589. Fuchsf. hist. 102. Hort.
cliff. 215.*

*Aquilegia flore simplici. Bauh. hist. III. 484. R. Syn.
273.*

Aquilegia sylvestris. Bauh. pin. 144.

Aquilegia simplex. Park. 1367.

Aquilegia flore cœruleo. Ger. Em. 1093.

Anglis, common Columbines.

Habitat in sylvis et dumetis. 4. VI.

Pharmac. Aquilegiæ Herba, Semina.

2. AQUI-

- alpina.* 2. **AQUILEGIA** nectariis rectis petalo lanceolato brevioribus. *Sp. pl.* 533.
Aquilegia montana magno flore. *Bauh. pin.* 144. *pr.* 75. *Burf. VII.* 108. *Bauh. hist. III.* 484. *R. hist.* 707.
Anglis, Mountain Columbines.
Habitat in sylvis montosis in comitatu Westmorlandico.
 4. VI.

HEXAGYNIA.

STRATIOTES. 607.

- Spatha* diphylla. *Perianthium* trifidum. *Petala* tria.
Bacca sexocularis, infera.
Aloides. 1. **STRATIOTES** foliis ensiformi-triangularis ciliato-aculeatis. *Sp. pl. Fl. suec.* 479.
Stratiotes foliis aloes, semine longo. *Gundelsh. ap. Johren. R. Syn.* 290.
Stratiotes sive *militaris aizoides.* *Park.* 1249.
Militaris aizoides. *Ger. Em.* 826.
Aloe palustris. *Bauh. pin.* 286.
Anglis, Water Aloe, or Fresh Water Soldier.
Habitat in fluentis pigrioribus et palustribus fossis in insula Eliensi copiose. R. Syn. 4. VI.

POLYGYNIA.

ANEMONE. 614.

- Cal. o. Petala* sex-novem. *Sem.* plura.
nemorosa. 1. **ANEMONE** feminibus acutis, foliolis incisfis, caule unifloro. *Hort. cliff.* 224. *Fl. suec.* 485. *Sp. pl.* 541.
Anemone nemorosa, flore majore. *Bauh. pin.* 176.
Anemone nemorum album. *Ger. Em.* 383. *R. Syn.* 259.
Ranunculus nemorosus albus simplex. *Park.* 325.
Anglis, common Wood Anemone.
Habitat in sylvis et ad sepes. 4. IV.

2. ANE.

2. ANEMONE feminibus acutis, foliolis incisfis, petalis apenninis. *lis lanceolatis numerosis. Sp. pl. 540.*
Anemone Geranii robertiani folio cœrulea. Baub. pin. 174.
Ranunculus nemorosus, flore purpureo-cœruleo. Park. 325. R. Syn. 259.
Ranunculus nemorosus, flore cœruleo, foliis majoribus apennini montis. Mentz. pug. t. 8. Tourn. 285.
Anglis, Mountain Wood Anemone.
Habitat in sylvis. In Wimbleton woods, Mr. Rand; near Harrow-on-the-hill, Mr. Dubois; in a wood near Luton-hoe, in Bedfordshire, Mr. T. Knowlton. R. Syn. 4. IV.
3. ANEMONE pedunculo involocrato, petalis rectis, *Pulsatilla* foliis bipinnatis. *Fl. jucc. 481. Sp. pl. 539.*
Pulsatilla folio crassiore et majore flore. Baub. pin. 177. R. Syn. 260.
Pulsatilla vulgaris. Park. 341. Ger. Em. 385.
Anglis, Pasque-flower.
Habitat in pratis montosis. On Gogmagog-hill's; on Bernack-heath, near Stamford, in great plenty; in pascuis montosis et siccis juxta Leadstone-hall, supra lacum loco the Close dicto abunde prope Pontemfractum, D. Richardson. R. Syn. in Cornbury park, near Charlbury, Oxfordshire. Blackf. spec. bot. 4. IV.

A D O N I S. 618.

Cal. pentaphyllus. Petala quinque plura absque nectario. Sem. nuda.

1. ADONIS floribus octopetalis, fructibus subcylindricis. *Hort. upf. 156. Sp. pl. 547.*
Adonis hortensis flore minore atro-rubente. Baub. pin. 178.
Flos Adonis. Park. par. 293. R. Syn. 251.
Flos adonis flore rubro. Ger. Em. 387.
Anglis, Adonis Flower, Pheasant's Eye, red Maithes, or red Morocco.
Habitat in arvis. In Cantio frequens; prope Acton, Dr. Watson. O. VI. VII.

RANUNCULUS. 619.

Cal. pentaphyllus. *Petala* quinque intra unques poro mellifero. *Sem.* nuda.

* *Foliis simplicibus.*

- × *Lingua.* 1. RANUNCULUS foliis lanceolatis, caule erecto. *Fl. suec.* 493. *Sp. pl.* 549.
 Ranunculus flammeus major. *Ger. Em.* 961. *R. Syn.* 250.
 Ranunculus palustris flammeus major. *Park.* 1215.
 Ranunculus longifolius palustris major. *Bauh. pin.* 180.
Anglis, great Spearwort.
Habitat in pascuis udis. In the bogs on Iwer-heath, et near Uxbridge. Blackf. spec. bot. Inter Rotherhithe Deptford copiose. 4. V.
- × *Flammula.* 2. RANUNCULUS foliis ovato-lanceolatis petiolatis, caule declinato. *Fl. suec.* 494. *Sp. pl.* 549.
 Ranunculus flammeus minor. *Ger. Em.* 961. *R. Syn.* 250.
 Ranunculus flammeus ferratus. *Ger. Em.* 961. *Park.* 1214.
 Ranunculus palustris flammeus minor five angustifolius. *Park.* 1214.
 Ranunculus longifolius palustris minor, item palustris ferratus. *Bauh. pin.* 180.
 β Ranunculus flammeus, latiori plantaginis folio; marginibus pilosis. *Pluk. alm.* 312.
Anglis, lesser Spearwort.
Habitat in paludosis ubique. 4. VI.—IX.

** *Foliis dissectis et divisis.*

- repens.* 3. RANUNCULUS calycibus patulis, pedunculis sulcatis, stolonibus repentibus, foliis compositis. *Fl. suec.* 505. *Sp. pl.* 554.
 Ranunculus foliis ternatis, foliolis petiolatis trifidis incis, caule productiore multifloro *Hort. cliff.* 230.
 Ranunculus pratensis repens. *Park.* 329. *R. Syn.* 247.
 hirsutus. *Bauh. pin.* 179.

Ranunculus

- Ranunculus pratensis. *Ger. Em.* 951.
- β Ranunculus repens major. *Pet. h. br. t.* 38. *f.* 8.
Anglis, creeping Crowfoot, or Butter-cups.
Habitat in pratis et pascuis. 4. V.
4. RANUNCULUS calycibus retroflexis, pedunculis *bulbosus*. ×
 sulcatis caule erecto, foliis compositis. *Fl. suec.*
 504. *Sp. pl.* 554.
 Ranunculus bulbosus. *Ger. Em.* 953. *Park.* 329.
R. Syn. 247.
 Ranunculus pratensis radice verticilli modo rotunda.
Bauh pin. 179.
 Ranunculus tuberosus. *Dod. pempt.* 431.
 β Ranunculus rectus foliis pallidioribus hirsutus. *Bauh.*
hist. III. 417. *R. Syn.* 247.
Anglis, bulbose Crowfoot, or Butter-cups.
Habitat in pratis et pascuis. 4. V.
Pharmac. Ranunculi bulbosi *Herba*.
5. RANUNCULUS calycibus patulis pedunculis tereti- *acris*.
 bus, foliis tripartito multifidis: fummis linearibus.
Fl. suec. 507. *Sp. pl.* 554.
 Ranunculus foliis peltatis quinquangularibus, multi-
 partitis laciniis linearibus. *Hort. cliff.* 231. *It. oel.*
 104.
 Ranunculus pratensis erectus acris. *Bauh. pin.* 178.
R. Syn. 248. *vulgaris.* *Park.* 329.
 Ranunculus surrectis cauliculis. *Ger. Em.* 951.
Anglis, upright Meadow Crowfoot.
Habitat in pratis et pascuis ubique. 4. VI. VII.
6. RANUNCULUS foliis radicalibus reniformibus cre- *auricomus*.
 natis incis; caulinis digitatis linearibus, caule
 multifloro. *Hort. cliff.* 229. *Fl. suec.* 498. *Sp. pl.*
 551.
 Ranunculus auricomus. *Ger. Em.* 954.
 Ranunculus nemorosus dulcis, II. Tragi. *Park.* 326.
R. Syn. 248.
 Ranunculus nemorosus vel sylvaticus folio rotundo.
Bauh. pin. 178.
Anglis, sweet Wood Crowfoot, or Goldilocks.
Habitat in sylvis et ad sepes frequens. 4. IV.

sceleratus. 7. RANUNCULUS foliis inferioribus palmatis, summis digitatis, fructibus oblongis. *Hort. cliff.* 230. *Fl. suec.* 499 *Sp. pl.* 551.

Ranunculus palustris rotundifolius. *Ger. Em.* 962. *R. Syn.* 249.

Ranunculus palustris fardonius lævis. *Park* 1215.

Ranunculus palustris apii folio lævis. *Bauh. pin.* 180.

Anglis, round-leaved Water Crowfoot.

Habitat in aquosis ubique. ☉. V. VI.

Pharmac. Ranunculi palustris *Herba*.

parviflorus. 8. RANUNCULUS feminibus muricatis, foliis simplicibus laciniatis acutis hirsutis, caule prostrato. *Syst. nat* 1087.

Ranunculus hirsutus annuus flore minimo. *R. Syn. ed. II.* 135. *Pluk. ph. t.* 55. *f.* 1. *Pet. b. br. t.* 38. *f.* 9. *R. Syn.* 248 *t.* 12. *f.* 1.

Ranunculus arvensis annuus hirsutus, florum omnium minimo *Hist. ox. II.* 440. *f.* 4. *t.* 28. *f.* 21.

Ranunculus annuus foliis Geranii columbini. *Merr. pin.*

Anglis, small-flowered Crowfoot.

Habitat in pratis, et arvis solo glareoso passim. ☉. V.

arvensis. 9. RANUNCULUS feminibus aculeatis, foliis superioribus decompositis linearibus. *Sp. pl.* 555. *Fl. suec.* 508.

Ranunculus arvorum *Park.* 328. *Ger. Em.* 951. *R. Syn.* 249.

Ranunculus arvensis echinatus. *Bauh. pin.* 179. *hist. III.* 859.

Anglis, Corn Crowfoot.

Habitat in arvis. ☉. VI.

hederaceus. 10. RANUNCULUS foliis subrotundis trilobis integerrimis, caule repente *Hort. cliff.* 231 *Roy. lugdb.* 492. *Dalib. paris.* 167. *Gort. gelr.* 325. *Sp. pl.* 556.

Ranunculus aquatilis hederaceus albus. *R. Syn.* 249.

Ranunculus aquaticus hederaceus luteus. *Bauh. pin.* 180.

Ra-

Ranunculus hederaceus rivulorum se extendens, atra macula notatus. *Bauh. hist. III. 782.*

Ranunculus hederaceus aquaticus. *Park. 1216.*

Ranunculi aquatilis varietas altera. *Ger. Em. 830.*

Anglis, Ivy-leaved Water Crowfoot.

Habitat in fossis et rivulis lentè fluentibus passim. *Æ. V.*

II. RANUNCULUS foliis submersis capillaceis, emer. *aquatilis.* sis peltatis. *Sp. pl. 556.*

Ranunculus aquatilis. *Ger. Em. 829. R. Syn. 249.*

Ranunculus aquaticus, hepaticæ facie. *Park. 1216.*

Ranunculus aquaticus, folio rotundo et capillaceo.

Bauh. pin. 180.

β Ranunculus foliis capillaceis, circumscriptione vaga brevioribus. *Hall. helv. 328.*

Ranunculus aquaticus capillaceus. *Bauh. pin. 180.*

Ranunculus aquaticus omnino tenuifolius. *Bauh. hist. III. 781.*

γ Ranunculus foliis omnibus capillaceis, circumscriptione oblongis. *Roy lugdb. 492.*

Millefolium aquaticum, foliis foeniculi, Ranunculi flore et capitulo. *Bauh. pin. 141.*

δ Ranunculus foliis omnibus capillaceis, circumscriptione rotundis. *Hort. cliff. 231. Roy lugdb. 492.*

Ranunculus aquaticus albus, circinatis tenuissime divisis foliis, floribus ex alis longis pediculis innixis.

Pluk. ph. t. 55. f. 2.

Anglis, Various leaved Water Crowfoot.

Habitat α. β. δ. in fossis et stagnis. γ. in rivulis. *Æ. IV. V.*

F I C A R I A.

CAL. *Perianthium* triphyllum: *foliolis* ovatis, concavis, deciduis.

COR. *Petala* circiter octo lanceolata nitida. *Nectarium* fovea in singulo petalo supra unguem squamula emarginata clausum.

STAM. *Filamenta* plurima brevissima. *Antheræ* rectæ, oblongæ, didymæ.

PIST. *Germina* numerosa, in capitulum collecta. *Styli* nulli. *Stigmata* reflexa minima.

PER. nullum. *Receptaculum* pedunculis minutissimis femina adnecit.

SEM. plurima, oblonga, apice reflexo.

verna.

I. FICARIA. *Dill. cat. giff.* 39.

Ranunculus foliis cordatis angulatis petiolatis. *Fl. suec.* 296. *Sp. pl.* 550.

Chelidonium minus. *Ger. Em.* 816. *Park.* 617. *R. Syn.* 246.

Chelidonia rotundifolia minor. *Bauh. pin.* 309.

Anglis, Pilewort, or lesser Celandine.

Habitat in pratis et pascuis ubique. 4. IV.

Pharmac. Chelidonii minoris. *Radix, Herba.*

TROLLIUS. 620.

Cal. o. *Petala* circiter quatuordecim. *Caps.* plurimæ, ovatæ, monospermæ.

europæus. I. TROLLIUS corollis conniventibus, nectariis longitudine staminum. *Fl. suec.* 510. *Sp. pl.* 556.

Helleborus flore clauso erecto petiolato, caule simplicissimo. *Fl. suec. ed. I.* 474.

Pseudo-Helleborus ranunculoides, flore globofo. *Hist. ox.* III. 467. *f.* 12. *t.* 2. *f.* 2.

Ranunculus globosus. *Ger. Em.* 955. *Park. par.* 218. *R. Syn.* 272.

Ranunculus flore globofo, quibusdam Trollius flos. *Bauh. hist. III.* 419.

Ranunculus montanus, aconiti folio, flore globofo. *Bauh. pin.* 182.

Anglis, the Globe-flower, or Locker-gowlons.

Habitat ad latera montium, inque pratis montosis in septentrionalibus Angliæ et Cambriæ copiosissime. 4.

V. VI.

CALTHA. 623.

Cal. o. *Petala* quinque. *Nectararia* o. *Caps.* plures, polyspermæ.

palustris. I. CALTHA. *Fl. lapp.* 227. *suec.* 511. *Sp. pl.* 558.

Caltha palustris. *Bauh. hist. III.* 419. flore simplici. *pin.* 276.

Caltha palustris vulgaris simplex. *Park.* 1213.

Caltha

Caltha palustris major. Ger. Em. 817.

Populago. Tab. ic. 750. R. Syn. 272.

Anglis, Marsh Marigold.

Habitat in aquis et ad ripas fluviorum. 4. IV.

HELLEBORUS. 622.

Cal. nullus. *Petala* quinque seu plura. *Nectaria* bilabiata tubulata. *Caps.* polyspermæ, erectiusculæ.

1. HELLEBORUS caule multifloro folioso, foliis pedatis. *Sp. pl.* 558.

Helleborus caule inferne angustato multifolio multifloro, foliis caule brevioribus. *Hort. cliff.* 227. *Roy. lugdb.* 484.

Helleborus niger foetidus. *Bauh. pin.* 185.

Helleboraster maximus. Ger. Em. 976. R. Syn. 271. f. confiligo. *Park.* 212.

Anglis, great bastard black Hellebore, Bearsfoot, or Setterwort.

Habitat in pratis pascuis et ad sepes. Near Cherry-Hinton, Cambridgeshire; in the parish of Brundish in Suffolk, Mr. J. Sherard; upon the downs in Sussex, towards Chichester, D. Dillenius. R. Syn. 4. III.

2. HELLEBORUS caule multifloro folioso, foliis digittatis. *Sp. pl.* 558.

Helleborus caule æquali folioso, foliis radicalibus caulem tandem superantibus. *Hort. cliff.* 227. *ups.* 158. *Roy. lugdb.* 484.

Helleborus niger hortensis flore viridi. *Bauh. pin.* 185. R. Syn. 271.

Helleboraster minor, flore viridante. *Park.* 212.

Helleborastrum. Ger. Em. 976.

Anglis, wild black Hellebore.

Habitat in pratis montosis. In Bigwyn closes near Cambridge; about Arundel-castle in Sussex; in Sir Tho. D'Aeth's woods at Knowlton, in East Kent; in the woods near Stoken-church, Oxfordshire. R. Syn. circa Ditton et Whitwell, in comitatu Cantabrigienfi, D. Lyons. 4. IV.

THALICTRUM. 617.

Cal. o. Petala quinque. Semina nuda.

- flavum.* 1. THALICTRUM caule folioso sulcato, panicula multiplici erecto. *Fl. suec.* 488 *Sp. pl.* 546.
 Thalictrum majus vulgare. *Park.* 263.
 Thalictrum seu Thalitrum majus. *Ger. Em.* 1251. *R. Syn.* 203.
 Thalictrum majus filiqua angulosa aut striata. *Bauh. pin.* 336.
Anglis, Meadow Rue.
Habitat in pratis et pascuis humidis et ad ripas fluviorum. 4. VI.
- minus.* 2. THALICTRUM foliis sexpartitis, floribus cernuis. *Sp. pl.* 546. *Fl. suec.* 489.
 Thalictrum caule folioso æquali, foliolis caulinis acutis, panicula simplici, floribus nutantibus. *It. goth.* 180.
 Thalictrum minus. *Ger. Em.* 1251. *Park.* 264. *Bauh. pin.* 337. *R. Syn.* 203.
 β Thalictrum montanum minus, foliis latioribus. *R. Syn.* 204.
Anglis, the lesser Meadow-rue.
Habitat in pratis montosis humidiusculis. On the chalky grounds about Newmarket, Linton, and elsewhere in Cambridgeshire; in comitatibus Eboracensi, Westmorlandico, Cumberlandico, frequens. 4. VII. VIII.
- alpinum.* 3. THALICTRUM caule simplicissimo subnudo racemo-simplici laxo terminali. *Fl. suec.* 491. *Sp. pl.* 545.
 Thalictrum minimum montanum atro-rubens foliis splendentibus. *R. Syn.* 204. *Boerb. ind. alt.* 44. *ic.*
 Thalictrum montanum minimum præcox, foliis splendentibus. *Hist. ex. III.* 325. *f. 9. t. 20. f. 14.*
Anglis, Mountain Meadow-rue.
Habitat in udis scopulis et ad rivulos in alpibus in Arvoniam planta est vulgatissima, D. Lhwyd. R. Syn. 4. VI.

CLE.

C L E M A T I S. 616.

Cal. o. *Petala* quatuor. *Sem.* caudata.

1. CLEMATIS foliis pinnatis : foliolis cordatis scanden- *Vitalba*.
 tibus. *Hort. cliff.* 225. *Roy. lugdb.* 486. *Hall helv.*
 334. *Gron. virg.* 62. *Gort. gelr.* 315. *Sp. pl.* 544.
Clematis sylvestris latifolia. *Bauh. pin.* 300. f. *Vi-*
orna. Park. 380.
Clematis latifolia seu Atragene quibusdam. *Bauh. hist.*
II. 125. *R. Syn.* 258.
Viorna. Ger. Em. 736.
Anglis, great wild Climber, or Travellers Joy.
Habitat in sepibus. h. VII.

Classis

*Classis XIV.*DIDYNAMIA
GYMNOSPERMIA.* *Calyces subquinquefidi.*

TEUCRIUM. 625.

Corollæ labium superius (nullum) ultra basin bipartitum, divaricatum ubi stamina.*Scordium.* I. TEUCRIUM foliis oblongis sessilibus dentato-ferratis, floribus geminis axillaribus, caule diffuso. *Sp. pl.* 565. *Fl. suec.* 513.Teucrium foliis ovato-lanceolatis ferrulatis sessilibus, floribus sepius binatis. *Hort. cliff.* 302. *Mat. med.* 289.*Scordium.* *Baub. pin.* 247. *hist. III.* 292. *R. Syn.* 246.*Scordium majus et minus.* *Ger. Em.* 661.*Scordium legitimum.* *Park.* 111.*Anglis,* Water Germander.Habitat in palustribus Eliensis insulæ copiose. *R. Syn.* 24. VIII.*Pharmac.* Scordii Herba.*Chamæ-* *2. TEUCRIUM foliis ovatis inciso-crenatis petiolatis, *drys.* floribus subverticillatis ternis, caulibus procumbentibus. *Syst. nat.* 1095. *Sp. pl.* 565.Chamædryes minor repens. *Baub. pin.* 248.Chamædryes major latifolia. *Ger. Em.* 616.*Anglis,* Germander.Habitat on the ruins of Winchelsea castle. *R. Syn.* 231. 24. V. VI.*Pharmac.* Chamædryos Herba.

3. TEU-

3. TEUCRIUM foliis trifidis linearibus integerrimis, *Chamæ-*
floribus sessilibus lateralibus solitariis, caule diffuso. pitys.
Sp. pl. 562.

Chamæpitys vulgaris. Park. 283. R. Syn. 244.

Chamæpitys lutea vulgaris f. folio trifido. Bauh. pin.
249.

Chamæpitys mas. Ger. Em. 525.

Anglis, Ground-pine.

Habitat in agris restibilibus. On the lays about the bor-
ders of Triplow-heath in Cambridgeshire; about
Rocheſter, Dartford, &c. R. Syn. 244. VI. VII.

Pharmac. Chamæpityos Herba.

4. TEUCRIUM foliis cordatis ferratis petiolatis, race- *Scorodo-*
mis lateralibus secundis, caule erecto. Sp. pl. 564. nia.

Scorodonia f. Salvia agrestis. Ger. Em. 626. R. Syn.
245.

Scorodonia f. Scordium alterum quibusdam et Salvia
agrestis. Park. 111.

Scordium alterum seu Salvia agrestis. Bauh. pin. 247.
Anglis, Wood Sage.

Habitat in sylvis dumetis et ericetis frequens. 24. VII.
Pharmac. Scorodonix Herba.

A J U G A. 624.

Corollæ labium superius minimum. Stamina labio su-
periore longiora.

1. AJUGA stolonibus reptantibus. *Sp. pl. 561. reptans.*

Ajuga foliis obovatis crenatis, caule simplicissimo,
stolonibus reptatricibus. Gort. gelr. 330.

Bugula. Dod. p. 135. Ger. Em. 631. R. Syn. 245.

Bugula vulgaris. Park. 525.

Consolida media pratensis cœrulea. Bauh. pin. 260.
Anglis, Bugle.

Habitat in pratis et pascuis humidis. 24. V.

Pharmac. Bugulæ f. consolidæ mediæ Herba.

2. AJUGA tetragono pyramidalis. *Fl. suec. 512. Sp. pyramida-*
pl. 561. lis.

Bugula

DIDYNAMIA GYMNOSPERMIA.

Bugula cœrulea alpina. *Park.* 525. *Pluk. ph. t.* 18.
f. 3. *R. Syn.* 245

Consolida media cœrulea alpina. *Bauh. pin.* 260.
Anglis, Mountain Bugle, or Sicklewort.

Habitat on Carnedd Llewellyn in Carnarvonshire, Dr.
Johnfon. R. Syn. 4. VI.

N E P E T A. 629.

Corollæ labium inferius lacinula intermedia crenata :
faux margine reflexo. *Stamina* aproximata.

Cataria. 1. *NĒPETA* floribus spicatis, verticillis subpedicellatis,
foliis petiolatis cordatis dentato-ferratis. *Sp. pl.* 570.
Fl. suec. 514.

Nepeta major vulgaris. *Park.* 38. *R. Syn.* 237.

Mentha cataria vulgaris et major. *Bauh. pin.* 228.

Mentha felina f. cataria. *Ger. Em.* 682.

Anglis, Nap, or Cat-mint.

Habitat in sepibus. 4. VII.

Pharmac. *Nepetæ Herba.*

B E T O N I C A. 631.

Cal. aristatus. *Corollæ* labium superius adscendens
planiusculum ; tubus cylindricus.

officinalis. 1. *BETONICA* spica interrupta, corollarum lacinia la-
bii intermedia emarginata. *Roy. lugdb.* 316. *Sp. pl.*
573. *Fl. suec.* 515.

Betonica. *Ger. Em.* 714. *R. Syn.* 238. purpurea.
Bauh pin. 235.

Betonica vulgaris flore purpureo. *Park.* 614.

Anglis, Wood Betony.

Habitat in ericetis et sylvis. 4. VII. VIII.

Pharmac. *Betonicæ Herba, Flores.*

hirta. 2. *BETONICA* verticillata villosa, calycibus subspino-
fis. *Syst. nat.* 1097.

Betonica foliis inter verticillos diffitos brevibus in a-
ristam terminatis integris. *Hall. helv.* 646.

Sideritis arvensis latifolia hirsuta lutea. *R. Syn.* 242.

Sideritis lutea. *Pet. h. br. t.* 33. f. 10.

Anglis, yellow Betony.

Habitat

Habitat *in arvis. In the west riding of Yorkshire, about Wakefield, Darfield, Sheffield, &c. among the corn plentifully. R. Syn. ☉. VIII.*

M E N T H A. 633.

Corolla subæqualis, quadrifida : lacinia latiore emarginata. Stamina erecta, distantia.

1. MENTHA spicis solitariis interruptis, foliis lanceolatis ferratis sessilibus. *Sp. pl. 576. Hort. cliff. 306. spicata.*

Mentha angustifolia spicata. Bauh. pin. 227.

Mentha angustifolia spicata, glabra, folio rugosiore, odore graviore. R. Syn. 233.

- β *Mentha spicata angustifolia, glabra, spica latiore. D. Dale. R. Syn. 233.*

- γ *Mentha spicata glabra, latiore folio. D. Dale. R. Syn. 234.*

Anglis, Spear-mint.

Habitat ad ripas fluviorum Found by Mr. Dale, in Essex, by Bocking river side, below the fulling-mill; on the river Medway in Kent, near Maidstone, D. Plukenet. R. Syn. 4. VIII.

Pharm. Menthæ vulgaris Herba.

2. MENTHA floribus spicatis confertis, foliis oblongis longifolia. ferratis tomentosis sessilibus.

Mentha sylvestris folio longiore. Bauh. pin. 227.

Menthastrum spicatum folio longiore candicante.

Bauh. hist. III. 2. 221. R. Syn. 234.

- β *Menthastrum aquaticum genus hirsutum spica latiore. Bauh. hist. III. 2. 222. R. Syn. 234.*

Anglis, long-leaved Horse-mint.

Habitat in palustribus et aquosis. 4. VIII.

Pharmac. Menthastrum Herba.

3. MENTHA floribus spicatis, foliis rotundis crenatis rotundifolia. rugosis sessilibus.

Mentha sylvestris rotundiore folio. Bauh. pin. 227.

Menthastrum folio rugoso rotundiore spontaneum, flore spicato, odore gravi. Bauh. hist. III. 2. 219.

R. Syn. 234.

Menthastrum. Ger. Em. 683.

Anglis,

Anglis, round-leaved Horse-mint.

Habitat in aquosis sed rarius. In Hornsey and Harefield church yards. Blackf. spec. bot. 4. VIII.

piperita. 4. MENTHA floribus capitatis solitariis terminalibus, foliis ovato-oblongis ferratis, staminibus corolla brevioribus. Syst. nat. 1099.

Mentha floribus capitatis, foliis lanceolatis ferratis subpetiolatis. Sp. pl. 576.

Mentha spicis brevioribus et habitioribus, foliis Menthæ fuscæ, sapore fervido piperis. R. Syn. 234. t. 10. f. 2.

Mentha piperata acuta. Pet. h. br. t. 31. f. 10.

β Mentha piperata obtusa. Pet. h. br. t. 31. f. 11.

Anglis, Pepper-mint.

Habitat in aquosis et rivulos; inter Newington et Hornsey, D. Alchorne. 4. VIII.

Pharmac. Menthæ piperitæ Herba.

gentilis. 5. MENTHA floribus verticillatis, foliis ovatis acutis ferratis staminibus corolla brevioribus. Sp. pl. 577.

Mentha fusca f. vulgaris. Park. 31. R. Syn. 232.

Mentha cardiaca. Ger. Em. 680.

Mentha hortensis verticillata, ocygni odore. Bauh. pin. 227.

β Mentha arvensis verticillata folio rotundiore, odore aromatico. R. Syn. 239.

Anglis, Red Mint.

Habitat in aquosis. 4. VIII.

verticillata. 6. MENTHA floribus verticillatis, pedunculis pilosis, foliis ovatis ferratis, staminibus corolla longioribus. Syst. nat. 1099.

Mentha verticillata. Riv. irr. mon. ic. R. Syn. 232.

Mentha crispa. Park. 31. verticillata. Bauh. pin. 227.

Mentha fativa rubra. Ger. Em. 680.

Anglis, Curled Mint.

Habitat in palustribus et ad ripas fluviorum. In Peckham fields, and on Hackney river, near the ferry-house. R. Syn. 4. VIII.

7. MENTHA floribus verticillatis, foliis ovatis acutis *arvensis*.
 ferratis staminibus corollam æquantibus. *Hort. cliff.*
 307. *Sp. pl.* 577. *Fl. suec.* 516. *Dalib. parif.* 178.
 Mentha arvensis verticillata. *Baub. pin.* 229.
 Mentha f. Calamintha aquatica. *Ger. Em.* 684. *R.*
Syn. 232.
Anglis, Corn Mint.
 Habitat in arvis humidis. 4. VIII. IX.
8. MENTHA floribus verticillatis, foliis lanceolato-o-*exigua*.
 vatis glabris acutis integerrimis. *Cent. pl.* 159. *Syst.*
nat. 1099.
 Mentha aquatica exigua. *Trag. lib. i. c. 6. ic.* *R.*
Syn. 232.
 Mentha arvensis verticillata f. aquatica Belgarum Lo-
 belii. *Park.* 36.
Anglis, smooth Mint.
 Habitat ad rivulos. In the hop-ground at Bocking
 plentifully, Mr. Dale ; by the side of the new-river,
 near the upper end of Stoke-Newington, Mr. Buddle.
R. Syn. 4. VIII.
9. MENTHA floribus capitatis, foliis ovatis ferratis pe-*aquatica*.
 tiolatis, staminibus corolla longioribus. *Sp. pl.* 576.
 Mentha aquatica f. Sifymbrium. *Ger. Em.* 684. *Baub.*
hist. III. 223. *R. Syn.* 233.
 Mentha aquatica rubra. *Park.* 1243.
 Mentha rotundifolia palustris f. aquatica major. *Baub.*
pin. 227.
- β Mentha Sifymbrium dicta hirsuta glomerulis ac foliis
 minoribus ac rotundioribus. *R. Syn.* 233 t. 10.
 f. 1.
- γ Sifymbrium hirsutum. *R. Syn.* 233.
Anglis, Water Mint.
 Habitat in aquosis et ad ripas fluviorum. 4. VII.
 VIII.
- Pharmac.* Menthæ aquaticæ f. Sifymbrii Herba.
10. Mentha floribus capitatis globosis, foliis ovatis fer-*hirsuta*.
 ratis, staminibus corolla brevioribus.
Anglis, round-headed Mint.

Habitat

Habitat in aquosis prope Marybone copiose; et inter Newington et Hornsey. 4. VIII.

- Pulegium.* 11. MENTHA floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus. *Hort. cliff.* 307. *Sp. pl.* 577. *Dalib. parif.* 178. *Gort. gelr.* 344.
Pulegium. *Bauh. hist.* III. 256. *R. Syn.* 235.
Pulegium vulgare. *Park* 29.
Pulegium regium. *Ger. Em.* 671.
Pulegium latifolium. *Bauh. pin.* 222.
Anglis, Pennyroyal.
Habitat in inundatis. 4. VIII. IX.
Pharmac. *Pulegii Herba.*

GLECHOMA. 634.

Antherarum singulum par in formam crucis connivens.
Cal. quinquefidus.

- hederacea.* 1. GLECHOMA foliis reniformibus crenatis. *Hort. cliff.* 307. *Fl. suec.* 518. *Sp. pl.* 578.
Hedera terrestris. *Ger. Em.* 856. vulgaris. *Park.* 676. *Bauh. pin.* 306.
Calamintha humilior, folio rotundiore. *Tourn.* 197. *R. Syn.* 243.
Hedera terrestris montana. *Bauh. pin.* 306. *Park.* 677.
Anglis, Ground-Ivy.
Habitat in umbrosis et ad sepes. 4. V.
Pharmac. *Hederæ terrestris Herba.*

- arvensis.* 2. GLECHOMA foliis cordatis oblongis crenatis, caule hispidulo erecto. *Sp. pl.* 578.
Sideritis calycibus subinermibus hispidis. *Fl. suec.* 488.
Sideritis humilis lato obtuso folio. *Ger. Em.* 699. *R. Syn.* 242.
Sideritis Hederulæ folio. *Park.* 587.
Sideritis alfinis trixaginis folio. *Bauh. pin.* 233.
Anglis, upright Ground-Ivy.
Habitat in arvis. ☉. VIII.

L A-

arvensis. 2. *Glechoma* *Sc.*
Stachys arvensis. *Lin.*

L A M I U M. 636.

Corollæ labium superius integrum, fornicatum; labium inferius bilobum; faux utrinque margine dentata.

1. LAMIUM foliis cordatis acuminatis ferratis petiolaribus. *Hort. cliff.* 314. *Sp. pl.* 579. *Fl. suec.* 520.

Lamium album. *Ger. Em.* 782. *R. Syn.* 240.

Lamium album non foetens, folio oblongo. *B. pin.* 231.

Lamium vulgare album f. Archangelicum flore albo. *Park.* 604.

Anglis, white dead Nettle, or Archangel.

Habitat ad sepes, et in cultis, ruderatis. 4. V.

Pharmac. *Lamii albi Herba, Flores.*

2. LAMIUM foliis cordatis obtusis petiolatis. *Hort. rubrum, cliff.* 314. *Fl. suec.* 521. *Sp. pl.* 579.

Lamium rubrum. *Ger. Em.* 703. *R. Syn.* 240.

Lamium vulgare folio subrotundo, flore rubro. *Park.* 604.

Lamium purpureum foetidum, folio subrotundo f. Galeopsis dioscoridis. *Bauh. pin.* 230.

- β *Lamium rubrum minus, foliis profunde incis.* *R. Syn.* 240.

Anglis, red Archangel, or dead Nettle.

Habitat in ruderatis et arvis. 5. V.

Pharmac. *Lamii rubri Folia, Flores.*

3. LAMIUM foliis floribus sessilibus amplexicaulibus amplexi-obtusis. *Hort. cliff.* 314. *Sp. pl.* 579. *Fl. suec.* 522. *caule.*

Lamium folio caulem ambiente majus et minus.

Bauh. pin. 231. *R. Syn.* 240.

Alfine hederula altera. *Ger. Em.* 616.

Alfine hederulæ folio major. *Park.* 762.

Anglis, great Henbit.

Habitat in arvis. 6. VI.

G A L E O P S I S. 637.

Corollæ labium superius crenatum, fornicatum; labium inferius supra bidentatum.

1. GALEOPSIS internodiis caulinis æqualibus: verti *Ladanum.* cillis omnibus remotis. *Sp. pl.* 579. *Fl. suec.* 524.

Q

Sideritis

DIDYNAMIA GYMNOSPERMIA.

Sideritis arvensis rubra. *Park.* 587. *R. Syn.* 242.

Sideritis VII. Ger Em. 699.

Sideritis arvensis angustifolia rubra. *Bauh pin.* 233.
Anglis, narrow-leaved Allheal.

Habitat in arvis. ☉. VII. VIII.

Tetrabit. 2. *GALEOPSIS* internodiis caulinis superne incrassatis, verticillis summis subcontiguis. *Sp. pl.* 579. *Fl. suec.* 523.

Lamium cannabino folio vulgare. *R. Syn.* 240.

Cannabis spuria. *Ger. Em.* 709. *Park.* 599.

Urtica aculeata, foliis ferratis. *Bauh. pin.* 232.

β *Lamium cannabinum floribus albis, verticillis purpurascens.* *R. Syn.* 241.

γ *Cannabis spuria flore albo magno, eleganti.* *Merr. pin.*

♫ *Lamium cannabino folio, flore amplo luteo: labio purpureo.* *R. Syn.* 241.

Anglis, Nettle-hemp, or Hemp-leaved dead Nettle.

Habitat in arvis. & inter segetes prope Kighley, et per totum Cravoniæ tractum; Dr. Richardson. *R. Syn.*

☉. VIII.

Galeobdolon. 3. *GALEOPSIS* verticillis sexfloris: involucri tetraphyllo. *Sp. pl.* 580. *Fl. suec.* 525.

Leonurus foliis ovatis ferratis acutis. *Hort. cliff.* 313.

Lamium luteum. *Ger. Em.* 671. *Park.* 606. *R. Syn.* 240.

Lamium folio oblongo, luteum. *Bauh. pin.* 231.

Anglis, yellow Nettle-hemp.

Habitat in sylvis sepibus et umbrosis. 4. V.

S T A C H Y S. 638.

Corollæ labium superius fornicatum; labium inferius lateralibus reflexum: intermedia majore emarginata. *Stamina* deflorata versus latera reflexa.

germanica. 1. *STACHYS* verticillis multifloris, foliorum ferraturis imbricatis caule lanato. *Sp. pl.* 581.

Stachys Fuchsi. *Bauh. hist III.* 319. *Ger. Em.* 795. *R. Syn.* 239.

Stachys major germanica. *Bauh. pin.* 236.

Anglis, bafe Horehound.

Habitat

arvensis. vide Glechoma arvensis.

Habitat in *sepibus*. Nigh Witney park in Oxfordshire, and thereabout plentifully. *R. Syn.* Found by Dr. Richardson four miles south of Grantham, near the London road opposite Easton, the seat of H. Chumley, Esq; 4. VII.

2. STACHYS verticillis sexfloris, foliis cordatis petio- *sylvatica*.
latis. *Sp. pl.* 580. *Fl. suec.* 526.

Galeopsis legitima Dioscoridis. *Park.* 608. *R. Syn.* 237.

Galeopsis vera. *Ger. Em.* 709.

Lamium maximum sylvaticum foetidum. *Bauh pin.* 231.

- β Lamium sylvaticum spicatum foetidum, folio angulosa
minus. *Pluk. alm.* 128.

Anglis, hedge Nettle.

Habitat in *sylvis et sepibus*. VII. VIII.

3. STACHYS verticillis sexfloris, foliis lineari-lanceo- *palustris*.
latis semiamplexicaulibus. *Sp. pl.* 580. *Fl. suec.* 528.

Stachys palustris foetida. *Bauh. pin.* 236.

Sideritis anglica strumosa radice. *Park.* 587. *R. Syn.* 242.

Panax coloni. *Ger. Em.* 1005.

Anglis, Clown's Allheal.

Habitat in *aquosis, et ad ripas fluviorum*. 4. VIII.

B A L L O T A. 639.

Cal. hypocrateriformis, quinquedentatus, decemfria-
tus. *Corollæ* labium superius crenatum concavum.

1. BALLOTA foliis cordatis indivisis ferratis, calycibus *nigra*.
acuminatis. *Sp. pl.* 582. *Fl. suec.* 529.

Ballota *Matth.* 825. *R. Syn.* 244.

Marrubium nigrum. *Ger. Em.* 701.

Marrubium nigrum foetidum, Ballotte dictum. *Park.* 1230.

Marrubium nigrum foetidum, Ballotte Dioscoridis.
Bauh. pin. 230.

- β Ballota foliis cordatis indivisis ferratis, calycibus sub-
truncatis. *Fl. suec.* 530.

Ballota flore albo. *Tourn.* 185.

Anglis, stinking Horehound.

Habitat in *runderatis*. ☉. VII.

Pharmac. Marrubii nigri *Herba*.

MARRUBIUM. 640.

Cal. hypocrateriformis, rigidus, decemstriatus. *Corollæ* labium superius bifidum, lineare, rectum.

vulgare. I. MARRUBIUM dentibus calycinis setaceis uncinatis.

Sp. pl. 585. *Mat. med.* 305. *Hall. helv.* 648.

Marrubium album. *Bauh. hist.* III. 316. *Ger. Em.* 693. *R. Syn.* 239.

Marrubium album vulgare. *Bauh. pin.* 230. *Park.* 44.

Anglis, white Horehound.

Habitat in *runderatis et ad vias*; in *ericeto* Hampstediano. ♀. VII.

Pharmac. Marrubii albi. f. *Præssii Herba*.

LEONURUS. 641.

Antheræ glandulis nitidis adspersæ.

Cardiaca. I. LEONURUS foliis caulinis lanceolatis trilobis. *Hort.*

cliff. 313. *Fl. suec.* 532. *Sp. pl.* 584.

Cardiaca. *Bauh. hist.* III. 320. *Park.* 41. *R. Syn.* 239.

Marrubium cardiaca dictum. *Bauh. pin.* 230.

Anglis, Mother wort.

Habitat in *finetis et ruderatis passim*. ♂. VII.

Pharmac. Cardiacæ *Herba*.

** *Calycis bilabiati.*

CLINOPODIUM. 644.

Involucrum multifetum, verticello subjectum.

vulgare. I. CLINOPODIUM capitulis subrotundis hispidis, bracteis setaceis. *Sp. pl.* 587.

Clinopodium organo simile. *Bauh. pin.* 224. *R. Syn.* 239.

Acinos. *Ger. Em.* 675. f. *clinopodium majus*. *Park.* 22.

Anglis, great wild Basil.

Habitat *ad sepes et in pratis aridis*. ♀. VII.

O R I-

ORIGANUM. 645.

Strobilus tetragonus, spicatus, calyces colligens.

1. ORIGANUM spiciis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. *Sp. pl. vulgare.*

590. *Fl. suec.* 534.

Origanum vulgare spontaneum. Bauh. hist. III. 236.

R. Syn. 236.

Origanum anglicum. Ger. Em. 666.

Origanum sylvestre. Bauh. pin. 223.

Majorana sylvestris. Park. 12.

Anglis, wild Marjoram.

Habitat ad sepes et in dumetis. 4. VII.

Pharmac. Origani vulgaris Herba.

2. ORIGANUM spicis oblongis aggregatis hirsutis, foliis cordatis tomentosis. *Sp. pl. 590. onites.*

Origanum onites. Bauh. pin. 223. *R. Syn.* 236.

Majorana major anglica. Ger. Em. 664.

Majorana latifolia, f. major anglica. Park 12.

Anglis, Pot-marjoram.

Habitat in pratis aridis. On the left hand of the road

from Braintree to Raine, beyond the bridge. Mr.

Dale. R. Syn. 4. VIII.

THYMUS. 646.

Calycis bilabiate faux villis clausa.

1. THYMUS floribus capitatis caulibus repentibus foliis planis obtusis basi ciliatis. *Mat. med. 282. Sp. serpyllum.*

pl. 590. Fl. suec. 535.

Serpyllum vulgare. Ger. Em. 570. *Bauh. hist. III.*

269. R. Syn. 230.

Serpyllum vulgare minus. Bauh. pin. 220. *Park.* 8.

β *Serpyllum vulgare, flore albo. Tourn.* 197.

γ *Serpyllum vulgare, flore amplo. R. Syn.* 230.

δ *Serpyllum vulgare majus. Bauh. pin.* 220.

ε *Serpyllum citratum. Ger. Em.* 575.

ζ *Serpyllum angustifolium glabrum. Bauh. pin.* 220.

η *Serpyllum vulgare hirsutum. R. Syn.* 231.

θ *Serpyllum villosum fruticosius floribus dilute rubenti-*

bus. R, Syn. 231.

DIDYNAMIA GYMNOSPERMIA.

Serpyllum hirsutum, minus, repens, inodorum. *Pluk. alm.* 344.

Anglis, Mother of Thyme, *ε* Lemon Thyme, *η* Hoary Thyme.

Habitat *in montosis ubique*. 4. VII. VIII.

Pharmac. Serpylli *Herba*.

Acinos. 2. THYMUS floribus verticillatis, pedunculis unifloris, caulibus erectis subramosis, foliis acutis serratis. *Sp. pl.* 591. *Fl. suec.* 536.

Acinos multis. *Bauh. hist.* III. 259. *R. Syn.* 238.

Ocimum sylvestre. *Ger. Em.* 675.

Clinopodium arvense *ocymi* facie. *Bauh. pin.* 225.

Clinopodium minus f. vulgare. *Park.* 21.

Anglis, wild Basil.

Habitat *in collibus aridis glareosis et cretaceis passim*. ☉. VII. VIII.

MELISSA. 647.

Cal. aridus, supra planiusculus : lab. superiore subfastigiato. *Corollæ* lab. superius fornicatum bifidum; labii inferioris lobo medio cordato.

Calamintha. 1. MELISSA pedunculis axillaribus dichotomis longitudine foliorum. *Sp. pl.* 593.

Calamintha vulgaris. *Park.* 36. *R. Syn.* 243. *off.* *Ger. Em.* 687.

Calamintha vulgaris, vel officinarum Germaniæ. *Bauh. pin.* 228.

Anglis, common Calamint.

Habitat *ad vias et in agrorum marginibus sed rarius*. 4. VIII.

Pharmac. Calaminthæ montanæ *Herba*.

nepeta. 2. MELISSA pedunculis axillaribus dichotomis folio longioribus, caule decumbente. *Sp. pl.* 593.

Calamintha odore *Pulegii*. *Ger. Em.* 687. *R. Syn.* 243.

Calamintha Pulegii odore f. *Nepeta*. *Bauh. pin.* 228.

Calamintha altera odore *Pulegii*, foliis maculosis. *Park.* 36.

Anglis, Field Calamint.

Habitat

Habitat in marginibus viarum, agrorumque. 4. VIII.
Pharmac. Calaminthæ officinalis Herba.

MELITTIS. 650.

Cal, tubo corollæ amplior. Corollæ labium superius planum; lab. inferius crenatum. Antheræ cruciatæ.

1. MELITTIS. Hort. cliff. 390. Sp. pl. 597. *melisso-*
Melissa Fuchsi. Lob. ic. 515. Ger. Em. 690. R. Syn. *phyllum*.
242.

Melissophyllum Fuchsi. Park. 41.

Lamium montanum melissæ folio. Bauh. pin. 231.

Anglis, bastard Balm.

Habitat in sylvis et sepibus. In several woods in the west of England, as about Totness in Devonshire, and Haverford-west in Pembroke-shire. R. Syn. 4. VI.

PRUNELLA. 654.

Filamenta bifurca, altero apice antherifera.

1. PRUNELLA foliis ovato-oblongis petiolatis. Sp. pl. vulgaris.
600. Fl. suec. 540.

Prunella bracteis cordatis. Hort. cliff. 316.

Prunella. Ger. Em. 631. R. Syn. 238. vulgaris.
Park. 1680.

Prunella major folio non dissecto. Bauh. pin. 260.

Anglis, Self-heal.

Habitat in pratis et pascuis. 4. VIII.

Pharmac. Prunellæ Herba.

SCUTELLARIA. 653.

Cal. ore integro, post florescentiam clauso, operculato.

1. SCUTELLARIA foliis cordato-lanceolatis rugosis, galericulatis, floribus axillaribus. Fl. lapp. 239. Hort. lata.
cliff. 316. Sp. pl. 599.

Cassida palustris vulgatio flore cæruleo. Tourn. 182.

R. Syn. 244.

Q 4

Lyfimachia

DIDYNAMIA GYMNOSPERMIA.

Lyfimachia cærulea galericulata, seu *gratiola cærulea*. *Baub. pin.* 246.

Lyfimachia galericulata. *Ger. Em.* 477.

Lyfimachia cærulea, f. *latifolia major*. *Park.* 221.

Anglis, Hooded Willow-herb.

Habitat *in aquosis et ad ripas fluviorum*. 4. VIII.

minor.

2. SCUTELLARIA foliis cordato-ovatis subintegerimis floribus axillaribus.

Cassida palustris minima, flore purpurascente. *Tourn.* 182. *R. Syn.* 244.

Lyfimachia galericulata minor. *R. Syn. II.* 132.

Gratiola latifolia. *Ger. Em.* 581. f. *nostras minor*. *Park.* 221.

Anglis, the lesser hooded Willow-herb.

Habitat *in palustribus*. 4. VII. VIII.

ANGIOSPERMIA.

* *Calyces bifidi.*

OROBANCHE. 697.

Cal. bifidus. *Cor.* ringens. *Capsula* unilocularis, bivalvis, polysperma. *Glandula* sub basi germinis.

major.

1. OROBANCHE caule simplicissimo pubescente, staminibus subexsertis. *Syst. nat.* 1119. *Sp. pl.* 632.

Orobanche major Garyophyllum olens. *Baub. pin.* 87. *R. Syn.* 288.

Rapum Genistæ. *Ger. Em.* 1311. *Park.* 229.

- β *Orobanche* flore minore. *Baub. hist. II.* 781.

Anglis, Broom-rape.

Habitat *in pratis ficcis*. 4. V. VI.

ramosa.

2. OROBANCHE caule ramoso corollis quinquefidis. *Sp. pl.* 633.

Orobanche ramosa. *Baub. pin.* 88. *Park.* 1363. *R. Syn.* 288.

Anglis, branched Broom-rape.

Habitat

Habitat *in arvis*. Found near Beccles in Suffolk, by Mr. J. Sherard. R. Syn. VI.

** *Calyces quadrifidi*.

L A T H R Æ A. 661.

Cal. quadrifidus. *Glandula* depresso, ad basin futuræ germinis. *Capsf.* unilocularis.

1. LATHRÆA caule simplicissimo, corollis pendulis: *squamalabio inferiore trifido*. *Sp. pl.* 606. *Fl. suec.* 550. *ria*.
Anblatum cordi f. *Aphyllon*. *Bauh. hist.* II. 783.
R. Syn. 288.

Orobanche radice dentata major. *Bauh. pin.* 88.

Orobanche radice dentata, f. *Dentaria major* Matthiolo. *Ger. Em.* 1322.

Anglis, Tooth-wort.

Habitat *in umbrosis et ad radices montium sed rarius*.
In the woods about Maidstone in Kent, Mr. J. Sherard. *R. Syn.* *in a shady lane leading from Harefield town to the river, plentifully*. *Blackf. spec. bot. in dumetis infra Conzick-scar, prope Kendal in Westmorlandia*. 4. IV. V.

B A R T S I A. 657.

Cal. bilobus, emarginatus, coloratus. *Corolla* minus ipso calyce colorata; labio superiore longiore.
Capsf. bilocularis.

1. BARTSIA foliis superioribus alternis ferratis, flori-
bus lateralibus. *Sp. pl.* 602.

Euphrasia major lutea latifolia palustris. *R. Syn.* 285.

Pluk. ph. t. 27. *f.* 5. *Hist. ox.* III. 432.

Anglis, Marsh Eye-bright Cow-wheat,

Habitat *in palustribus*. *In boggy and watery places towards the further end of Cornwall*. *R. Syn.* ☉. VIII.

2. BARTSIA foliis oppositis cordatis obtuse ferratis. *alpina*.
Hort. cliff. 325. *Sp. pl.* 602. *Fl. suec.* 541.
Clinopodium alpinum hirsutum. *Bauh. pin.* 225.
Pluk. ph. t. 163. *f.* 5.

Euphrasia

DIDYNAMIA ANGIOSPERMIA.

Euphrasia rubra Westmorlandica, foliis brevibus obtusis. *R. Syn.* 285.

Anglis, Mountain Eye-bright Cow-wheat.

Habitat *ad ripas fluviorum, locis apricis duris. Prope Orton vicum in Westmorlandia, juxta rivulum qui decurrit secus viam qua inde Crosbeiam itur. R. Syn.* 2. VIII.

RHINANTHUS. 658.

Cal. quadrifidus, ventricosus. *Caps.* bilocularis, obtusa, compressa.

Crista galli.

1. RHINANTHUS corollum labio superiore compresso brevior. *Fl. lapp.* 248. *Sp. pl.* 603. *Fl. suec.* 542. Pedicularis f. *Crista galli* lutea. *Park.* 713. *R. Syn.* 284.

Pedicularis pratensis lutea five *Crista galli*. *Bauh. pin.* 163.

Crista galli. *Ger. Em.* 1071.

- β Pedicularis major angustifolia ramosissima, flore minore luteo, labello purpureo. *R. Syn.* 284.

Anglis, yellow Rattle, or Cock's-comb.

Habitat *in pratis et pascuis.* ☉. VI. VII.

EUPHRASIA. 659.

Cal. quadrifidus, cylindricus. *Capsula* bilocularis, ovato-oblonga. *Antheræ* inferiores altero lobo basi spinosa.

- officinalis.* 1. EUPHRASIA foliis ovatis argute dentatis. *Sp. pl.* 604. *Fl. suec.* 543.

Euphrasia. *Bauh. hist. III.* 432. *Ger. Em.* 563. *R. Syn.* 284.

Euphrasia vulgaris. *Park.* 1329.

Euphrasia officinarum. *Bauh. pin.* 233.

Anglis, Eye-bright.

Habitat *in pratis et pascuis.* ☉. VIII. IX.

Pharmac. *Euphrasie Herba.*

- odontites.* 2. EUPHRASIA foliis linearibus: omnibus ferratis. *Hort. cliff.* 326. *Roy. lugdb.* 299. *Sp. pl.* 604.

Euphrasia

Euphrasia pratensis rubra. *Bauh. pin.* 234. *R. Syn.*

284. *major.* *Park.* 1329.

Cratægonon Euphrosine. *Ger. Em.* 91.

Anglis, red Eye-bright.

Habitat in arvis et pascuis. ☉. VIII. IX.

MELAMPYRUM. 660.

Cal. quadrifidus. *Cor.* labium superius compressum, margine replicato. *Caps.* bilocularis obliqua, hinc dehiscens. *Sem.* duo, gibba.

1. MELAMPYRUM spicis quadrangularibus compactis obtusis imbricatis. *Sp. pl.* 605. *Fl. suec.* 545.

Melampyrum cristatum, flore albo et purpureo. *Bauh. hist.* III. 440. *R. Syn.* 286.

Melampyrum luteum angustifolium. *Bauh. pin.* 234.

Anglis, crested Cow-wheat.

Habitat in sylvis Cantabrigiensibus et Bedfordiensibus copiose. *R. Syn.* ☉. VII.

2. MELAMPYRUM spicis conicis laxis laceris. *Sp. pl. arvense.* 605. *Fl. suec.* 511.

Melampyrum purpurascens coma. *Bauh. pin.* 234. *R. Syn.* 286.

Anglis, purple Cow-wheat.

Habitat in arvis. *In the corn, on the right hand, just before you come to Lycham in Norfolk.* *R. Syn.* *On a hill at Horsley bath, near Beeston-castle, Cheshire, Mr. Vernon.* *Blackf. spec. bot.* ☉. VII.

3. MELAMPYRUM florum conjugationibus remotis, *pratense.* corollis clausis. *Fl. suec.* 548. *Sp. pl.* 605.

Melampyrum latifolium flore albo; labio inferiore duabus maculis luteis distincto. *Hist. pl. par. R. Syn.* 286.

Anglis, Meadow Cow-wheat.

Habitat in pratis et pascuis. *In the wood by Dr. Richardson's house at North Bierley.* *R. Syn.* ☉. VII. VIII.

- sylvaticum.* 4. MELAMPYRUM corollis hiantibus. *Sp. pl.* 605. *Fl. suec.* 549.
 Melampyrum sylvaticum flore luteo, sive Satureja lutea sylvestris. *Bauh. hist. III.* 441. *R. Syn.* 286.
 Melampyrum luteum latifolium. *Bauh. pin.* 234.
 Cratæogonon album. *Ger. Em.* 91.
 Cratæogonon vulgare. *Park.* 1326.
Anglis, yellow Cow-wheat.
 Habitat in *sylvis*. ☉. VI. VIII.

*** *Calyces quinquefidi.*

P E D I C U L A R I S. 664.

- Cal.* quinquefidus. *Capf.* bilocularis, mucronata, obliqua. *Semina* tunicata.
- sylvatica.* 1. PEDICULARIS caule ramoso, calycibus oblongis angulatis lævibus, corollis labio cordato. *Sp. pl.* 607. *Fl. suec.* 552.
 Pedicularis caule ramoso, floribus solitariis, calycibus quinquefidis crenatis. *Fl. suec.* I. 504.
 Pedicularis. *Ger. Em.* 1071. *pratensis* purpurea. *Bauh. pin.* 163.
 Pedicularis *pratensis* rubra vulgaris. *Park.* 713. *R. Syn.* 284.
Anglis, common Louse-wort.
 Habitat in *pratibus et ericetis uliginosis*. 24. V. VI.
- palustris.* 2. PEDICULARIS caule ramoso, calycibus calloso-punctatis, corollis labio obliquis. *Sp. pl.* 607. *Fl. suec.* 551.
 Pedicularis caule ramoso erecto, calycibus bifidis crenatis. *Hall. herb.* 621.
 Pedicularis *palustris* rubra elatior. *R. Syn.* 284.
 Pedicularis *danica* maxima. *Lob. ill.* 147.
Anglis, Marsh Louse-wort.
 Habitat in *pratibus et pascuis humidioribus*. VI.

A N T I R R H I N U M. 668.

- Cal.* pentaphyllus. Corollæ basis deorsum prominens, nectarifera. *Capf.* bilocularis.

* *Foliis*

* *Foliis angulatis.*

- *1. ANTIRRHINUM foliis cordatis quinquelobis al- *Cymba-*
ternis caulibus procumbentibus. *Syst. nat.* 1111. *laria.*

Hort. cliff. 32. *upf.* 175. *Sp. pl.* 612.

Linaria hederaceo folio glabro, f. Cymbalaria. Tourn.
169. *R. Syn.* 282.

Cymbalaria. Bauh. pin. 306.

Anglis, Ivy-leaved Toadflax.

Habitat in muris antiquis circa Londinum frequens. 4.
VI. X.

2. ANTIRRHINUM foliis hastatis alternis, caulibus *Elatine.*
procumbentibus, corollis calcaratis. *Sp. pl.* 612.

Linaria Elatine dicta folio acuminato. R. Syn. 282.

Elatine folio acuminato, in basi auriculato, flore lu-
teo. Bauh. pin. 253.

Elatine folio acuminato. Park. 553.

Elatine altera. Ger. Em. 623.

Anglis, sharp-pointed Fluellin.

Habitat in arvis. ☉. VIII. X.

Pharmac. Elatines Herba.

3. ANTIRRHINUM foliis ovatis alternis, caulibus *spurium.*
procumbentibus, corollis calcaratis. *Sp. pl.* 613.

Linaria Elatine dicta folio subrotundo. R. Syn. 282.

Elatine folio subrotundo. Bauh. pin. 252. *Park.* 553.

Veronica fœmina Fuchsi, f. Elatine. Ger. Em. 625.

Anglis, round-leaved Fluellin.

Habitat in arvis. ☉. VIII.

** *Foliis oppositis verticillatis.*

4. ANTIRRHINUM foliis sublinearibus: inferioribus *arvense.*
verticillatis, calycibus piloso-viscidis, floribus spi-
catis, caule erecto. *Sp. pl.* 614.

Antirrhinum foliis linearibus alternis recurvis, ramis
caule brevioribus. Roy. lugdb. 297.

Linaria arvensis cærulea. Bauh. pin. 213. *Dill. elth.*
199. *t.* 163. *f.* 198.

Linaria cærulea foliis brevioribus et angustioribus. R.
Syn. 282.

Anglis, Corn blew Toadflax.

Habitat in arvis et pratis aridis. ☉. VII. VIII.

5. AN-

repens.

5. ANTIRRHINUM foliis sublinearibus inferne quaternis calycibus capsulæ æqualibus. *Sp. pl.* 614.
 Antirrhinum foliis linearibus alternis sæpius oppositis acuminatis floribus laxè spicatis. *Roy. lugdb.* 297.
Linaria angustifolia, flore cinereo striato. *Dill. elth.* 198. *t.* 163. *f.* 197.
Anglis, creeping Toadflax.
 Habitat in Cornubia non longe ab oppido Penryn occidentem versus, in sepibus. *R. Syn.* 4. VIII.

*** *Foliis alternis.**monspes-
fulanum.*

6. ANTIRRHINUM foliis linearibus confertis, caule nitido paniculato, pedunculis spicatis nudis. *Roy. lugdb.* 297. *Sp. pl.* 616.
Linaria odorata monspessulana. *Bauh. hist. III.* 459. *R. Syn.* 282.
Linaria capillaceo folio odora. *Bauh. pin.* 213.
Anglis, sweet-smelling Toadflax.
 Habitat in arvis prope Gads-hill, in Cantio. 4. VII. VIII.

Linaria.

7. ANTIRRHINUM foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus imbricatis. *Sp. pl.* 616.
Linaria lutea vulgaris. *Ger. Em.* 550. *Bauh. hist. III.* 456. *R. Syn.* 281.
Linaria vulgaris nostras. *Park.* 458.
Linaria vulgaris lutea flore majore. *Bauh. pin.* 212.
 β *Peloria.* *Amoen. acad. I.* 282. *t.* 3.
Anglis, common yellow Toadflax.
 Habitat in pratis et pascuis sterilioribus β in pratis arenosis prope Clapham, in comitatu Surriensi. 4. VII. *Pharmac. Linariæ Herba.*

minus.

8. ANTIRRHINUM foliis lenceolatis obtusis alternis, caule ramosissimo diffuso. *Sp. pl.* 617.
Antirrhinum arvense minus. *Bauh. pin.* 212.
Antirrhinum minimum repens. *Ger. Em.* 549.
Antirrhinum sylvestre minimum. *Park.* 1334.
Linaria antirrhinum dicta. *R. Syn.* 283.
Anglis, the least Toadflax.
 Habitat in arvis solo glareoso. ☉. VI. X.

**** *Corollis*

**** *Corollis hiantibus aut ecaudatis, foliis alternis integris.*

*9. ANTIRRHINUM corollis ecaudatis, calycibus rotundatis. *Sp. pl.* 617.

Antirrhinum foliis lanceolatis petiolatis, calycibus brevissimis, racemo terminali. *Vir. cliff.* 61. *Roy. lugdb.* 294.

Antirrhinum majus alterum, folio longiore. *Bauh. pin.* 211.

Antirrhinum purpureum. *Ger. Em.* 549.

Anglis, the greater Snapdragon.

Habitat in muris antiquis; in clivis cretaceis inter Northfleet et Gravesend copiose. *Æ.* VI.

10. ANTIRRHINUM corollis ecaudatis, calycibus orontium. corolla longioribus. *Sp. pl.* 617.

Antirrhinum angustifolium sylvestre. *Bauh. hist.* III. 464. *R. Syn.* 283.

Antirrhinum arvense majus. *Bauh. pin.* 212.

Antirrhinum sylvestre medium. *Park.* 1334.

Antirrhinum minus. *Ger. Em.* 549.

Anglis, the least Snapdragon, or Calf's-snout.

Habitat in arvis. *Æ.* VII. VIII.

SCROPHULARIA. 674.

Cal. quinquefidus. *Corolla* subglobosa, resupinata. *Capf.* bilocularis.

1. SCROPHULARIA foliis cordatis basi transversis, nodosa. caule obtusangulo. *Sp. pl.* 619.

Scrophularia major. *Ger. Em.* 715. *R. Syn.* 283.

Scrophularia vulgaris. *Park.* 610.

Scrophularia nodosa foetida. *Bauh. pin.* 235.

Anglis, knobby-rooted Fig-wort.

Habitat in sylvis et sepibus humidis. *Æ.* VII.

Pharmac. Scrophulariæ Folia, Radix.

2. SCROPHULARIA foliis cordatis petiolatis decurrentibus, caule membranis angulato, racemis terminalibus. *Sp. pl.* 620.

Scrophularia aquatica major. *Bauh. pin.* 235. *R. Syn.* 283.

Betonica

Betonica aquatica. *Ger. Em.* 715. *major. Park.* 613.
Anglis, Water Fig-wort, or Betony.

Habitat in aquosis. 4. VII.

Pharmac. Scrophulariæ Δ . f. *Betonicæ aquaticæ*
Folia.

scorodo-
nia.

habitat in jersey

3. SCROPHULARIA foliis cordatis duplicato-ferratis,
racemo composito. *Roy. lugdb.* 296. *Sp. pl.* 620.

Scrophularia scorodonix folio. *Hort. R. Bl. Hist. ox.*
II. 482. *Pet. h. br. t.* 35. *f.* 11. *R. Syn.* 283.

Scrophularia melissæ folio. *Tourn.* 166.

Anglis, Balm-leaved Fig-wort.

Habitat ad sepes et rivulos. Found near the sea-shore
about St. Ives in Cornwall by Mr. Ed. Lhwyd. *R.*

Syn. 4. VIII. *Pluk. Alm.* 338. *t.* 59. *f.* 5.

vernalis.

*4. SCROPHULARIA foliis cordatis, pedunculis axil-
laribus solitariis dichotomis. *Sp. pl.* 620. *Roy.*
lugdb. 293.

Scrophularia flore luteo. *Ger. Em.* 717. *Bauh. pin.*
236. *pr.* 112. *Park.* 610.

Anglis, yellow Fig-wort.

Habitat ad sepes prope Mitcham, *in comitatu* Surrien-
si, *copiose.* δ . IV. V.

D I G I T A L I S. 676.

Cal. quinquepartitus. *Cor.* campanulata, quinquefi-
da, ventricosa. *Capf.* ovata bilocularis.

purpurea. 1. DIGITALIS calycinis foliolis ovatis acutis, corollis
obtusis labio superiore integro. *Hort. ups.* 178. *Sp.*
pl. 621.

Digitalis purpurea. *Bauh. hist. II.* 812. *Ger. Em.*
790. *R. Syn.* 283.

Digitalis purpurea, folio aspero. *Bauh. pin.* 243.

Digitalis purpurea vulgaris. *Park.* 1653.

β *Digitalis alba*, folio aspero. *Bauh. pin.* 243.

Anglis, purple Fox-glove.

Habitat ad vias et sepes passim. δ . VII.

Pharmac. *Digitalis Folia, Flores.*

SIBTHORPIA. 693.

Cal. quinquepartitus. *Corolla* quinquepartita æqualis.

Stam. paribus remotis. *Capsf.* compressa, orbicularis, bilocularis : dissipimento-transverso.

1. SIBTHORPIA foliis reniformi-subpeltatis crenatis. *europæa.*
Sp. pl. 631.

Alfina spuria pusilla repens foliis saxifragæ aureæ. Cat.
pl. ang. Pluk. ph. t. 7. f. 6. R. Syn. 352.

Chrysosplenium cornubienfe. Pet. herb. br. t. 6. f. 11.
Anglis, bastard Money-wort.

Habitat ad rivulos et scaturagines. In Cornubia et
Devonia frequens. R. Syn. circa Longsledale in
Westmorlandia. 4. VIII.

LIMOSELLA. 694.

Cal. quinquefidus. *Cor.* quinquefida, æqualis. *Stamina* per paria approximata. *Capsf.* unilocularis, bivalvis, polyspermia.

1. LIMOSELLA. *Fl. lapp.* 249. *suec.* 563. *Sp. pl.* 631. *aquatica.*
Plantaginella palustris. Baub pin. 190. *R. Syn.* 278.

Alfina palustris repens, foliis lanceolatis, floribus per-
exiguis. Pluk. ph. t. 74. f. 4.

Anglis, bastard Plantain.

Habitat in inundatis et ubi aquæ pluviales stagnant. 4.
VIII. IX.

Classis XV.

TETRADYNAMIA
SILICULOSA.* *Silicula integra, nec apice emarginata.*

MYAGRUM. 713.

*Silicula stylo conico terminata, loculo submonospermo.**fativum.* * I MYAGRUM filiculis ovatis pedunculatis polyspermis. *Sp. pl.* 641. *Fl. suec.* 564.*Myagrum. Ger. Em.* 273. *R. Syn. fativum. Baub. pin.* 109.*Myagrum sylvestre f. Pseudo-myagrum. Park.* 868.*Sesamum. Trag. II.* 656.*Camelina five Myagron. Dod. pempt.* 532.*Anglis, Gold of Pleasure.**Habitat in agris inter Linum sæpe copiose, cum quo ab exteris terris introductum. O. VI.*

SUBULARIA. 716.

*Silicula integra, ovata, valvis ovatis, concavis, dissepimento contrariis. Stylus filicula brevior.**aquatica.* I. SUBULARIA. *Fl. lapp.* 253. *suec.* 566. *Sp. pl.* 642. *Subularia erecta, junci foliis acutis mollibus. R. Syn.* 307.*Juncifolia sub aquis nascens, cochleariæ capsulis. R. Syn. ed. II.* 281.*Graminifolia aquatica, thlaspeos capitulis rotundis, septo medio filiculam dirimente. Pluk. ph. t.* 188. *f. 5.**Anglis, Awl-wort.**Habitat sub aquis, solo glareos, in lacu Hibernico Lough Neagh. R. Syn.* 4.

VELLA.

V E L L A. 714.

Silicula dissepimento valvulis duplo majore, extra ovato.

1. VELLA foliis pinnatifidis, filiculis pendulis. *Sp. pl. annua.*
641.

Nasturtium sylvestre erucæ affine. Baub. pin. 105.
R. Syn. 304.

Nasturtium sylvestre valentinum Clusio. Baub. hist. II.
920. *Park. 830.*

Eruca nasturtio cognata tenuifolia. Ger. Em. 247.
Anglis, Cresse Rocket.

Habitat on Salisbury-plain, not far from Stonehenge,
Mr. Lawson. R. Syn. ☉. VI.

D R A B A. 717.

Silicula integra, ovali-oblonga : valvis planiusculis :
dissepimento parallelis. *Stylus nullus.*

1. DRABA scapis nudis, foliis subserratis. *Syst. nat. verna.*
1227.

Draba caulibus nudis, foliis subdentatis, petalis bipartitis. Fl. suec. 567. Sp. pl. 642.

Draba vulgaris caule nudo, polygoni folio hirsuto.
Dill. gen. 122. Cat. giff 40.

Paronychia vulgaris. Ger. Em. 624. R. Syn. 292.

Paronychia vulgaris alines folio. Park. 556.

Bursa pastoris minor loculo oblongo. Baub. pin. 108.
Anglis, common Whitlow-grass.

Habitat ubique fere in ficioribus. ☉. IV.

2. DRABA caule ramoso, foliis cordatis dentatis am- muralis,
plexicaulibus. *Roy. lugdb. 333. It. gotl. 192. Fl.*
suec. 569. Sp. pl. 642.

Bursa pastoris major loculo oblongo. Baub. pin. 108.
pr. 50. R. Syn. 292.

Thlaspi veronicæ folio. Park. 843. Th. Robinsf. H.
of West. 95.

Anglis, Speedwel-leaved Whitlow-grass.

Habitat in fissuris rupium et pratis montosis in comitatibus
Westmorlandico, et Eboracensi passim. ☉. f.
3. V.

TETRADYNAMIA SILICULOSA.

- incana.* 3. DRABA foliis caulinis numerosis incanis, filiculis oblongis. *Sp. pl.* 643. *Fl. suec.* 568.
 Draba caule ramoso folioso foliis dentatis. *Fl. lapp.* 254.
 Lunaria contorta major. *Pet. Herb. br. t.* 48. γ . 3.
R. Syn. 291.
 Lunaria vasculo sub longo intorto. *R. Syn. ed. II.* 164.
Pluk. ph. t. 42. *f.* 1.
 β Paronychiæ similis sed major, perennis alpina repens.
R. Syn. 292.
Anglis, wreathen-podded Whitlow-grafs.
 Habitat in fissuris rupium et pratis montosis subhumidis, in comitatibus Eboracensi, Westmorlandico, Cumberlandico, passim et in montibus Arvonix. δ . V.

** *Silicula emarginata apice.*

LEPIDIUM. 717.

Silicula emarginata, cordata, polysperma: valvulis carinatis contrariis.

- latifolium.* 1. LEPIDIUM foliis ovato-lanceolatis integris ferratis.
Fl. suec. 571. *Sp. pl.* 644.
 Lepidium latifolium. *Baub. pin.* 97. *R. Syn.* 304.
 Piperitis f. Lepidium vulgare. *Park.* 855.
 Raphanus sylvestris officinarum. *Ger. Em.* 241.
Anglis, Dittander, or Pepperwort.
 Habitat in pratis et pascuis sed variis. Near the Hythe at Colchester, and at Heybridge near Maldon, by the water-side, plentifully; plentifully in the corn betwixt Benington and Mitton, in the north riding of Yorkshire. *R. Syn.* In the marshes near Grays in Essex, Dr. Watson. 4. VI. VII.
Pharmac. Lepidii Herba.

- ruderales.* 2. LEPIDIUM floribus diandris apetalis, foliis radicalibus dentato-pinnatis, ramiferis linearibus integerrimis. *Hort. cliff.* 331. *Fl. suec.* 572. *Sp. pl.* 645
 Nasturtium sylvestre, osyridis folio. *Baub. pin.* 105
Park. 829. *R. Syn.* 303. capsulis minimis. *Hist. ex. II.* 301. *f.* 3. *t.* 19. *f.* 7.

Nastur

Nasturtium angustifolium. *Fuchf. hist.* 307. *Dalech. hist.* 1181.

Thlaspi minus. *Ger. Em.* 263.

Anglis, narrow-leaved wild Cress, or Dittander.

Habitat in maritimis. At Maldon in Essex, Lynne in Norfolk, and Truro in Cornwall. *R. Syn.* ☉. VI.

3. *LEPIDIUM* foliis pinnatis integerrimis, petalis emarginatis calyce minoribus. *Syst. nat.* 1127. *Fl. suec. petræum.* 573. *Sp. pl.* 644.

Nasturtium montanum annuum tenuissime divisum. *D. Bobart. R. Syn.* 304.

Nasturtium pumilum vernal. *Bauh. pin.* 105.

Nasturtium petræum annuum nostras. *Pluk. ph. t.* 204. f. 5.

Cardamine pusilla saxatilis montana discoides. *Col. Ec. I.* 275. t. 274.

Anglis, Mountain Dittander.

Habitat on S. Vincent's rock. *R. Syn.* ☉. IV. V.

THLASPI. 719.

Silicula emarginata, obcordata, polysperma: valvulis navicularibus, marginato-carinatis.

1. *THLASPI* filiculis subrotundis, foliis sagittatis dentatis incanis. *Fl. suec.* 575. *Sp. pl.* 646. *campestre.*

Thlaspi vulgatus. *Bauh. hist. II.* 921. *R. Syn.* 305.

Thlaspi arvense, vaccariæ folio, majus. *Bauh. pin.* 106.

Thlaspi vulgatissimum. *Ger. Em.* 262.

Thlaspi mithridaticum f. *vulgatissimum, vaccariæ folio.* *Park.* 835.

Anglis, Mithridate, Mustard, or Bastard Cress.

Habitat in arvis solo arenoso. ☉. VI. VII.

Pharmac. *Thlaspeos vulgaris semina.*

2. *THLASPI* filiculis orbiculatis, foliis oblongis dentatis *arvense.* glabris. *Fl. lapp.* 237. *suec.* 574. *Sp. pl.* 646. *Mat. med.* 323.

Thlaspi arvense, filiquis latis. *Bauh. pin.* 105.

Thlaspi Dioscoridis. *Ger. Em.* 262. *R. Syn.* 305.

Thlaspi Drabæ folio. Park. 836.
Anglis, Treacle-mustard, or Penny-cress.
Habitat in arvis sed rarius. ☉. VI. VII.
Pharmac. Thlaspeos semina.

perfolia-
tum.

3. **THLASPI** filiculis subrotundis, foliis amplexicaulibus cordatis subserratis. *Sp. pl. 646.*
Thlaspi foliis radicalibus ovatis, caulinis amplexicaulibus lanceolatis. Guett. stamp. 142. Dalib. parif. 196.
Thlaspi perfoliatum minus. Bauh. pin. 106. Park. 837. R. Syn. 305.
Thlaspi minus Clusii. Ger. Em. 268.
Anglis, perfoliate Treacle-mustard.
Habitat in pratis montosis. Among the stone-pits between Whitney and Burford, in Oxfordshire On moist Limestone pastures in Yorkshire, Westmoreland, and Cumberland. R. Syn. ☉. VI.

hirtum.

4. **THLASPI** filiculis subrotundis pilosis, foliis caulinis sagittatis hirsutis. *Roy. lugdb. 333. Sauv. monsp. 120. Sp. pl. 646.*
Thlaspi vaccariæ incano folio perenne. R. Syn. 305.
Thlaspi capsulis hirsutis. Bauh. hist. II. 922.
Thlaspi villosum, capsulis hirsutis. Bauh. pin. 106. pr. 212.
Anglis, Perennial Mithridate Mustard.
Habitat in montosis, Cambro-britanniæ et alibi observati. R. Syn. 4. VII.

montanum

5. **THLASPI** filiculis obcordatis, foliis integerrimis glabris: radicalibus cuneiformibus; caulinis amplexicaulibus. *Syst. nat. 1128.*
Thlaspi filiculis obcordatis, foliis radicalibus cuneiformibus integerrimis. Sp. pl. 647.
Thlaspi foliis globulariæ. Bauh. hist. II. 926. R. Syn. 305.
Thlaspi montanum, glabro folio minus. Park. 842. Bauh. pin. 106.
Thlaspi alpinum, bellidis cœruleæ folio. Bauh. pin. 106.
Thlaspeos albi supini varietas. Ger. Em. 268.
Anglis,

Anglis, Mountain Mithridate Mustard.

Habitat in pascuis montosis. In the pastures about the ebbing and flowing well, a mile from Settle in Yorkshire, towards Ingleborough, and between Malham and Settle. R. Syn. 4. VII.

6. THLASPI filiculis obcordatis, foliis radicalibus pin-natifidis. *Sp. pl.* 647. *Bursa pastoris.* *Baub. hist.* II. 936. *Ger. Em.* 276. *R. Syn.* 306. *Bursa pastoris major vulgaris.* *Park.* 866. *Bursa pastoris major folio sinuato.* *Baub. pin.* 108.
- β *Bursa pastoris minor foliis integris.* *Celf. upf.* 13. *Anglis*, Shepherd's-purse. Habitat ubique in agris, cultis, ruderatis, muris viis. ☉. III. IV. *Pharmac.* *Bursæ pastoris Herba.*

COCHLEARIA. 720.

Silicula emarginata, turgida, scabra : valvulis gibbis obtusis.

1. COCHLEARIA foliis radicalibus subrotundis, caulinis oblongis subsinuatis. *Fl. lapp.* 256. *suec.* 577. *Mat. med.* 320. *Sp. pl.* 647. *Cochlearia.* *Baub. hist.* II. 942. *R. Syn.* 302. *rotundifolia.* *Ger. Em.* 401. *Cochlearia rotundifolia f. Batavorum.* *Park.* 285. *Cochlearia folio subrotundo.* *Baub. pin.* 110.
- β *Cochlearia minor rotundifolia nostras.* *Park.* 286. *Anglis*, common or Garden Scurvygrafs. Habitat in littoribus maritimis, et in montosis. Near Castleton in the Peak, upon Penigent and Ingleborough hills, and upon Stanemore in Yorkshire, near the Spittle. *R. Syn.* ♂. IV. V. *Pharmac.* *Cochleariæ Herba.*
2. COCHLEARIA foliis reniformibus integris. *Hort. cliff.* 498. *Roy. lugdb.* 335. *Sp. pl.* 647. *Cochlearia rotundifolia.* *Merr. pin.* *R. Syn.* 302. *Cochlearia minima ex montibus Walliæ.* *Boerb. ind.* alt. 10. *groenlandica.*

R 4

Anglis,

D^r. Von Troil in his letters on Iceland says, the food of people in that island is grass & herbs; & scurvy-grass (cochlearia) in particular makes them so fat, that they yield more than twenty pounds of fat. Quære, what species?

Anglis, Groenland Scurvygrafs.

Habitat in montibus Arvoniciis. ♂. IV. V.

- anglica.* 3. COCHLEARIA foliis ovato-lanceolatis finuatis.
 Cochlearia foliis radicalibus lanceolatis integerrimis,
 caulinis subfinuatis. *Mill. Dict. Hort.*
 Cochlearia folio sinuato. *Bauh pin.* 110. *R. Syn.* 303.
 Cochlearia vulgaris. *Park.* 285.
 Cochlearia britannica. *Ger. Em.* 401.
Anglis, English or common Sea Scurvygrafs.
 Habitat in littoribus maritimis. ♂. V.
Pharmac. Cochleariæ marinæ f. britannicæ *Herba.*
- danica.* 4. COCHLEARIA foliis hastatis angulatis. *Fl. suec.*
 578. *Sp. pl.* 647.
 Cochlearia marina folio anguloso parvo. *D. Lawson.*
R. Syn. 303.
 Cochlearia aremorica. *Tourn.* 215.
 Thlaspi hederaceum. *Lob. ic.* 615.
Anglis, Danish Scurvygrafs.
 Habitat in Walney Lancastriæ insula; near Llanba-
 drick church, Mr Lhwyd. *R. Syn.* ♂. V.
- Coronopus* 5. COCHLEARIA foliis pinnatifidis. *Fl. suec.* 579.
Sp. pl. 648.
 Coronopus Ruellii. *Ger. Em.* 427.
 Coronopus recta vel repens Ruellii. *Park.* 502.
 Nasturtium supinum capsulis verrucosis. *R. meth. em.*
 98. *syn.* 304.
 Ambrosia campestris repens. *Bauh. pin.* 138.
Anglis, Swine's Cressies.
 Habitat in uliginosis et ad vias. ☉. VI. VIII.
Pharmac. Coronopi ruellii *Herba.*
- Armoracia.* 6. COCHLEARIA foliis radicalibus lanceolatis crena-
 tis caulinis incis. *Fl. suec.* 580 *Sp. pl.* 648. *Mat.*
med. 321.
 Cochlearia folio cubitali. *Tourn.* 215.
 Raphanus rusticus. *Ger. Em.* 241. *Park.* 860.
Bauh. pin. 96. *R. Syn.* 301.
 Raphanus sylvestris f. Armoracia multis. *Bauh. pin.*
 851.

Anglis,

Anglis, Horfe-radish.

Habitat *in ruderatis et ad fossas*. Found plentifully about Alnwick, and elsewhere in Northumberland, and Dr. Lister upon the banks of Skiptonbeck, and elsewhere, in Bolland in Craven. *R. Syn.* 4. V.

Pharmac. Raphani rufficani Radix.

I B E R I S. 721.

Cal. irregularis : *petalis* duobus exterioribus majoribus. *Silicula* polysperma, emarginata.

1. IBERIS foliis sinuatis, caule nudo simplici. *Hort. nudicaulis cliff.* 328. *It. oeland.* 139. *Fl. suec.* 581. *Sp. pl.* 650.

Nasturtium petræum. *Tabern. ic.* 451. *Ger. Em.* 251. *R. Syn.* 303.

Nasturtium petræum, foliis burfæ pastoris. *Bauh. pin.* 104.

Burfæ pastoris minor, foliis incis. *Bauh. pin.* 108.

Burfæ pastoris minor. *Park.* 866.

Anglis, Rock Cresse.

Habitat *in glareosis*. Near Hampton-court, Richmond, and Barnes Commons, Mr. Doody. *R. Syn.* 303. *in ericeto* Putnejano, D. Ph. Miller. ☉. V.

S I L I Q U O S A.

* *Calyx clausus* : *folioliis longitudinaliter conniventibus*.

D E N T A R I A. 726.

Siliqua elastice diffiliens, valvulis revolutis. *Stigma* emarginatum. *Calyx* longitudinaliter connivens

*1. DENTARIA foliis inferioribus pinnatis, summis bulbifera. simplicibus *Fl. suec.* 584 *Sp. pl.* 653.

Dentaria bulbifera. *Ger. Em.* 984.

Dentaria heptaphyllos baccifera. *Bauh. pin.* 322.

Anglis, Coral-wort.

Habitat *in nemorosis*. In Old Park wood, near Harefield, abundantly. *Blackf. fascic. pl.* 23. 4. IV.

CHEIRANTHUS. 730.

Germen utrinque denticulo glandulato. *Cal.* clausus :
foliolis duobus basi gibbis. *Semina* plana.

Cheiri.

1. CHEIRANTHUS foliis lanceolatis acutis glabris,
ramis angulatis. *Syst. nat.* 1133. *Hort. cliff.* 334.
upf. 187. *Mat. med.* 335. *Sp. pl.* 661. *Roy. lugdb.*
337. *Gort. gelr.* 382. *Guett. stamp. II.* 156.

Leucojum luteum, vulgo *Cheiri* flore simplici. *Bauh.*
hist. II. 872. *R. Syn.* 291.

Leucojum luteum vulgare. *Bauh. pin.* 202.

Keiri f. *Leucojum* vulgare luteum. *Park.* 625.

Viola lutea. *Ger. Em.* 456.

Anglis, Wall-flower, or wild *Cheir*.

Habitat in muris antiquis et tectis. 4. IV. V.

Pharmac. *Cheiri Flores.*

tricuspidatus.

2. CHEIRANTHUS filiquis apice tridentatis. *Hort.*
cliff. 335. *Sp. pl.* 663.

Leucojum marinum majus. *Park.* 622. *R. Syn.* 291.

Leucojum maritimum sinuato folio. *Bauh. pin.* 201.

Anglis, Sea Stockgilly-flower.

Habitat in maritimis Cambriæ littoribus arenosis, and
on the coast of Cornwall. *R. Syn.* 4. VI.

ERYSIMUM. 729.

Siliqua columnaris, exacte tetraedra. *Cal.* clausus.

- efficinale.* 1. ERYSIMUM filiquis spicæ appressis. *Fl. suec.* 598.
Sp. pl. 660.

Erysimum vulgare. *Bauh. pin.* 298.

Erysimum Dioscoridis Lobelio. *Ger. Em.* 254.

Ioio f. *Erysimum.* *Dod. pempt.* 714. *Lob. ic.* 206.
vulgare. *Park.* 833.

Eruca hirsuta, filiqua caule appressa, *Erysimum* dicta.
R. Syn. 298.

Anglis, Hedge Mustard.

Habitat juxta muros, inque plateis et ruderatis passim.

©. V.

Pharmac. *Erysimi Herba.*

2. ERY-

2. ERYSIMUM foliis integris lanceolatis. *Fl. lapp. Cherian-*
 263. *suec.* 601. *Sp. pl.* 661. *thoides.*
Myagro affinis planta siliquis longis. *Baub. hist. II.*
 894. *R. Syn.* 298.
Myagrum siliqua longa. *Baub. pin.* 109.
Camelina. *Ger. Em.* 273. f. *myagrum* alterum ama-
 rum. *Park.* 867.
Anglis, Treacle Wormseed.
Habitat in insula Eliensi in salicetis prope urbem Ely
copiosam observavimus; item prope Ashburn oppidum
in Pecco Derbieni dicto. In the corn-fields about
Elden. *R. Syn.* ☉. VII.
3. ERYSIMUM foliis lyratis : extimo subrotundo. *Fl. Barbarea.*
suec. 599. *Sp. pl.* 660.
Erysimum foliis basi pinnato-dentatis apice subrotun-
 dis. *Fl. lapp.* 264.
Eruca lutea seu *Barbarea.* *Baub. pin.* 98. *R. Syn.* 297.
Barbarea. *Baub. hist. II.* 868. *Ger. Em.* 243. flore
 simplici. *Park.* 819.
 β *Barbarea* foliis minoribus et frequentius sinuatis. *R.*
hist. 809.
Anglis, Winter Cresses, or Rocket.
Habitat in fossis et aquosis frequens. ♀. V.
4. ERYSIMUM foliis cordatis. *Hort. cliff.* 333. *Fl. Alliaria.*
suec. 600. *Sp. pl.* 660.
Hesperis allium redolens. *Hist. ox. II.* 252. *Tourn.*
 222. *R. Syn.* 293.
Alliaria. *Ger. Em.* 796. *Park.* 112. *Baub. pin.* 110.
Till. ic. 60.
Anglis, Jack-by-the-hedge, or Sawce-alone.
Habitat ad sepes passim. ♀. V.

H E S P E R I S. 731.

Petala obliqua flexa. *Glandula* intra stamina brevi-
 ora. *Siliqua* stricta. *Stigma* basi bifurca apice
 connivente. *Cal.* clausus.

1. HESPERIS caule simplici erecto, foliis ovato lance-*matrona-*
 olatis denticulatis, petalis mucrone emarginatis.*lis.*
Hort.

TETRADYNAMIA SILIQUOSA.

Hort. cliff. 188. *Roy. lugdb.* 338. *Dalib. paris.* 197. *Sp. pl.* 663.

Hesperis sylvestris inodora. *Baub. pin.* 202. *R. Syn.* 293.

Hesperis pannonica inodora. *Baub. hist. II.* 878. *Park.* 628.

Anglis, unfavory Dame's-violet.

Habitat on the banks of the rivers about Dalehead and Grassmire in Westmorland. *R. Syn.* ♂. V.

R A P H A N U S. 736.

Cal. clausus. Siliqua torosa, subarticulata, teres. Glandula melliferæ duo inter stamina breviora et pistillum, totidem inter stamina longiora et calycem.

Raphanistrum. I. **RAPHANUS** filiquis teretibus articulatis lævibus unilocularibus. *Hort. cliff.* 340. *Sp. pl.* 669. *Fl. suec.* 612.

Raphanus sylvestris. *Ger. Em.* 240.

Raphanistrum filiqua articulata glabra, majore et minore. *Hist. Ox. II.* 265. *R. Syn.* 296.

Rapistrum album articulatum. *Park.* 863.

Rapistrum flore albo, filiqua articulata. *Baub. pin.* 95.

β *Rapistrum flore luteo, filiqua glabra articulata.* *R. Syn.* 296.

γ *Raphanus maritimus flore luteo, filiquis articulatis.* *R. Syn.* 296.

Anglis, α white flowered Charlock, β yellow flowered Charlock.

Habitat in arvis. ☉. VI. VII.

B R A S S I C A. 734.

Cal. erectus, connivens. Semina. globosa. Glandula inter stamina breviora et pistillum, interque longiora et calycem.

orientalis. I. **BRASSICA** foliis cordatis amplexicaulibus glabris; radicalibus scabris integerrimis, filiquis tetragonis. *Sp. pl.* 666.

Brassica campestris perfoliata flore albo. *Baub. pin.* 112. *R. Syn.* 293.

Perfoliata filiquosa. *Ger. Em.* 536. *Baub. hist.* 335. *Perfo.*

Perfoliata filiquosa vulgaris f. *Brassica campestris*.
Park. 580.

Anglis, perfoliate Cabbage.

Habitat in arvis et in clivis maritimis prope Harvicum
 oppidum in Efflexia et in clivis juxta Bardsey prope
 Orford in Suffolcia a D. Dale. *R. Syn.* ☉. VI.

2. BRASSICA radice caulescente tereti carnosa. *Hort. oleracea*.
cliff. 338. *Fl. suec.* 608.

Brassica maritima arborea f. procerior ramosa. *Hist.*
Ox. II. 208. *R. Syn.* 293.

Brassica maritima. *Pet. H. br. t.* 45. f. 6.
Anglis, Sea Cabbage.

Habitat in clivis maritimis. On Dover cliffs, &c. *R.*
Syn. prope castellum Harleck in comitatu Mervinia.
 ♂. IV.

3. BRASSICA radice caulescente orbiculari depresso *Rapa*.
 carnosa. *Hort. cliff.* 339. *ups.* 190. *Roy. lugd.*
 344. *Sp. pl.* 666.

Rapa sylvestris. *R. hist.* 800.

Rapa sativa rotunda. *Bauh. pin.* 89. *R. Syn.* 294.

Rapum majus. *Ger. Em.* 232.

Rapum sativum rotundum. *Bauh. hist.* 838.

- β *Rapa sativa* oblonga seu foemina. *Bauh. pin.* 90.
Anglis, Turneps.

Habitat in arvis et ad margines agrorum. ♂. IV.

Pharmac. Rapi Radix, semina.

4. BRASSICA radice caulescente fusiformi. *Hort. cliff. Napus*.
 339. *Fl. suec.* 609. *Sp. pl.* 666.

Napus sylvestris. *Bauh. pin.* 95. *hist. II.* 843. *R.*
Syn. 295. *It. goet.* 240. et *W.—goeth.* 132.

Bunias seu *Napus sylvestris* nostras. *Park.* 865.

Bunias sylvestris Lobelii. *Ger. Em.* 235.

Anglis, wild Navew.

Habitat inter segetes et in aggeribus fossarum. ♂. V.

5. BRASSICA foliis dentato-pinnatifidis, caule hispido, *Erucaf-*
 filiquis lævibus. *Sp. pl.* 667. *trum.*

Eruca sylvestris. *Ger. Em.* 247. *R. Syn.* 296. vul-
 gator. *Park.* 818.

Eruca

TETRADYNAMIA SILIQUOSA.

Eruca major lutea, caule aspero. Bauh. pin. 98.

Anglis, wild Rocket.

Habitat in muris antiquis et ruderalis frequens. 4. V.

T U R R I T I S. 733.

Siliqua longissima, angulata. Cal. connivens, erectus. Cor. erecta.

- glabra.* 1. TURRITIS foliis radicalibus dentatis hispidis, caulinis integerrimis amplexicaulibus glabris. *Fl. suec. 606. Sp. pl. 666.*
Turritis foliis inferioribus cichoraceis, reliquis perfoliatis. Tourn. 224.
Brassica sylvestris foliis circa radicem cichoraceis. Bauh. pin. 109.
Brassica sylvestris foliis circa radicem cichoraceis asperis, caulibus autem adhærentibus planis, seu glabris. Hist. Ox. II.
 β *Brassica sylvestris foliis integris et hispidis. Bauh. pin. 112.*
Turritis. Ger. Em. 272. R. Syn. 293. vulgarior. Bauh. hist. II. 836.
Anglis, great Tower-mustard.
Habitat in pratis glareosis. Frequent in Surry and Kent, in pits and uncultivated places, Dr. Watson, near Colchester, M. Newton, ☉. V.

- hirsuta.* 2. TURRITIS foliis omnibus hispidis, caulinis amplexicaulibus. *Fl. suec. 607. Sp. pl. 666.*
Turritis muralis minor. Pet. H. br. t. 47. 12. R. Syn. 294.
Erysimo similis hirsuta, non laciniata, alba. Bauh. pin. 101. pr. 42. ic.
Erysimo similis hirsuta planta. Park. 834.
Anglis, hairy Tower-mustard.
Habitat in rupibus et muris antiquis. ♂. VI.

A R A B I S. 732.

Glandulæ nectariferæ quatuor, singulæ intra calycis foliola, squamæ instar reflexæ.

- italiana.* 1. ARABIS foliis petiolatis lanceolatis integerrimis. *Virid.*

rid. cliff. 64. *Fl. suec.* 985. *Roy. lugdb.* 339. *Dalib. parif.* 200. *Sp. pl.* 665.

Turritis foliis lanceolatis integris petiolatis, ad exortum ramorum solitariis. *Hort. cliff.* 339.

Turritis vulgaris ramosa. *Tourn.* 224. *R. Syn.* 294.

Brassica spuria minima, foliis hirsutis et glabris. *R. Syn. ed. II.* 166.

Bursæ pastoris similis siliquosa major et minor. *Bauh. pin.* 108.

Pilosella siliquosa. *Thal. tab.* 7.

Paronychia major et altera minor. *Park.* 556.

β Turritis minor foliosa. *Pet. Herb. br. t.* 48. *f.* 2.

Anglis, coded Mouse-ear.

Habitat in muris tectis et pratis aridis. ☉. V.

** *Calyx hians : foliolis superne distantibus.*

C A R D A M I N E. 727.

Siliqua elastice diffiliens valvulis revolutis. *Stigma* integrum. *Cal.* subhians.

* *foliis simplicibus.*

1. CARDAMINE foliis simplicibus ovatis integerrimis, bellidifolii petiolis longissimis. *Fl. lapp.* 260. *t.* 9. *f.* 2. *suec. lia.* 590. *Sp. pl.* 654.

Cardamine pumila bellidis folio, alpina. *Ger. Em.* 260. *R. Syn.* 301.

Plantula cardamines alterius æmula Clusii. *Park.* 812.

Nasturtium alpinum bellidis folio minus. *Bauh pin.* 105.

Anglis, Daïsle-leaved Ladies-smock.

Habitat in variis locis juxta Rippon ad monasterium ejus et in Denbighshire. *Merr. pin.* Found by Mr. Newton on S. Vincent's rock near Bristol. *R. Syn.* 2. IV. V.

2. CARDAMINE foliis simplicibus oblongis dentatis. *petræa.* *Sp. pl.* 654. *Fl. suec.* 591.

Cardamine petræa cambrica, nasturtii facie. *Dill. elth.* 70. *t.* 61. *f.* 71.

Nasturtium petræum. *Merc. bot. Pluk. ph. t.* 101. *f.* 3. *R. Syn.* 300.

Anglis, mountain Ladies-smock.

Habitat in excelsis rupibus Arvonix et Mervinix; in rupibus udis supra lacum Layn-du copiose. *D. Lhwyd.* *R. Syn.* 2. V. ** *Foliis*

** *Foliis pinnatis.*

- pratensis.* 3. CARDAMINE foliis pinnatis : foliolis radicalibus subrotundis ; caulinis lanceolatis. *Sp. pl.* 656. *Fl. suec.* 585.
 Cardamine. *Ger. Em.* 259. *R. Syn.* 299.
 Nasturtium pratense magno flore. *Bauh. pin.* 104.
 Nasturtium pratense majus f. Cardamine latifolia. *Park.* 825.
 Flos cuculi. *Dod. pempt.* 592.
Anglis, common Ladies-smock, or Cuckow-flower.
 Habitat in pratis et pascuis subhumidis. 4. IV.
Pharmac. Cardamines Herba.
- amara.* 4. CARDAMINE foliis pinnatis : axillis stoloniferis.
Sp. pl. 656. *Fl. suec.* 586.
 Cardamine foliis pinnatis : foliolis subrotundo-angulosis. *Hall. helv.* 558.
 Cardamine flore majore elatior. *Tourn.* 224. *R. Syn.* 299.
 Nasturtium aquaticum majus et amarum. *Bauh. pin.* 104.
 Nasturtium aquaticum amarum. *Park.* 1239.
Anglis, bitter Cresses, or Ladies-smock.
 Habitat in pratis et pascuis humidis. Circa Lewisham, D. Alchorne ; by the river-side at Harefield, and about Uxbridge, plentifully. *Blackf. spec. bot.* 4. IV. V.
- impatiens.* 5. CARDAMINE foliis pinnatis : foliolis laciniatis, petalis fugatissimis.
 Cardamine foliis pinnatis : floribus apetalis. *Sp. pl.* 655. *Fl. suec.* 588.
 Cardamine impatiens, vulgo Sium minus impatiens. *Ger. Em.* 260. *R. Syn.* 299.
 Cardamine annua. exiguo flore. *Tourn.* 224.
 Sium minimum, Noli me tangere dictum, f. impatiens nasturtii sylvestris folio. *Park.* 1241.
Anglis, impatient Ladies smock.
 Habitat in pratis montosis ad rivulos et in rupibus humidis. In comitatibus Eboracensi et Westmorlandica passim. 6. IV.

6. CAR-

6. CARDAMINE foliis pinnatis exstipulatis, foliolis *parviflora* lanceolatis obtusis floribus corollatis. *Syst. nat.*

1131.

Cardamine pratensis, parvo flore. *Tourn.* 224.

Cardamine annua glabra, parvo flore, lobis foliorum vix sinuatis. *Hort. chelf. ind.*

Cardamine quarta Dalechampii. *Lugd. hist.* 654.

Anglis, small-flowered Ladies-smock.

Habitat in pratis, pascuis et ad rivulos frequens. ☉.

III. V.

7. CARDAMINE foliis pinnatis floribus pentandris. *hirsuta*.

Sp. pl. 655. *Fl. suec.* 587.

Cardamine impatiens altera hirsutior. *R. Syn.* 300.

Nasturtium aquaticum minus. *Bauh. pin.* 104.

Anglis, hairy Ladies-smock.

Habitat in pratis pascuis et locis umbrosis, passim. ☉.

IV.—VIII.

SISYMBRIUM. 728.

Siliqua dehiscens valvulis rectiusculis. *Calyx* patens.

Corolla patens.

* *Siliquis* declinatis brevibus.

1. SISYMBRIUM filiquis declinatis, foliis pinnatis: *Nasturtium* foliolis subcordatis. *Hort. cliff.* 336. *Fl. suec.* 592.

Mat. med. 331. *Sp. pl.* 657.

Sisymbrium Cardamine, seu Nasturtium aquaticum.

Bauh. hist. II. 884. *R. Syn.* 300.

Nasturtium aquaticum vulgare. *Park.* 1239.

Nasturtium aquaticum supinum. *Bauh. pin.* 104.

Nasturtium aquaticum f. *Cratevæ* Sium. *Ger. Em.*

257.

- β Nasturtium aquaticum foliis minoribus præcocius.

R. Syn. 301.

- γ Nasturtium aquaticum pinnulis paucioribus. *Dood.*

R. Syn. 301.

• *Anglis*, Water-creffes.

Habitat in aquosis et rivulis. ♂. 4.

• *Pharmac.* Nasturtii aquatici *Herba.*

S

2. SY-

- sylvestre.* 2. *SISYMBRIUM* filiquis declinatis, foliis pinnatis: foliolis lanceolatis ferratis. *Roy. lugdb.* 336. *Dalib. paris.* 204. *Sauv. monsp.* 228. *Boehm. lips.* 118. *Sp. pl.* 657.
Eruca aquatica. *Ger. Em.* 248. *Park.* 1242. *R. Syn.* 297.
Eruca sylvestris minor, luteo parvoque flore. *Bauh. pin.* 98.
Anglis, Water Rocket.
Habitat in palustribus et aquosis præsertim solo glareoso.
In Tothil-fields, Westminster, plentifully. *Blackf. spec. Bot.* 4. VI.

- ambili-* 3. *SISYMBRIUM* filiquis declinatis oblongo-ovatis, fo-
um. liis pinnatifidis ferratis. *Sp. pl.* 657. *Fl. suéc.* 593.
Sisymbrium foliis simplicibus dentatis ferratis. *Hort. cliff.* 336.
aquaticum Lin. 6 *Rhaphanus aquaticus alter.* *Bauh. pr.* 38. *Park.* 1229. *Fulham about the Bishop's Stairs.*
Rhaphanus aquaticus rapistrifolio. *Bauh. pin.* 97.
Rapistrum aquaticum. *Ger. Em.* 240.
 β *Sisymbrium foliis pinnatifidis ferratis.* *Fl. suéc.* 551.
Sisymbrium foliis infimis capillaceis, summis pinnatifidis. *Hort. cliff.* 337.
Rhaphanus aquaticus foliis in profundas lacinias divis. *Bauh. pin.* 97. *pr.* 38. *R. Syn.* 301.
Anglis, Water Radish.
Habitat in aquosis. 4. VI.

** *Caule nudo.*

- murale.* 4. *SISYMBRIUM* foliis ferrato-dentatis lævibus, caule nudo adscendente scabro. *Syst. nat.* 1132.
Sisymbrium foliis lyratis denticulatis, filiquis sessilibus. *Sp. pl.* 658.
Eruca monensis laciniata lutea. *R. Syn.* 297.
Eruca sylvestris minor lutea, burfæ pastoris folio. *Bauh. pin.* 98. *pr.* 39. *Pluk. ph. t.* 86. *f.* 7. *Barr. ic.* 1016.
Anglis, Wall Rocket.
Habitat in arenosis. *Found by Mr. Lawson in Sella-fields, Seabank, Cumberland; also between Marsh-grange*

grange and the isle of Walney, but not plentifully,
by Mr. Lhwyd, near Abermeney ferry in Angle-
sea. *R. Syn.* 4. VI.

5. *SISYMBRIUM* foliis dentato-pinnatis lævibus, caule *monense*,
nudo adscendente glabro. *Syst. nat.* 1132.

Sisymbrium foliis pinnatifidis integerrimis, caulibus
subnudis simplicibus procumbentibus. *Sp. pl.* 658.

Eruca monensis laciniata, flore luteo majore. *Dill.*
elth. 135. *t.* 111. *f.* 135.

Anglis, yellow Rocket.

Habitat in arenosis in Monæ Insula. D. Dillenius. 3.
4. VI.

*** *Foliis bipinnatis.*

6. *SISYMBRIUM* foliis pinnato-hastatis dentatis nudis, *Irio*,
caule lævi, filiquis erectis. *Sp. pl.* 659. *Fl. suec.*
596.

Erysimum foliis lyrato-pinnatis : extimo hastato. *Fl.*
suec. ed. I. 556.

Erysimum latifolium Neapolitanum. *Park.* 834. *R.*
Syn. 1298.

Erysimum latifolium majus glabrum. *Bauh. pin.* 101.

Irio lævis Apulas, erucæ folio. *Col. eip. I.* 264. *t.*
265.

Anglis, broad-leaved Rocket, or Hedge Mustard.

Habitat in ruderatis et muris antiquis circa Londinum
frequens. ☉. V.—VIII.

7. *SISYMBRIUM* petalis calyce minoribus, foliis de- *Sophia*,
composito pinnatis. *Fl. suec.* 595. *Mat. med.*
332. *Sp. pl.* 659.

Erysimum Sophia dictum. *R. Syn.* 298.

Sophia Chirurgorum. *Ger. Em.* 1068. *Park.* 830.

Nasturtium sylvestre tenuissime divisum. *Bauh. pin.*
105.

Anglis, Flix-weed.

Habitat in maceris, ruderatis, tectis, frequens. ☉.
VII.

Pharmac. *Sophiæ semina.*

S I N A P I S. 735.

Cal. patens. *Cor.* unguis recti, *Glandula* inter stamina breviora et pistillum, interque longiora et calycem.

nigra.

1. SINAPIS filiquis glabris tetragonis. *Hort. cliff.* 338. *Lech scan.* 51. *Fl. suec.* 611. *Mat. med.* 326. *Sp. pl.* 668.

Sinapi rapi folio. *Bauh. pin.* 99.

Sinapi fativum secundum. *Ger. Em.* 243. *R. Syn.* 295.

Sinapi filiqua latiuscula glabra, semine rufo, f. vulgare. *Bauh. hist.* II. 885.

Anglis, common Mustard.

Habitat in arvis aggeribus fossarum et juxta vias. ☉. VI.

Pharmac. Sinapis Semina.

alba.

2. SINAPIS filiquis hispidis : rostro obliquo longissimo. *Hort. cliff.* 338. *ups.* 191. *Roy. lugdb.* 343. *Dalib. paris.* 200. *Sp. pl.* 668.

Sinapi album filiqua hirsuto, semine albo vel ruffo.

Bauh. hist. II. 856. *R. Syn.* 295.

Sinapi album. *Ger. Em.* 244.

Sinapi apii folio. *Bauh. pin.* 99.

Anglis, white Mustard.

Habitat in agris et ad vias. ☉. VIII.

Pharmac. Sinapis albæ semina.

arvensis.

3. SINAPIS filiquis multangulis toroso-turgidis rostro longioribus. *Fl. suec.* 610. *Sp. pl.* 668.

Rapistrum arvorum. *Ger. Em.* 233. *Park.* 862. *R. Syn.* 295.

Rapistrum flore luteo. *Bauh. pin.* 95. *hist.* II. 844.

Anglis, Wild-mustard, or Charlock.

Habitat in arvis et marginibus agrorum ubique. ☉. V.

C R A M B E. 739.

Filamenta quatuor longiora, apice bifurca : altero antherifero. *Bacca* sicca globosa decidua.

- maritima.* 1. CRAMBE foliis cauleque glabris. *Fl. suec.* 615. *Sp. pl.* 671.

Crambe foliis cordatis crispis carnosis. *Hort. cliff.* 340.

Crambe

Crambe maritima brassicæ folio. Tourn. 211. R. Syn.

307.

Brassica maritima monospermos. Bauh. pin. 112.

Brassica marina monospermos. Park. 270.

Brassica marina anglica. Ger. Em. 315.

Anglis, Sea Colewort.

Habitat in littoribus arenosis maritimis frequens. 4. V.

B U N I A S. 737.

Silicula decidua, tetraëdra, angulis inæqualibus acuminatis muricatis.

1. *BUNIAS filiculis ovatis lævibus ancipitibus. Fl. suec. Cakile. 613. Sp. pl. 670.*

Raphanus filiquis ovatis angulatis monospermis. Hort. cliff. 340.

Cakile quibusdam, aliis Eruca marina et Raphanus marinus. Bauh. hist. II. 867. R. Syn. 307.

Eruca marina. Ger. Em. 248. anglica. Park. 821.

Eruca maritima Italica, filiqua hastæ cuspidi simili. Bauh. pin. 99.

Anglis, Sea Rocket.

Habitat in littoribus maritimis passim. 6. VI.

I S A T I S. 738.

Siliqua lanceolata, unilocularis, monosperma, decidua, bivalvis: valvulis navicularibus.

*1. *ISATIS foliis radicalibus crenatis, caulinis sagittatis tinctoria. filiculis oblongis. Fl. suec. 614. Sp. pl. 67.*

Isatis sylvestris, f. angustifolia. Bauh. pin. 113.

β *Isatis fativa f. latifolia. Bauh. pin. 113.*

Glaustum fativum. Ger. Em. 49. Park. 600. R. Syn. 307.

Anglis, Woad.

Habitat in arvis et marginibus agrorum inter Bath et Bristol, copiose. 8. VII.

Classis XVI.

MONADELPHIA
DECANDRIA.

GERANIUM. 746.

Monogyna. *Stigmata* quinque. *Capf.* rostrata, pentacocca.

* *Staminibus* quinque fertilibus.

- cicutari-* 1. GERANIUM pedunculis multifloris, calycibus pentaphyllis, floribus pentandris, foliis pinnatis incis obtusis. *Hort. cliff.* 344. *Fl. suec.* 625. *Sp. pl.* 680.
um. Geranium cicutæ folio inodorum. *Ger. Em.* 945. *R. Syn.* 357.
Geranium cicutæ folio, minus et supinum. *Bauh. pin.* 319.
Geranium moschatum inodorum. *Park.* 708.
β Geranium inodorum album. *Ger. Em.* 946.
Anglis, Hemlock-leaved Cranebill.
Habitat in ruderatis, muris, et juxta vias, ubique. ☉.
IV.—VI.
- moscha-* 2. GERANIUM pedunculis multifloris, calycibus pentaphyllis, floribus pentandris, foliis pinnatis incis, cotyledonibus pinnatifidis. *Syst. nat.* 1143.
tum. Geranium moschatum. *Ger. Em.* 941. *Park.* 706. *R. Syn.* 358.
Geranium cicutæ folio, moschatum. *Bauh. pin.* 319.
β Geranium pimpinellæ folio. *Cat. giff.* 173.
Anglis, musked Cranebill, or Muscovy.
Habitat in pratis siccioribus. In comitatibus Westmorlandico et Eboracensi passim. ☉. V.
Pharmac. Geranii Moschati Herba.

3. GERANIUM pedunculis subtrifloris, foliis cordatis *maritimo*-crenato-incisus subvillosis, caulibus procumbentibus. *mun.*
Syst. nat. 1143.

Geranium pedunculis bifloris, foliis ovatis hirsutis crenato-incisus, caulibus procumbentibus. *Roy. lugdb.* 391.

Geranium pusillum supinum maritimum Althææ aut Betonicæ folio nostras. *R. Syr.* 357. *Pluk. ph. t.* 31. f. 4.

- Geranium minimum anglicum procumbens, foliis betonicæ. *Hist. ox. II.* 512. t. 35.

Anglis, Sea Cranesbill.

Habitat in arenosis et glareosis prope littus occidentale, v. g. in insulis Mona et Prestholma; in Wallia circa Carnarvon; Cornubia circa Pensans et alibi copiosissime. In Sufflexiæ ora prope Brackelsham, D. Dillenius. *R. Syn.* 4. VI. VII.

*** *Staminibus decem fertilibus.*

4. GERANIUM pedunculis bifloris, foliis caulinis trinodofum. lobis integris serratis, summis subseffilibus. *Hort. cliff.* 343. *Roy. lugdb.* 350. *Sp. pl.* 681.

Geranium V. nodosum Plateau. *Clus. hist.* 101. *Ger. Em.* 947. *R. Syn.* 361.

Geranium nodosum. *Bauh. pin.* 318. *Park.* 704.

Geranium magnum folio trifido. *Bauh. hist. III.* 478. *Anglis*, knotty Cranesbill.

Habitat in montosis Cumberlandiæ sponte ortum affer-mavit. D. Archergen. *R. Syn.* 4. VII. VIII.

5. GERANIUM pedunculis bifloris foliisque alternis, *phæum*. calycibus subaristatis, caule erecto, petalis undulatis. *Sp. pl.* 681.

Geranium montanum fuscum. *Bauh. pin.* 318. *R. Syn.* 361.

Geranium maculatum five fuscum. *Ger. Em.* 942.

Geranium pullo flore. *Park.* 704.

Anglis, spotted Cranesbill.

Habitat in montosis. Found by Mr. Drayton apothecary of Maidstone, at Tovel, in the valley by the old fulling-mill. *R. Syn.* circa Clapham et Ingle-ton in agro Eboracensi. 4. IV. V.

S 4

6. GE-

G. malacoides. pedunculis multifloris, floribus pentandris, foliis cordatis sublobatis.

Habitat in Angliæ maritimis ©. Lin.

- sylvaticum* 6. GERANIUM pedunculis bifloris, foliis subpeltatis multifidis inciso-ferratis, caule erecto, petalis emarginatis. *Fl. lapp.* 266. *suec.* 617. *Sp. pl.* 684.
 Geranium batrachoides montanum nostras. *Ger. Em.* 942.
 Geranium batrachoides minus. *Park.* 704.
 Geranium batrachoides folio aconiti. *Bauh. pin.* 317. *Anglis*, Mountain Cranesbill.
 Habitat in pratis montosis et dumetis agri Westmoriandici et Eboracensis copiose. *R. Syn.* 4. VII. VIII.
- pratense*. 7. GERANIUM pedunculis bifloris, foliis subpeltatis multipartitis rugosis pinnato-laciniatis acutis, petalis integris. *Fl. suec.* 618. *Sp. pl.* 681.
 Geranium batrachoides. *Ger. Em.* 942. *Bauh. hist.* III. 475. *R. Syn.* 360.
 Geranium batrachoides flore cæruleo. *Park.* 704.
 Geranium batrachoides, Gratia Dei Germanorum, Lobelio. *Bauh. pin.* 318.
Anglis, Crowfoot Cranesbill.
 Habitat in pratis et pascuis subhumidis et ad margines agrorum. Circa Battersea. *R. Syn.* 4. VI. VII.
- robertianum*. 8. GERANIUM pedunculis bifloris, calycibus pilosis decemangulatis. *Sp. pl.* 681. *Fl. suec.* 619.
 Geranium pedunculis bifloris, foliis quinque-tripartitis : lobis pinnatifidis. *Hort. cliff.* 344. *Mat. med.* 336.
 Geranium robertianum. *Ger. Em.* 939. *Bauh. pin.* 319. *R. Syn.* 358.
 Geranium robertianum vulgare. *Park.* 710.
 β Geranium lucidum faxatile, foliis geranii robertiani. *R. Syn.* 358.
Anglis, Herb Robert.
 Habitat in sepibus, muris, ruderatis, frequens. ♂. IV. —VI.
Pharmac. Geranii robertiani Herba.
- lucidum*. 9. GERANIUM pedunculis bifloris, calycibus pyramidalis

dati angulatis rugosis, foliis quinquelobis rotundatis. *It. gotl.* 228. *Fl. suec.* 620. *Sp. pl.* 682.

Geranium saxatile. Ger. Em. 938. *Park.* 707. *R. Syn.* 361.

Geranium lucidum saxatile. Bauh. pin. 318.

Anglis, shining Dovesfoot Cranesbill.

Habitat in tectis, muris, et umbrosis. ☉. VI. VII. V.

10. GERANIUM pedunculis bifloris, petalis integris *rotundifolium.* longitudine calycis, caule prostrato, foliis reniformibus incis. *Sp. pl.* 683. *Fl. suec.* 621.

Geranium folio malvæ rotundo. Bauh. pin. 318.

Geranium columbinum majus, flore minore cæruleo. R. Syn. 358.

Pes columbinus. Dod. pempt. 61.

Anglis, round-leaved Cranesbill.

Habitat in muris, tectis et ad sepes circa Bath et Bristol copiose, inter Battersea et Wandsworth. ☉. VII.

Pharmac Geranii columbini Herba.

11. GERANIUM pedunculis bifloris, foliis inferioribus *perenne.* quinque-partito-multifidis rotundatis, superioribus trilobis, caule erecto.

Geranium columbinum, pyrenaicum perenne, maximum. Tourn. 268.

Anglis, perennial Dovesfoot Cranesbill.

Habitat in pratis montosis. Ad ripam fluvii inter Bingley et Kighley in agro Eboracensi; prope Enfield, et inter Hyde-Park et Little-Chelsea. 4. V. —VIII.

12. GERANIUM pedunculis bifloris, foliisque floralibus alternis, caule ramoso diffuso, calycibus muticis. *Sp. pl.* 682. *Fl. suec.* 624.

Geranium pedunculis bifloris, foliis quinque-partito-multifidis rotundatis, laciniis obtusiusculis, capsulis hirsutis. Fl. suec. ed. I. 577.

Geranium columbinum villosum, petalis bifidis purpureis. Vaill. paris. 79. t. 15. f. 3.

Geranium columbinum. Ger. Em. 938. *R. Syn.* 359. vulgare. *Park.* 706.

Anglis, common Dovesfoot Cranesbill.

Habitat

Habitat in arvis, pratis, pascuis et ad sepes. ☉. V.
VI.

pufillum. 13. GERANIUM pedunculis bifloris, petalis bifidis, caule prostrato, foliis reniformibus palmatis linearibus acutis. *Syst. nat.* 1144.

Geranium columbinum humile, flore cæruleo minimo. *R. Syn.* 359. *t.* 16. *f.* 2.

Geranium malacoides f. columbinum minimum. *Park.* 707.

Anglis, small-flowered Dovesfoot Cranebill.

Habitat in arvis et ad vias præsertim solo arenoso. ☉. VI.

columbinum. 14. GERANIUM pedunculis bifloris, foliis quinquepartito-multifidis angulatis: laciniis acutis, capsulis glabris, calycibus aristatis. *Fl. suec.* 623. *Sp. pl.* 682.

Geranium columbinum dissectis foliis, pediculis florum longissimis. *R. Syn.* 359. *Vaill. parif.* 79. *t.* 15. *f.* 4.

Geranium columbinum annuum minus, folio tenuius laciniato, flore pediculo longissimo insistente. *Hist. ox. II.* 512. *f.* 5. *t.* 15. *f.* 3.

Anglis, long-stalked Dovesfoot Cranebill.

Habitat in arvis et pratis montosis. Circa Charleton in Cantio. ☉. VII.

dissectum. 15. GERANIUM pedunculis bifloris, foliis quinquepartito-multifidis, petalis emarginatis longitudine calycis, capsulis villosis. *Cent. pl.* 61. *Fl. suec.* 622.

Geranium columbinum majus dissectis foliis. *Ger. Em.* 938. *R. Syn.* 359.

Geranium malacoides f. columbinum alterum. *Park.* 706.

Geranium columbinum maximum, foliis dissectis. *Pl. hist. nat. ox.* 145. *t.* 9. *f.* 3. *R. Syn.* 360.

Geranium columbinum majus foliis imis longe usque ad pediculum divisis. *Hist. ox. II.* 511.

Anglis, jagged-leaved Dovesfoot Cranebill.

Habitat in pratis et pascuis. ☉. V. VII.

16. GERANIUM pedunculis unifloris, foliis quinquepartitis trifidis orbiculatis. *Fl. suec.* 616. *Sp. um. pl.* 685.

Geranium hæmatodes. *Park.* 705. *R. Syn.* 360.

Geranium sanguineum. *Ger. Em.* 945.

Geranium sanguineum maximo flore. *Baub. pin.* 318.

- β Geranium hæmatodes foliis majoribus, pallidioribus et altius incisis. *R. Syn.* 360.

- γ Geranium hæmatodes Lancastrienſe flore eleganter ſtriato. *R. hiſt.* 1061. *ſyn.* 360.

Anglis, bloody Craneſbill.

Habitat in rupibus, pratis ſiccioribus et ad ſepes. γ. in inſula Walney. 4. VII. VIII.

P O L Y A N D R I A.

A L T H Æ A. 749.

Cal. duplex : exterior novemfidus. *Caps.* plurimæ, monospermæ.

1. ALTHÆA foliis ſimplicibus tomentofis. *Hort. cliff. officinalis.* 348. *upſ.* 204. *Mat. med.* 338. *Gort. gelr.* 410. *Sp. pl.* 686.

Althæa vulgaris. *Park.* 303. *R. Syn.* 252.

Althæa Ibiſcus. *Ger. Em.* 933.

Althæa Dioſcoridis et Plinii. *Baub. pin.* 315.

- β Althæa vulgari ſimilis folio reſuſo brevi. *R. Syn.* 252. *Anglis*, Marſhmallow.

Habitat in paluſtribus maritimis frequens. 4. VIII.

Pharmac. Althææ Radix, Herba, ſemina.

L A V A T E R A. 752.

Cal. duplex : exterior trifidus. *Caps.* plurimæ, monospermæ.

1. LAVATERA caule arboreo, foliis ſeptemangulatis tomentofis plicatis, pedunculis confertis unifloris axillaribus. *Hort. upſ.* 202. *Sp. pl.* 690.

Malva arborea minima noſtras, *Merr. pin.* *R. Syn.* 252.

Anglis, Sea-tree Mallow.

Habitat

Habitat *ad littoribus maritimis. At Hurst-castle, over against the Isle of Wight, in Portland island, on the rocks of Caldey island, and on the Basse island near Edinburgh. R. Syn. ♂. VIII.*

M A L V A. 751.

Cal. duplex: exterior triphyllus. Caps. plurimæ, monospermæ.

sylvestris. 1. MALVA caule erecto herbaceo, foliis lobatis obtusis, pedunculis petiolisque pilosis. *Fl. suec. 627. Sp. pl. 689.*

Malva caule erecto, foliis subpalmatis obsolete serratis. *Hort. cliff. 347.*

Malva sylvestris. *Ger. Em. 930. folio sinuato. Baub. pin. 314.*

Malva vulgaris. *Park. 299. R. Syn. 251.*

Anglis, common Mallow.

Habitat *ad sepes et semitas inque locis ruderalis. ♂. V.—X.*

Pharmac. Malvæ vulgaris Herba, Radix, semina.

*rotundi-
folia.* 2. MALVA caule prostrato, foliis cordato-orbiculatis obsolete quinquelobis. *Fl. suec. 626. Sp. pl. 688.*

Malva sylvestris pumila. *Ger. Em. 930.*

Malva sylvestris folio rotundo. *Baub. pin. 314.*

Malva sylvestris minor. *Park. 299. R. Syn. 250.*

Anglis, Dwarf Mallow.

Habitat *in ruderalis et ad vias. ☉. VI.—X.*

*parviflo-
ra.* 3. MALVA caule patulo, foliis angulatis, floribus axillaribus glomeratis, calycibus glabris patentibus. *Amoen. III. 416.*

Malva minor, flore parvo cæruleo. *R. Syn. 251.*

Malva sylvestris, foliis sinuatis minoribus, flosculis minimis, Anglica. *Boerb. ind. alt. 268.*

Anglis, small-flowered Mallow.

Habitat *in arenosis. Found by Mr. J. Sherard, in company with Mr. Rand, at Hithe in Kent. R. Syn. ♂. VI.*

Alcea. 4. MALVA caule erecto, foliis multipartitis scabriusculis. *Fl. suec. 628. Sp. pl. 689.*

Alcea

Alcea vulgaris. *Clus. hist. II. 25. R. Syn. 252.*

Alcea vulgaris major. *Bauh. pin. 316.*

Alcea vulgaris f. Malva verbenacea. *Park. 301.*

Malva verbenacea. *Ger. Em. 930.*

Anglis, Vervain Mallow.

Habitat in sepibus et ad margines agrorum. In comitatibus Warwicensi, Leicestriensi, et Eboracensi, 24.

VII. IX.

Pharmac. Alceæ Herba.

5. *MALVA* foliis radicalibus reniformibus incis, cau- *moschata.*
linis quinquepartitis pinnato-multifidis. *Hort. ups.*
202. *Fl. suec. 629. Sp. pl. 690.*

Malva montana f. Alcea rotundifolia laciniata. *Col. ecphr. I. 148. t. 147. f. 147.*

Alcea folio rotundo laciniato. *Bauh. pin. 316.*

Alcea tenuifolia crispa. *Bauh. hist. II. ap. 1067.*

Anglis, jagged-leaved Vervain Mallow.

Habitat in pratis pascuis sepibus. 3. VIII.

Classis

Classis XVII.

DIADELPHIA
HEXANDRIA.

FUMARIA. 760.

Cal. diphyllus. *Cor.* ringens. *Filamenta* duo membranacea singula. *Antheris* tribus.

officinalis. 1. FUMARIA pericarpis monospermis racemosis, caule diffuso. *Sp. pl.* 700. *Fl. suec.* 630.

Fumaria vulgaris. *Bauh. hist.* III. 101. *Park.* 287. *R. Syn.* 204.

Fumaria purpurea. *Ger. Em.* 1088.

Fumaria officinarum et Dioscoridis. *Bauh. pin.* 143. *Anglis*, Fumitory.

Habitat in agris, arvis, satis, aggeribus, frequens. ☉. IV.—VI.

Pharmac. Fumarizæ Herba, Semina.

capreolata 2. FUMARIA pericarpis monospermis racemosis, foliis scandentibus subcirrhosis. *Sp. pl.* 701.

Fumaria major scandens flore pallidiore. *R. Syn.* 204. *Anglis*, ramping Fumitory.

Habitat in sylvis, sepibus, in agro Westmorlandico passim. ☉. VIII.

claviculata. 3. FUMARIA filiquis linearibus, foliis cirrhiferis. *Sp. pl.* 701.

Fumaria alba latifolia. *Park.* 288. *R. Syn.* 335: claviculata. *Ger. Em.* 1088.

Fumaria claviculis donata. *Bauh. pin.* 143. *Hist. Ox.* II. 266. f. 3. t. 12. f. 3.

Anglis, climbing Fumitory.

Habitat in sylvis, sepibus, subhumidis, præsertim solo glareoso. ☉. VI.

OCTANDRIA

POLYGALA. 761.

Cal. pentaphyllus: foliolis duobus alæformibus coloratis. *Legumen* obcordatum, biloculare.

1. POLYGALA floribus cristatis racemosis, caulibus *vulgaris*. herbaceis simplicibus procumbentibus, foliis linearilanceolatis. *Amen. acad. II.* 136. *Sp. pl.* 702. *Fl. suec.* 632.

Polygala. *Ger. Em.* 563. *R. Syn.* 287. minor. *Park.* 1332.

Polygala vulgaris. *Bauh. pin.* 215.

β Polygala violacea. *Boerb. ind. alt. I.* 236.

γ Polygala carnea. *Boerb. ind. alt. I.* 236.

δ Polygala alba. *Tourn.* 174.

ε Polygala major. *Bauh. pin.* 215.

ζ Polygala myrtifolia palustris humilis et ramosior. *R. Syn.*

Anglis, Milkwort.

Habitat in ericetis. 4. V. VI.

DECANDRIA

SPARTIUM. 765.

Stigma longitudinale, supra villosum. *Filamenta* carinæ adhærentia. *Cal.* deorsum productus.

1. SPARTIUM foliis ternatis solitariisque, ramis inerscoparium: mibus angulatis. *Hort. cliff.* 356. *Sp. pl.* 709. *Fl. suec.* 633.

Genista angulosa et Scoparia. *Bauh. pin.* 395.

Genista angulosa trifolia. *Bauh. hist. I.* 388. *R. Syn.* 475.

Genista. *Ger. Em.* 1311. vulgaris et Scoparia. *Park.* 228.

Anglis, common Broom.

Habitat in pascuis siccioribus. 5. V. VI.

Pharmac. Genistæ Herba, Flores, Semina.

G E N I S T A. 766.

Cal. bilabiatus. *Vexillum* oblongum a pistillo staminibusque deorsum reflexum.

* *Inermes.*

tinctoria. 1. GENISTA foliis lanceolatis glabris, ramis striatis teretibus erectis. *Mat. med.* 347. *Sp. pl.* 710. *Fl. suec.* 634.

Genista tinctoria germanica. *Bauh. pin.* 395.

Genista tinctoria vulgaris. *Park.* 228.

Genistella tinctoria. *Ger. Em.* 1316. *R. Syn.* 474. *Anglis*, Green-wood, Dyer's-weed, or Wood-waxen.

Habitat in pascuis et ad agrorum margines. *h.* VII.

** *Spinosæ.*

anglica. 2. GENISTA spinis simplicibus, ramis stoloniferis inermibus foliis lanceolatis. *Hort. cliff.* 355. *Roy. lugdb.* 371. *Sp. pl.* 710.

Genista minor aspalathoides, f. Genista spinosa anglica. *Bauh. pin.* 395. *R. Syn.* 475.

Genista aculeata. *Ger. Em.* 1320.

Genistella aculeata. *Park.* 1004.

Anglis, Needle-furze, or Petty-whin.

Habitat in ericetis subhumidis. *h.* V. VI.

U L E X. 786.

Cal. diphyllus. *Legumen* vix calyce longius.

europæus. 1. ULEX foliis villosis acutis, spinis sparsis. *Sp. pl.* 741.

Ulex folio sub singulis spinis subulato plano acuto.

Hort. cliff. 356. *ups.* 212. *Roy. lugdb.* 372.

Genista spinosa vulgaris. *Ger. Em.* 1319. *R. Syn.* 475.

Genista spinosa vulgaris, f. Scoparius Theophrasti, quem Gaza Nepam transtulit. *Park.* 1003.

β Genista spinosa minor. *Park.* 1003.

Anglis, Furze, Whins, or Gorse.

Habitat in ericetis. *h.* IV. V.

ONONIS. 772.

Cal. quinquepartitus : laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum, sessile.

1. ONONIS floribus racemosis solitariis, foliis ternatis, *spinosa*. ramis spinosis. *Syst. nat.* 1159.

Ononis floribus racemosis geminatis, foliis ternatis, ramis spinosis. *Sp. pl.* 716. *Fl. suec.* 637.

Anonis spinosa flore purpureo. *Bauh. pin.* 389. *Park.* 994. *R. Syn.* 332.

Anonis f. Resta bovis. *Ger. Em.* 1323. *Anglis*, Restharrow, or Cammock.

Habitat in pascuis sterilioribus. 4. VI.—VIII. *Pharmac.* Anonidis Radix.

2. ONONIS floribus racemosis geminatis, foliis ternatis *arvensis*. ramis inermibus. *Syst. nat.* 1159.

Ononis floribus subsessilibus solitariis lateralibus, ramis inermibus. *Hort. cliff.* 359. *Fl. suec.* 637.

Anonis non spinosa purpurea. *Ger. Em.* 1323. *Park.* 993. *R. Syn.* 332.

Anonis spinis carens purpurea. *Bauh. pin.* 389. *Anglis*, Corn Restharrow, or Cammock.

Habitat in pascuis sterilioribus. 4. VII. VIII.

3. ONONIS caulibus diffusis, ramis erectis, foliis caulinis ternatis, stipulis ovatis. *Syst. nat.* 1159. *Sp. pl.* 717.

Anonis maritima procumbens, foliis hirsutiae pubescentibus. *R. Syn.* 332. *Dill. elth.* 29. t. 25. f. 28.

Anglis, creeping Restharrow.

Habitat in littoribus maritimis frequens, in pratis et pascuis circa Charleton in Cantio copiose. 4. VI. VII.

ANTHYLLIS. 773.

Cal. ventricosus. *Legumen* subrotundum, testum.

2. ANTHYLLIS herbacea, foliis pinnatis inæqualibus, *Vulneraria*. capitulo duplicato. *Sp. pl.* 719. *Fl. suec.* 638.

T Anthyllis

DIADELPHIA DECANDRIA.

Anthyllis foliis pinnatis: foliolis pluribus, terminatrice majore. Roy. lugdb. 389.

Anthyllis leguminosa. Ger. Em. 1240. vulgaris. Park.

1093.

Vulneraria rustica. Bauh. hist. II. 362. R. Syn. 323.

Loto affinis vulneraria pratensis. Bauh. pin. 332.

β *Vulneraria supina flore coccineo. R. Syn. 323.*

Anglis, Kidney-vetch, or Ladies Finger.

Habitat in pascuis siccioribus præsertim cretaceis. 4. VII.

P I S U M. 779.

Stylus triangulus, supra carinatus pubescens. Calycis laciniae superiores duabus breviores.

marinum. 1. PISUM petiolis supra planiusculis, caule angulato, stipulis sagittatis, pedunculis multifloris. It. W—goth. 190. Sp. pl. 727. Fl. suec. 639.

Pisum marinum. Ger. Em. 1250. R. Syn. 319.

Pisum spontaneum maritimum anglicum. Park. 1059. Anglis, Sea Pease.

Habitat in littoribus maritimis. On the long baick of stones running from Aldburgh towards Orford, in Suffolk, on the end next to Orford, abundantly; we found it also near Hastings in Suffex; among the stones on the west side of Dengeness, near Lyd in Kent. R. Syn. 4. VII.

O R O B U S. 780.

Stylus linearis. Cal. basi obtusus: laciniis superioribus profundioribus, brevioribus.

tuberosus. 1. OROBUS foliis pinnatis lanceolatis, stipulis semisagittatis caule simplici. Sp. pl. 728. Fl. suec. 642.

Orobus sylvaticus foliis oblongis glabris. Tourn. 393. R. Syn. 324.

Astragalus sylvestris foliis oblongis glabris. Bauh. pin. 351.

Astragalus sylvaticus. Ger. Em. 1237.

Lathyrus sylvestris lignosior. Park. 1072.

Anglis, Wood Pease, or Heath Pease.

Habitat in pratis, pascuis, nemoribus et ad sepes. 4. V.

2. ORO-

2. OROBUS caulibus decumbentibus hirsutis ramosis. *salvaticus*.
Cent. pl. 67.

Orobis sylvaticus nostras. *R. Syn. 324.*

Anglis, English Wood Vetch, or bitter Vetch.

Habitat in sylvis et sepibus. At Gamlesby in Cumberland about six miles from Pereth, in the way to Newcastle, in the hedges and pastures, plentifully; observed by Tho. Willisel below Brecknock hills, in the way to Caerdyff, and in Merionethshire, near Bala. *R. Syn. 4. VII.*

LATHYRUS. 781.

Stylus planus, supra villosus, superne latior. Cal. laciniae superiores duabus breviores.

* *Pedunculis unifloris.*

1. LATHYRUS pedunculis unifloris, foliis simplicibus, *Nissolia*, stipulis subulatis. *Sp. pl. 729.*

Lathyrus sylvestris minor. *Bauh. pin. 344.*

Vicia folio gramineo, filiqua porrectissima. *Merr. pin.*

Catanance leguminosa quorundam. *Bauh. hist. II. 309. R. Syn. 323.*

Ervum sylvestre. *Ger. Em. 1249. f. Catanance. Park. 1079.*

Anglis, Crimfon Grafs-vetch.

Habitat in agrorum marginibus non infrequens. ☉. V.

2. LATHYRUS pedunculis unifloris, cirrhis aphyllis, *Aphaca*, stipulis sagittato-cordatis. *Sp. pl. 729.*

Aphaca. Ger. Em. 1250. Park. 1067. R. Syn. 320.

Vicia lutea, foliis convolvuli minoris. *Bauh. pin. 345.*

Anglis, yellow Vetchling.

Habitat inter segetes sed rarius. ☉. VI. VII.

** *Pedunculis bifloris.*

3. LATHYRUS pedunculis bifloris, cirrhis diphyllis, *hirsutus*, foliolis lineari-lanceolatis, leguminibus hirsutis, seminibus scabris. *Roy. lugdb. 363. Dalib. parif. 215. Sp. pl. 732.*

DIADELPHIA DECANDRIA.

Lathyrus filiqua hirsuta. *Bauh. hist. II.* 205. *R. Syn.* 320.

Lathyrus angustifolius filiqua hirsuta. *Bauh. pin.* 344. *Anglis*, rough-codded Chickling-vetch.

Habitat in arvis. *In the fields about Hockley and Raleghe, and elsewhere in Rochford hundred in Essex.* *R. Syn.* *In the border of a field between the road and Munden church; likewise among bushes within the gate leading to Laydon-hall, both in Dengy hundred, Essex, Mr. S. Dale.* *Blackf. spec. bot.* 4. VII.

*** *Pedunculis multifloris.*

latifolius. 4. **LATHYRUS** *pedunculis multifloris, cirrhis diphyllis, foliolis lanceolatis, internodiis membranaceis.* *Fl. suec.* 645. *Sp. pl.* 733.

Lathyrus major latifolius. *Ger. Em.* 1229. *R. Syn.* 319.

Lathyrus latifolius. *Bauh. pin.* 344.

Lathyrus major perennis. *Park.* 1061.

Anglis, broad-leaved Pease-everlasting.

Habitat in sylvis et sepibus. *In Madingley and other woods near Cambridge; Mr. Lawson observed it on the rocks by Red-neefe by Whitehaven, plentifully.* *R. Syn.* 4. VII.

sylvestris. 5. **LATHYRUS** *pedunculis multifloris, cirrhis diphyllis; foliolis ensiformibus, internodiis membranaceis.* *Sp. pl.* 738. *Fl. suec.* 644.

Lathyrus sylvestris major. *Bauh. pin.* 344.

Lathyrus sylvestris. *Dod. pempt.* 523. *Clus. hist. II.* 129. *Park.* 1061.

Lathyri majoris species flore rubente et albido minore dumetorum, f. Germanicus. *Bauh. hist. II.* 302. *R. Syn.* 319.

Anglis, narrow-leaved Pease-everlasting.

Habitat in sylvis et sepibus. *Mr. Dale observed it near Cattle-Campes, in the hedges by the way that leads thence to Bartlow in Cambridgeshire.* *R. Syn.* *Circa Rochester in Cantio; inter Bath et Bristol copiose.* 4. VII. VIII.

6. LA-

6. LATHYRUS pedunculis multifloris, cirrhis poly-*palustris*.
phyllis stipulis lanceolatis. *Fl. suec.* 648. *Sp. pl.*
733.

Lathyrus peregrinis, foliis viciæ, flore subcæruleo
pallidive purpurascente. *Bauh. pin.* 344.

Lathyrus viciæ formis f. Vicia Lathyroides nostras.
R. Syn. 320.

Vicia lathyroides f. Lathyrus viciæ formis. *R. pluk.*
ph. t. 71. *f.* 2.

Anglis, Marsh Chickling-vetch.

Habitat in pratis et pascuis subhumidis. In a wood
near Abingdon, Berkshire. *Blackf. spec. bot.*; upon
the bogs in Charley forest, near Bardon hill, Lei-
cesterhire, Mr. Pultney. *Phil. transf.* 4. VII.
VIII.

7. LATHYRUS pedunculis multifloris, cirrhis di-*pratensis*.
phyllis: foliolis lanceolatis: cirrhis simplicissimis.
Sp. pl. 733. *Fl. suec.* 647.

Lathyrus luteus sylvestris dumetorum. *Bauh. hist. II.*
304. *R. Syn.* 320.

Lathyrus sylvestris flore luteo. *Ger. Em.* 1231. *Park.*
1062.

Lathyrus sylvestris luteus, foliis viciæ. *Bauh. pin.*
344.

Anglis, Tare-everlasting, or common yellow Vetch-
ling.

Habitat in sylvis, sepibus, pratis, pascuis. 4. VII.
VIII.

V I C I A. 782.

Stigma latere inferiore transverse barbatum.

* *Pedunculis elongatis.*

1. VICIA pedunculis multifloris, floribus imbricatis, *cracca*.
foliolis lanceolatis pubescentibus, stipulis integris.
Sp. pl. 735. *Fl. suec.* 652.

Vicia multiflora. *Bauh. pin.* 345. *f.* spicata. *Park.*
1072.

Cracca, *Riv. tetr.* 49. *Rupp. jen.* 212. *R. Syn.* 322.

Anglis, tufted Vetch.

Habitat in nemorosis et sepibus. 2. VII. VIII.

sylvatica. 2. VICIA pedunculis multifloris, foliolis ovalibus, stipulis denticulatis. *Sp. pl.* 734. *Fl. succ.* 650.

Vicia sylvatica multiflora. *Ph. br.* R. Syn. 322.

Vicia multiflora maxima perennis, tetro odore, floribus albetibus lincis cæruleis striatis. *Fluk. ph. t.* 71. f. I.

Anglis, tufted Wood Vetch.

Habitat in sylvis et sepibus. In a wood near Caerwent in Monmouthshire; in the woods about Greta bridge, Yorkshire, Mr. Johnson; in Smoakhall wood by Bath, and at the Devizes, Wiltshire, Mr. Goodyer; supra pontem Kirby Lonsdale, Westmorlandiæ, Dr. Richardson. 2. VII. VIII.

** Floribus axillaribus subsessilibus.

sepium. 3. VICIA leguminibus pedicellatis subquaternis erectis, foliis ovatis integerrimis: exterioribus decrescen-
tibus. *Sp. pl.* 737. *Fl. succ.* 651.

Vicia sepium perennis. *Bauh. hist. II.* 713. R. Syn. 320.

Vicia sepium folio rotundiore acuto. *Bauh. pin.* 345.

Vicia maxima dumetorum. *Ger. Em.* 1727. *Park.* 1072.

β Vicia folio subrotundo, brevi, obtuse mucronato, pediculo brevi infidente, flore viciæ sepium seu dumetorum vulgaris. R. Syn.

Anglis, Bush Vetch.

Habitat in sylvis, sepibus, pratis et pascuis. 2. V.

fativa. 4. VICIA leguminibus sessilibus subbinatis erectis, foliis retusis, stipulis notatis. *Sp. pl.* 736. *Fl. succ.* 654.

Vicia leguminibus erectis, petiolis polyphyllis, foliolis acumine emarginatis, stipulis dentatis. *Hort. cliff.* 368.

Vicia. *Ger. Em.* 1227. R. Syn. 320. vulgaris fativa. *Park.* 1072.

Vicia fativa vulgaris, semine nigro. *Bauh. pin.* 344.

β Vicia fativa vulgaris, semine albo. *Bauh. pin.* 344.

Anglis, common Vetch, or Tare.

Habit 2x

Habitat in agris, inter segetes frequens, ☉. V. VI.
Pharmac. Viciæ semina.

5. VICIA leguminibus sessilibus solitariis erectis glabris *lathyroides*.
foliolis senis: inferioribus obcordatis. *Sp. pl.* 736.
Vicia sylvestris seu Cracca major. *Ger. Em.* 1227.
Vicia semine rotundo nigro. *Bauh. pin.* 345.
Vicia lathyroides, purpureo-cæruleis floribus. *Herm.*
par. 242. *t.* 242.
Aracus f. Cracca major. *Park.* 1070.
β Vicia sylvestris flore ruberrimo, filiqua longa nigra.
R. Syn. 321.
Anglis, Strangle Tare, or wild Vetch.
Habitat in arvis pratis et pascuis siccissimis præsertim
solo arenoso. ☉. V.

6. VICIA leguminibus sessilibus reflexis pilosis solitariis, *lutea*.
pentaspermis, corollæ vexillis glabris. *Sp. pl.* 736.
Vicia foliis subsessilibus obverse ovatis apice emarginatis
leguminibus solitariis villosis. *Dalib. parif.* 219.
Vicia sylvestris lutea filiqua hirsuta. *Bauh. pin.* 345.
Vicia luteo flore sylvestris. *Bauh. hist. II.* 313. *R.*
Syn. 321.
Anglis, yellow Vetch.
Habitat in pratis et pascuis. On Glastenbury torne
hill. *R. Syn.* ☉. VI. VII.

ERVUM. 784.

Calyx quinquepartitus longitudine corollæ.

1. ERVUM pedunculis subbifloris cirrhosis, petiolis a-*solonienſe*.
cuminatis, foliolis obrufis. *Cent. pl.*
Vicia minima præcox parisiensium. *Tourn.* 397.
R. Syn. 321.
Vicia minima. *Riv. teir.*
Vicia pratensis verna, f. præcox soloniensis semine
cubico f. hexahedron referente. *Hist. Ox. II.* 63.
f. 2. t. 4. f. 14.
Anglis, Spring Tare.

DIADELPHIA DECANDRIA.

Habitat *in pratis et pascuis solo cretaceo. On the chalky banks near Greenhithe, in Kent. R. Syn. ☉. IV. V.*

tetraspermum. 2. ERVUM pedunculis subbifloris, seminibus globosis quaternis. *Sp. pl. 738. Fl. succ. 655.*

Vicia segetum singularibus filiquis glabris. *Bauh. pin. 345.*

Cracca minor filiquis singularibus, flosculis cæruleiscentibus. *Hoff. cat. H. alt. R. Syn. 322.*

Anglis, smooth Tare.

Habitat *in arvis et ad marginibus agrorum. ☉. VI.*

hirsutum. 3. ERVUM pedunculis multifloris, seminibus globosis binis. *Sp. pl. 738. Fl. succ. 656.*

Vicia segetum cum filiquis plurimis hirsutis. *Bauh. pin. 345.*

Vicia sylvestris seu Cracca minima. *Ger. Em. 1028.*

Aracus f. Cracca minor. *Park. 1070.*

Cracca minor. *Riv. tetr. ic. R. Syn. 322.*

Anglis, hairy Tare.

Habitat *cum priore. ☉. VI.*

ORNITHOPUS. 790.

Legumen articulatum, teres, arcuatum.

perpusillus. 1. ORNITHOPUS foliis pinnatis, leguminibus subarticulatis. *Sp. pl. 743.*

Ornithopus foliis pinnatis, articulis leguminum subrotundis lævibus. *Hort. cliff. 364.*

Ornithopodium minus. *Ger. Em. 1241. Bauh. pin. 350.*

Ornithopodium radice nodosa. *Park. 1093.*

Ornithopodium radice tuberculis nodosa. *Bauh. pin. 350.*

Ornithopodium majus. *Bauh. pin. 350. Park. 1093.*

Anglis, Bird's-foot.

Habitat *in glareosis. 4. V.—VIII.*

HIPPOCREPIS. 791:

Legumen compressum, altera sutura pluries emarginatum, curvum.

1. HIP-

1. HIPPOCREPIS leguminibus pedunculatis confertis : *comosa*,
margine exteriori repandis. *Roy. lugdb.* 384. *Sp.*
pl. 744.

Ferrum equinum germanicum filiquis in summitate.
Bauh. pin. 349. *R. Syn.* 325.

Ferrum equinum comosum. *Park.* 1091.

Hedysarum glycyrrhizatum. *Ger. Em.* 1233.

Anglis, tufted Horfeshoe-vetch.

Habitat in pratis et pascuis solo cretaceo. On the chalky
grounds about Gogmagog hills, and elsewhere in
Cambridgeshire ; also on the northern mountains. In
collibus cretaceis inter Northfleet et Gravesend. *R.*
Syn. ☉. VII.

HEDYSARUM. 793.

Cor. carina transverse obtusa. *Legumen* articulis
monospermis.

1. HEDYSARUM foliis pinnatis, leguminibus monof- *Onobry-*
permis aculeatis, corollarum alis brevissimis, caule *chis.*
elongato. *Syst. nat.* 1171. *Sp. pl.* 751.

Onobrychis f. *Caput Gallinaceum.* *Ger. Em.* 1243.
R. Syn. 327.

Onobrychis vulgaris. *Park.* 1082.

Onobrychis foliis viciæ, fructu echinato major. *Bauh.*
pin. 350.

Polygalon Gefneri. *Bauh. hist. II.* 335.

Anglis, St. Foin, or Cock's-head.

Habitat in pratis et pascuis cretaceis. 4 VII.

ASTRAGALUS. 799.

Legumen biloculare, gibbum.

1. ASTRAGALUS caulescens prostratus, leguminibus *glycyphyl-*
subtriquetris arcuatis, foliis ovalibus pedunculo *los.*
longioribus. *Sp. pl.* 758. *Fl. suec.* 658.

Astragalus luteus perennis procumbens, vulgaris f.
sylvestris. *Hist. Ox. II.* 107.

Glaux vulgaris f. *Glycyrrhiza sylvestris.* *Park.* 1098.

Glycyrrhiza sylvestris floribus luteo pallescentibus.
Bauh. pin. 352.

Hedysarum glycyrrhizatum. *Ger. Em.* 1233.

Anglis,

DIADELPHIA DECANDRIA.

Anglis, wild Liquorice, or Liquorice Vetch.

Habitat in pratis, pascuis, et ad sepes. Circa Charle-
ton in Cantio copiose. 4. VI. VII.

arenarius. 2. ASTRAGALUS subcaulescens procumbens, floribus
subracemosis erectis, foliis tomentosis. *Sp. pl.* 759.
Fl. suec. 659.

Astragalus pedunculis folia villosa æquantibus, flori-
bus erectis, caulibus procumbentibus. *It. scan.* 108.

Astragalus incanus parvus purpureus nostras. *Pluk.*
alm. 59. *R. Syn.* 326. t. 12. f. 3.

Glaux exigua montana purpurea nostras. *R. Syn.*
ed. II. 192.

Anglis, Purple Mountain Milkwort.

Habitat in pratis et pascuis arenosis. On Gogmagog
hills, Newmarket heath, Royston heath, &c. About
Hafelwood, Yorkshire; Nullibi copiosius quam secus
viam agrorum Pigburnensium, non procul Doncas-
tria agri Eboracensis, D. Richardson. *R. Syn.* 4.
VI. VII.

T R I F O L I U M. 802.

Flores subcapitati. Legumen vix calyce longius, non
dehiscens, deciduum.

* *Meliloti leguminibus nudis polyspermis.*

m. offici. 1. TRIFOLIUM leguminibus racemosis nudis disper-
mis, caule erecto. *Hort. cliff.* 375. *Sp. pl.* 765.
Fl. suec. 663.

Melilotus officinarum germaniæ. *Bauh. pin.* 331.

Melilotus vulgaris. *Park.* 719. *R. Syn.* 331.

Melilotus germanica. *Ger. Em.* 1205.

Anglis, common Melilot.

Habitat in arvis et ad sepes. 8. VI, VII.

Pharmac. Meliloti Herba, Flores.

ornithopo- 2. TRIFOLIUM leguminibus nudis octospermis subter-
ides. nis calyce duplo longioribus, caulibus declinatis.
Sp. pl. 766.

Medicago leguminibus ternatis erectis recurvis de-
scendentibus, pedunculo communi. *Hort. cliff.* 376.

Trifo-

Trifolium filiquis ornithopodii nostras. *R. Syn. ed. II.*
195.

Trifolium filiquosum loto affine filiquis ornithopodii.
Pluk. ph. t. 68. f. 1.

Fœnumgræcum humile repens, ornithopodii filiquis
brevibus erectis. *R. Syn. 331.*

Anglis, Birdsfoot Trefoil.

Habitat in pratis et pascuis arenosis. Found by Mr. Th.
Willisell among the corn, half a mile on this side Tad-
caster, towards Sherborn, as one goeth up the hill;
also near Oxford, in the like grounds. Mr. New-
ton, in our company, found it on the sandy banks by
the sea-side, at Tottlesbury in Essex, plentifully.
Mr. Doody in Tothil-fields, Westminster. *R. Syn.*
©. VI.

** Lotoidea leguminibus testis polyspermis.

3. TRIFOLIUM capitulis umbellaribus, leguminibus repens.
tetraspermis, caule repente. *Sp. pl. 767. Fl. suec.*
665.

Trifolium capitulis subrotundis, flosculis pedunculatis,
leguminibus tetraspermis, caule procumbente. *Hort.*
cliff. 375.

Trifolium pratense album. *Baub. pin. 327. Park.*
1110. *R. Syn. 327.*

Trifolium minus pratense, flore albo. f. 2. *Ger. Em.*
1185.

- β Trifolium capitulis umbellaribus, leguminibus, tetra- hybridum
spermis, caule adscendente. *Sp. pl. 766. Fl. suec.*
664.

Anglis, white Trefoil, or Dutch Clover.

Habitat in pratis et pascuis ubique, β in pascuis humidis
prope Peckham. 4. V.—IX.

*** Lagopoda calycibus villosis monospermis.

4. TRIFOLIUM spicis globosis, corollis monopetalis, ochroleu-
calyculum infimo dente longissimo erecto, foliis cau- con.
leque hirsutis.

Trifolium pratense hirsutum majus, flore albo-sul-
phureo, seu ὀχρολεύκω. *R. Syn. 328.*

Trifolium

Trifolium lagapoides annuum hirsutum pallide luteum
feu ochroleucum. *Hist. ox. II.* 141. *f.* 2. *t.* 12.
f. 12.

Anglis, yellow flowered Trefoil.

Habitat in pratis et pascuis siccioribus, in comitatibus
Essexiensis, Cantabrigiensis, frequens. δ . V. VI.

pratense. 5. TRIFOLIUM spicis subvillosis cinctis stipulis oppositis membranaceis, corollis monopetalis. *Sp. pl.* 768. *Fl. suec.* 666.

Trifolium spicis villosis, caule diffuso, foliolis integerrimis. *Hort. cliff.* 375.

Trifolium pratense. *Ger. Em.* 1185. purpureum.
Bauh. pin. 327. *R. Syn.* 328. vulgare. *Park.* 1110.

β *Trifolium pratense* purpureum minus, foliis cordatis.
R. Syn. 328. *t.* 13. *f.* 1.

γ *Trifolium purpureum majus sativum*, *pratense* simile.
R. Syn. 328.

Anglis, purple or Honeysuckle Trefoil, γ Clover.

Habitat in pratis et pascuis. α . V.—IX.

Pharmac. Trifolii pratensis Herba.

medium. 6. TRIFOLIUM spicis villosis globosis terminalibus, corollis monopetalis foliolis lanceolatis subintegerrimis, caule erecto.

Trifolium purpureum majus, foliis longioribus et angustioribus, floribus saturatioribus. *R. Syn.* 328.

Trifolium montanum purpureum majus. *Bauh. pin.* 328.

Anglis, long-leaved purple Trefoil.

Habitat in pratis et pascuis montosis. α . VII.

maritimum. 7. TRIFOLIUM spicis villosis globosis, calycibus patulis æqualibus, caule ascendente foliolis cuneiformibus hirsutis.

Trifolium stellatum glabrum *Ger. Em.* 1208. *Pluk. ph.* 113 *f.* 4. *R. Syn.* 329.

Anglis, leaf-headed Trefoil

Habitat in pratis maritimis. Observed by Dr. Johnson in Dartford salt marsh; found it by the water-side at Lighe, and at Little-holland in Essex; at Tilbury-fort, Mr. Petiver; plentifully about Sheerness,

nefs, Mr. Doody. *R. Syn. prope Bristolium copiose;*
in pratis inter Greenhithe et Northfleet in Cantio.
4. VII.

8. TRIFOLIUM spicis villosis ovalibus, dentibus ca- *arvense.*
lycinis setaceis æqualibus. *Sp. pl. 769. Fl. suec.*
668.

Trifolium arvense humile spicatum f. Lagopus. *Bauh.*
pin. 328. R. Syn. 330.

Lagopus vulgaris. *Park. 1107.*

Lagopodium sive Pes leporis. *Ger. Em. 1192.*

- β Lagopus perpuillus supinus perelegans maritimus.
Ph. br. R. Syn. 330.

Anglis, Hare's-foot Trefoil.

Habitat in pascuis arenosis et arvis. ☉. VII. VIII.

9. TRIFOLIUM capitulis sessilibus sublateralibus ova- *striatum.*
tis, calycibus striatis rotundatis. *Sp. pl. 770. Fl.*
suec. 669.

Trifolium parvum hirsutum, floribus parvis dilute pur-
pureis in glomerulis mollioribus et oblongis, semine
magno. *R. Syn. 329. t. 12. f. 3.*

Anglis, knotted Trefoil.

Habitat in pascuis siccioribus. ☉. VI.

10. TRIFOLIUM capitulis sessilibus globosis, calyci- *glomeratum.*
bus striatis patulis æqualibus. *Sp. pl. 770.*

Trifolium cum glomerulis ad caulium nodos rotundis.
R. Syn. 329.

Anglis, round-headed Trefoil.

Habitat in pratis et pascuis arenosis. On Blackheath,
and near Greenhithe in Kent, Mr. Doody. *R. Syn.*
☉. VI.

11. TRIFOLIUM capitulis sessilibus ovatis, calycibus *scabrum.*
villosis inæqualibus. *Sp. pl. 770.*

Trifolium capitulo oblongo aspero. *Bauh. pin. 329.*
pr. 140.

Trifolium cujus caules ex geniculis glomerulos ob-
longos proferunt. *Bauh. hist. II. 378.*

Trifolium

Trifolium flosculis albis, in glomerulis oblongis asperis, cauliculis proxime adnatis. R. Syn. 329. Vaill. parif. 169. t. 33. f. 1.

Anglis, Oval-headed Trefoil.

Habitat in collibus cretaceis juxta Thamefin, inter Northfleet et Gravesend. R. Syn. in insula Shepey copiose. ☉. V. VI.

subterra- 12. TRIFOLIUM capitulis villosis subquinquefloris, in-
neum. *volucro centrali reflexo rigido fructum involvente. Hort. cliff. 374. Roy. lugdb. 379. Dalib. parif. 224. Sp. pl. 769.*

Trifolium pumilum supinum flosculis longis albis. Ph. br. R. Syn. 327.

Trifolium album tricoccum subterraneum reticulatum. Hist. ox. II. 138. f. 2. t. 14. f. 5.

Anglis, dwarf Trefoil.

Habitat in pascuis sterilioribus circa Londinum frequens. ☉. V.

**** *Vesicaria calycibus inflatis ventricosis.*

fragife- 13. TRIFOLIUM capitulis subrotundis. calycibus in-
rum. *flatis bidentatis reflexis, caulibus repentibus. Hort. cliff. 373. Fl. suec. 670. Sp. pl. 772.*

Trifolium fragiferum. Ger. Em. 1208. R. Syn. 329. friscum. Bauh. pin. 329. Clus. cur. app. 38. t. 39. Park. 1109.

β Trifolium fragiferum purpureum folio oblongo. Hist. ox. II. 144.

Anglis, Strawberry Trefoil.

Habitat in pratis et pascuis humidis. Ⅵ. VIII.

**** *Lupulina vexillis corollæ reflexis.*

agrarium. 14. TRIFOLIUM spicis ovalibus imbricatis, vexillis
deflexis persistentibus, calycibus nudis, caule erecto. Sp. pl. 772. Fl. suec. 671.

Trifolium pratense luteum, capitulo lupuli, vel agrarium. Bauh. pin. 328. R. Syn. 330.

Trifolium luteum minimum. Ger. Em. 1186. Lob. ii. 29. Park. III.

Anglis,

Anglis, Hop Trefoil.

Habitat in pratis glareosis. ☉. VI.

15. TRIFOLIUM spicis ovalibus imbricatis, vexillis procumbentibus, caulibus procumbentibus. *bens*.

Sp. pl. 772. *Fl. suec.* 673.

Trifolium lupinum alterum minus. *R. Syn.* 330. *t.*

14. *f.* 3.

Anglis, decumbent Trefoil.

Habitat in pascuis et pratis. ☿. V.—VIII.

16. TRIFOLIUM spicis subimbricatis, vexillis deflexis *filiforme*.
persistentibus, calycibus pedicellatis. caulibus procumbentibus. *Sp. pl.* 773. *Fl. suec.* 674.

Trifolium lupulinum minimum. *R. Syn.* 331. *t.* 14.
f. 4.

Anglis, small Trefoil.

Habitat in pratis arenosis passim. ☉. V. VI.

M E D I C A G O. 805.

Legumen compressum cochleatum. Carina corollæ
a vexillo deflexens.

1. MEDICAGO pedunculis racemosis. leguminibus *lufalcata*.
natis caule prostrato. *Sp. pl.* 779.

Medicago pedunculis laxè spicatis leguminibus contortis caule erecto glabris. *Hort. cliff.* 377.

Medica sylvestris. *Bauh. hist. II.* 383. *R. Syn.* 333.

Medica frutescens flavo flore Clusii. *Park.* 1114.

Trifolium luteum filiqua cornuta. *Ger. Em.* 1191.

Trifolium luteum sylvestre, filiqua cornuta et medica frutescens. *Bauh. pin.* 330.

Anglis, yellow Medick.

Habitat in aggeribus agrorum et ad vias in Norfolcia;
between Bushyhill and Watford, the footway, Mr.
Doody. *R. Syn.* ☿. VII.

- *2. MEDICAGO pedunculis racemosis, leguminibus *fativa*.
contortis, caule erecto glabro. *Hort. cliff.* 377. *Sp.*
pl. 778.

Medica legitima. *Clus. hist. II.* 242.

Medica fativa. *Hist. ox. II.* 150. *f.* 2. *t.* 16. *f.* 2.

Foenum

Foenum buigundiacum. *Loc. ic.* 2. p. 36.

Anglis, Lucern.

Habitat in pratis et pascuis. 4. VI. VII.

- lupulina.* 3. MEDICAGO spicis ovalibus, leguminibus reniformibus monospermis, caule procumbente. *Sp. pl.* 779. *Fl. suec.* 678.
 Medica polycarpus fructu minore compresso scabro. *R. Syn.* 333.
 Trifolium luteum lupulinum. *Ger. Em.* 1186. *R. Syn.* 331.
 Trifolium montanum lupulinum. *Park.* 1105.
 Trifolium pratense luteum, capitulo brevior. *Bauh. pin.* 328.
Anglis, Melilot Trefoil.
 Habitat in arvis, pratis et pascuis arenosis. ☉. V.—VIII.

- arabica.* 4. MEDICAGO pedunculis racemosis, leguminibus cochleatis spinosis, caule diffuso. *Sp. pl.* 780.
 Medica Arabica Camerarii, five Trifolium cordatum. *Park.* 1115.
 Trifolium cochleatum folio cordato maculato. *Bauh. pin.* 329. *R. Syn.* 333.
 Trifolium cordatum. *Ger. Em.* 1190.
 β Medica echinata minima. *Bauh. hist. II.* 386.
 γ Trifolium cochleatum modiolis spinosis. *R. Syn.* 333.
 δ Medica marina supina nostras, foliis viridibus ad summos ramulos villosis. *Pluk. alm.* 245.
Anglis, Heart Trefoil, or Claver.
 Habitat in pratis, pascuis, et inter segetes. ☉. V.

L O T U S. 803.

Legumen cylindricum, strictum. *Alæ* sursum longitudinaliter conniventes. *Cal.* tubulosus.

- corniculata.* 1. LOTUS capitulis depressis, caulibus decumbentibus, leguminibus cylindricis patentibus. *Sp. pl.* 775. *Fl. suec.* 675.
 Lotus corniculata glabra minor. *Bauh. hist. II.* 356. *R. Syn.* 334.

Lotus

Lotus f. melilotus pentaphyllos minor glabra. *Bauh. pin.* 332.

Trifolium filiquosum minus. *Ger. Em.* 1191.

β Lotus pentaphyllos minor angustioribus foliis fruticosior. *R. Syn.* 334.

γ Lotus pentaphyllos flore majore luteo splendente. *Bauh. pin.* 332.

δ Lotus corniculata major, minus hirsuta. *R. Syn.* 334.

ε Lotus corniculata minor, foliis subtus incanis. *R. Syn.* 334.

Anglis, Bird's-foot Trefoil.

Habitat in pratis pascuis et ad sepes. γ in sylvis et sepibus humidis. 24. VI.—VIII.

U

Classis

Classis XVIII.

POLYADELPHIA
POLYANDRIA.

HYPERICUM. 808.

Cal. quinquepartitus. *Petala* quinque. *Filamenta* multa, in quinque phalanges.

- perfora-*
tum. 1. HYPERICUM floribus trigynis, caule ancipiti, foliis obtusis pellucido-punctatis. *Hort. cliff.* 380. *Sp. pl.* 785. *Mat. med.* 368. *Fl. succ.* 681.
Hypericum. *Ger. Em.* 540. *R. Syn.* 342. vulgare. *Baub. pin.* 279. *Park.* 572.
Hypericum vulgare f. perforata, caule rotundo, foliis glabris. *Baub. hist.* III. 381.
Anglis, St John's Wort.
Habitat in dumetis et ad sepes. 4. VII.
Pharmac. Hyperici Herba Flores Semina.
- humif-*
sum, 2. HYPERICUM floribus trigynis axillaribus solitariis, caulibus ancipitibus filiformibus, foliis glabris. *Syst. nat.* 1184.
Hypericum floribus trigynis, caulibus ancipitibus prostratis. *Sp. pl.* 785. *Fl. succ.* 681.
Hypericum minus supinum *Park* 572. *R. Syn.* 343.
Hypericum supinum glabrum. *Ger. Em.* 541. *Baub. pin.* 279.
Anglis, trailing St. John's Wort.
Habitat in pratis et pascuis glareosis. 4. VII.
- fulchrum.* 3. HYPERICUM floribus trigynis, calycibus ferrato-glandulosis, foliis cordatis glabris, bracteis integerrimis, caule tereti. *Syst. nat.* 1184. *Sp. pl.* 786.
Hypericum foliis cordatis connatis, laciniis calycinis ovatis obtusis. *Guett. stamp.* II. 175.

Hypericum

- Hypericum foliorum basi latissima, foliis calycis brevioribus obtusis. Hall. herb. 361.*
Hypericum minus erectum. Bauh. pin. 279.
Hypericum pulchrum Tragi. Bauh. hist. III. 383.
R. Syn. 342.
Hypericum quintum s. pulchrum Tragi. Ger. Em. 540.
Anglis, upright St. John's Wort.
Habitat in sylvis, sepibus et ericetis. 4. VII.
4. **HYPERICUM** floribus trigynis, fructibus baccatis, *Androsæ-*
 caule fruticoso ancipiti. *Sp. pl. 785. mum.*
Hypericum floribus trigynis, fructu baccato, foliis
ovatis pedunculo longioribus. Hort. cliff. 380.
Hypericus maximum Androsæmum vulgare dictum.
R. Syn. 343.
Androsæmum vulgare. Park. 575.
Androsæmum maximum frutescens. Bauh. pin. 280.
Clymenum Italarum. Ger. Em. 548.
Anglis, Tutfan, or Park-leaves.
Habitat in sylvis et sepibus humidis frequens. 4. VII.
5. **HYPERICUM** floribus trigynis, calycibus ferrato- *montanum*
 glandulosis, caule erecto, foliis ovatis glabris. *Fl.*
succ. 683.
Hypericum elegantissimum non ramosum folio lato.
Bauh. hist. III. 383. R. Syn. 343.
Androsæmum campoclarense. Col. ecphr. I. 73. t. 74.
Afcyron s. Hypericum bifolium glabrum non perfora-
tum. Bauh. pin. 280.
Anglis, imperforate St. John's Wort.
Habitat in sylvis et dumetis montosis. In Charleton
wood. Blackf. spec. bot. 4. VII.
6. **HYPERICUM** floribus trigynis, calycibus ferrato- *hirsutum.*
 glandulosis, caule erecto, foliis ovatis pubescenti-
 bus. *Sp. pl. 786. Fl. succ. 682.*
Hypericum majus s. Androsæmum matthioli. R. hist.
100.
Androsæmum hirsutum. Bauh. pin. 280.
Androsæmum Afcyron dictum, caule rotundo hirsu-
to. Hist. ox. II. 271. f. 5. t. 6. f. 11.
Anglis, Tutfan, or hairy St. John's Wort.
Habitat in dumetis et sepibus. 4. VII.

- quadran-*
gulum. 7. **HYPERICUM** floribus trigynis, caule quadrato herbaceo *Sp. pl.* 785. *Fl. suec.* 679
Hypericum Ascyron dictum, caule quadrangulo.
Bauh. hist. III. 382 *R. Syn.* 344.
Ascyron. *Ger. Em.* 542. *Dod. pempt.* 78. vulgare.
Park. 575.
Anglis, St. Peter's Wort.
Habitat in sepibus humidis frequens. 24. VII.
- elodes.* 8. **HYPERICUM** floribus trigynis, foliis rotundis tomentosis, caule repente.
Ascyron supinum villosum palustre. *Bauh. pin.* 280.
Park. 574. *R. Syn.* 344.
Ascyron supinum *ἐλαδης* Clusii. *Ger. Em.* 542.
Anglis, Marsh St. Peter's Wort.
Habitat in palustribus solo putrido aut spongioso. In
Sherley common inter Croydon et Wickham copiose.
In ericeto Hamstediano. 24. VII.

Classis

Classis XIX.

SYNGENESIA:

POLYGAMIA ÆQUALIS.

* *Semiflosculosi Tournef. corollis ligulatis omnibus.*

TRAGOPOGON. 810.

Receptaculum nudum. Cal. simplex, polyphyllus.

Pappus plumosus.

1. TRAGOPOGON calycibus corollæ radium æquantibus, foliis integris strictis. *Sp. pl.* 789. *Fl. suec.* 685. *pratense.*

Tragopogon luteum. Ger. Em. 735. *Park.* 412. *R. Syn.* 171.

Tragopogon pratense luteum majus. Bauh. pin. 274. *Anglis, yellow Goat's-beard.*

Habitat in pratis et pascuis. ♂. VI.

2. TRAGOPOGON calycibus corollæ radio longioribus, foliis integris nudis, pedunculis superne incrassatis. *Syst. nat.* 1191. *Sp. pl.* 789. *porrifolium.*

Tragopogon purpureo-cæruleum, porrifolio, quod Artificivulgo. Bauh. pin. 274. *Hist. ox. III.* 8. *f. 1. t.* 9. *f. 5.*

Tragopogon purpureum. Ger. Em. 735. *Park.* 412. *R. Syn.* 172.

Anglis, purple Goat's-beard.

Habitat in pratis et pascuis. In many places of Cornwall, Dr. Gunthorp; In the fields about Carlisle and Rose-Castle, Cumberland. R. Syn. In the marshes near Long-Reach, below Woolwich, Mr. Jarvis. In some meadows near Edmonton, Mr. Hurlock. Blackf. spec. bot. ♂. VI.

P I C R I S. 812.

Recept. nudum. *Cal.* caliculatus. *Pappus* plumosus. *Sem.* transversim sulcata.

Echioides. 1. PICRIS perianthiis involucro maximo cinctis. *Hort. upf.* 241. *Sp. pl.* 792.

Picris calycibus simplicibus, perianthio maximo obvallatis. *Hort. cliff.* 387. *Roy. lugdb.* 128.

Hieracium echioides, capitulis cardui benedicti. *Baub. pin.* 128. *R. Syn.* 166.

Buglossum luteum. *Ger. Em.* 798. f. Lingua bovis. *Park.* 800.

Anglis, Ox's Tongue.

Habitat in marginibus agrorum. ☉. VII. VIII.

Hieracides. 2. PICRIS perianthiis imbricatis. *Hort. cliff.* 387. *Lech. scan.* 62. *Sp. pl.* 792. *Fl. suec.* 686.

Hieracium asperum, majore flore, in agrorum limitibus. *Baub. hist. II.* 1029. *R. Syn.* 167.

Hieracium asperum. *Ger. Em.* 298.

Cichorium pratense luteum hirsutiae asperum. *Baub. pin.* 126.

Cichorium pratense luteum asperum. *Park.* 777.

Anglis, yellow Succory, or rough Hawkweed.

Habitat in agrorum marginibus. ☉. VII.

S O N C H U S. 813.

Recept. nudum. *Cal.* imbricatus, ventricosus. *Pappus* plumosus.

oleraceus. 1. SONCHUS pedunculis tomentosis, calycibus lævibus. *Sp. pl.* 794. *Fl. suec.* 688.

Sonchus caule ramosissimo diffuso foliis summis amplexicaulibus, pedunculis inæqualibus. *Roy. lugdb.* 129

α Sonchus lævis. *Ger. Em.* 292. *R. Syn.* 162. vulgaris. *Park.* 805.

Sonchus lævis laciniatus latifolius. *Baub. pin.* 124.

β Sonchus lævis minor paucioribus laciniis. *Baub. pin.* 124.

γ Sonchus asper laciniatus. *Baub. pin.* 124. *Park.* 804.

δ Sonchus asper non laciniatus. *Baub. pin.* 123.

ε Sonchus

2. *Sonchus subrotundo folio nostras, lævissimis spinulis circa foliorum oras exasperatus. Pluk. alm. 354. t. 61. f. 5.*
Sonchus ἀφυλλόκαυλος angusto et oblongo folio nostras, per foliorum ambitum creberrimis spinulis asperatus. Pluk. ph. t. 62. f. 4.
Sonchus asper dentatus. Pet. h. br. t. 14. f. 4.
Sonchus lævis laceratus. Pet. h. br. t. 14. f. 10.
Anglis, common Sow-thistle.
Habitat in rudèratis, cultis, arvis. ☉. VI.—VIII.
Pharmac. Sonchi Herba.
2. **SONCHUS** pedunculis calycibusque hispìdis subum-arvensis. bellatis, foliis basi cordatis. *Sp. pl. 793. Fl. suec. 687.*
Sonchus foliis lanceolato-oblongis dentatis, floribus congestis hispìdis. Hort. cliff. 384.
Sonchus repens, multis Hieracium majus. Baub. hist. II. 1070. R. Syn. 163.
Sonchus arborescens. Ger. Em. 294.
Hieracium majus folio Sonchi. Baub. pin. 126.
Anglis, Tree Sow-thistle.
Habitat in arvis et ad sèpes. 4. VIII.
3. **SONCHUS** pedunculis calycibusque hispìdis subum-palustris. bellatis, foliis basi sagittatis. *Sp. pl. 793.*
Sonchus caule erecto, foliis pinnato-hastatis apiceque hastatis, floribus congestis villosis. Hort. cliff. 385.
Sonchus tricubitalis, folio cuspidato. Merr. pin. R. Syn. 163.
Sonchus arborescens alter. Ger. Em. 294.
Sonchus asper arborescens. Baub. pin. 124.
Anglis, Marsh Sow-thistle.
Habitat ad ripas Thamefis fluvii non procul Greenico, et circa Blackwall, R. Syn. 4. VIII.

L A C T U C A. 814.

Recept. nudum. Cal. imbricatus, cylindricus. Papus simplex, stipitatus.

1. **LACTUCA** foliis verticalibus carina aculeatis. *Syst. virofa. nat. 1193.*

SYNGENESIA : POLYGAMIA ÆQUALIS.

Lactuca sylvestris major odore opii. *Ger. Em.* 309.

R. Syn. 161.

Lactuca sylvestris odore viroso. *Bauh. pin.* 123.

Lactuca viroso. *Park.* 813.

β *Lactuca sylvestris* costa spinosa. *Bauh. pin.* 123.

γ *Lactuca sylvestris* folio non laciniato. *R. Syn.* 162.

Anglis, strong-scented wild Lettuce.

Habitat ad sepes et agrorum margines. ♂. VII. VIII.

saligna. 2. **LACTUCA** foliis hastato-linearibus sessilibus carina aculeatis. *Amoen. acad.* 1. 150. *Sp. pl.* 796.

Lactuca foliis imis pinnatifidis, cæteris ensiformibus basi hastatis, dorso ferratis. *Sauv. monsp.* 692.

Lactuca foliis radicalibus linearibus dentato-pinnatis : laciniis sursum dentatis : caulinis alato-lanceolatis, floribus linearibus sessilibus. *Guet. stamp.* 374.

Lactuca sylvestris minima. *R. cat. cant. syn.* 162.

Chondrilla viscosa humilis. *Bauh. pin.* 130. *pr.* 68.

Park. 783. *Ger. Em.* 287.

Anglis, the least wild Lettuce.

Habitat in aggeribus fossarum et pascuis solo cretaceo passim. ○. VIII.

P R E N A N T H E S. 816.

Recept. nudum. *Cal.* calyculatus. *Pappus* simplex, subseffilis, *Flosculi* simplici serie.

muralis. 1. **PRENANTHES** flosculis quinis, foliis lyrato-hastatis. *Hort. cliff.* 383 *Fl suec.* 692. *Sp. pl.* 797.

Lactuca sylvestris murorum flore luteo. *Bauh. hist.* II.

1004. *R. Syn.* 162.

Sonchus lævis muralis. *Ger. Em.* 293.

Sonchus lævis alter parvis floribus. *Park.* 805.

Sonchus lævis laciniatus muralis parvis floribus. *Bauh. pin.* 124.

Anglis, Ivy-leaved wild Lettuce.

Habitat in muris et sylvis umbrosis. Prope Hornsey. 4. VII.

L E O N T O D O N. 817.

Recept. nudum. *Cal.* imbricatus, squamis laxiusculis. *Pappus* plumosus.

I. L E.

1. LEONTODON calyce inferne reflexo. *Fl. lapp. Taraxa-*
280. *Mat. med.* 368. *Sp. pl.* 798. *Fl. suec.* 693. *cum.*
Dens leonis. *Ger. Em.* 290. *R. Syn.* 171. vulgaris.
Park. 780.
Dens leonis latiore folio. *Bauh. pin.* 126.
β Dens leonis angustiore folio. *Bauh. pin.* 126.
γ Dens leonis montanus angustifolius. *R. Syn.* 171.
Hieracium montanum angustifolium nonnihil incanum.
Bauh. pin. 129.
Anglis, Dandelion.
Habitat in pratis et pascuis. 4. IV.—VI.
Pharmac. Taraxaci Radix, Herba.
2. LEONTODON calyce toto erecto hispido, foliis *hispidum.*
hispidis dentatis integerrimis; setis furcatis. *Syst.*
nat. 1194. *Fl. suec.* 694. *Sp. pl.* 799.
Dens leonis hirsutus λεπτόκαυλον, Hieracium dictus.
R. Syn. 171.
Hieracium Dentis leonis folio hirsutum. *Ger. Em.*
303.
Hieracium asperum flore magno Dentis leonis. *Bauh.*
pin. 127.
Hieracium asperum foliis et floribus dentis leonis bul-
bosi. *Park.* 788.
β Crepis foliis lanceolatis dentato-sinuatis hispidis: setis
subulatis, scapo unifloro. *Sp. pl.* 805.
Hieracium pumilum saxatile asperum premorsa radice.
Bauh. pin. 128. *pr.* 66. *ic.* *R. Syn.* 167.
Anglis, rough Dandelion.
Habitat in pratis et pascuis. 4. V. VI.
3. LEONTODON foliis glabris dentatis integerrimis, *autumnale*
calycibus erectis, pedunculis squamosis. *Sp. pl.* 798.
Fl. suec. 695.
Crepis foliis longis dentatis linearibus, caule declinato
subnudo. *Hort. cliff.* 389.
Hieracium minus præmorsa radice. *Park.* 794. *R. Syn.*
165.
Hieracium minus f. leporinum. *Ger. Em.* 296.
Hieracium chondrillæ folio glabro, radice succisa,
majus. *Bauh. pin.* 127.

β Hieracium

- β Hieracium præmorsum laciniatum. *Pet. h. br. t.*
12. f. 2.
Anglis, yellow Devil's-bit.
Habitat in pratis et pascuis. 4. VIII.

HIERACIUM. 818.

Recept. nudum. Cal. imbricatus, ovatus. Pappus simplex, fessilis.

* *Scapo nudo unifloro.*

alpinum. 1. **HIERACIUM** foliis oblongis integris dentatis, caule subnudo unifloro, calyce piloso. *Sp. pl. 800. Fl. suec. 696.*

Hieracium alpinum pumilum, folio lanuginoso. *Bauh. pin. 129.*

Hieracium villosum alpinum, flore magno singulari. *R. Syn. 169. t. 6. f. 2.*

Anglis, Mountain Hawkweed.

Habitat in editioribus Venedotiæ et Provisiæ rupibus passim. D. Lhwyd. R. Syn. 4. VII.

Pilosella. 2. **HIERACIUM** foliis integerrimis ovatis subtus tomentosis, caule repente, scapo unifloro. *Sp. pl. 800. Fl. suec. 697.*

Pilosella repens. Ger. Em. 638. R. Syn. 170.

Pilosella minor vulgaris repens. Park. 690.

Pilosella major repens hirsuta. Bauh. pin. 262.

Anglis, common creeping Mouse-ear.

Habitat in pratis et pascuis siccissimis frequens. 5. V.

Pharmac. Pilosellæ s. Auriculæ muris Herba.

** *Scapo nudo multifloro.*

dubium. 3 **HIERACIUM** foliis integris ovato-oblongis, stolone repente, scapo nudo multifloro. *Sp. pl. 800. Fl. suec. 698.*

Hieracium foliis virentibus pilosis ovato-lanceolatis, scapo simplici multifloro. *Hall. helv. 743.*

Hieracium piloselloides vulgare. *Vaill. æt. Gal. 1721. p. 238.*

Pilosella major repens minus hirsuta. Bauh. pin. 262.

Anglis, Mouse-ear Hawkweed.

Habitat

Habitat in monte Fairfield dicta, prope Rydal in comitatu Westmorlandico. 4. VII. VIII.

4. HIERACIUM foliis integerrimis lanceolatis, scapo *Auricula*. nudo multifloro. *Fl. lapp.* 289. *Sp. pl.* 800. *Fl. suec.* 699

Hieracium foliis integerrimis, caule repente, scapo nudo multifloro. *Hort. cliff.* 388.

Pilosella major erecta altera. *Bauh. pin.* 262.

Pilosella repens minor, caule pedali, polyanthos, foliis angustis oblongis. *Raj. app.* 147.

Anglis, narrow-leaved Hawkweed.

Habitat in pratis montosis supra Dalehead, non longe a Grassmere in Westmorlandia, sed sparsim. 4. VII.

Folia lanceolata, acuta subintegerrima. Scapus erectus nudus, vix semipedalis pilis minimis adpersus, Panicula a 3. ad 6. floribus luteis. Calycis foliola linearipilosa.

*** *Caule folioso.*

5. HIERACIUM caule ramoso, foliis radicalibus ovatis *murorum*. dentatis, caulino minore. *Hort. cliff.* 388. *Sp. pl.* 802. *Fl. suec.* 701.

Hieracium caule ramoso, foliis dentatis. *Fl. lapp.* 284.

Hieracium murorum foliis pilosissimo. *Bauh. pin.* 129. *R. Syn.* 168.

Hieracium murorum Bauhini, quod est Pulmonaria Gallorum Lobelii. *Park.* 801.

Pulmonaria gallica f. aurea latifolia. *Ger. Em.* 304.

- β Hieracium Pulmonarium dictum angustifolium. *R. Syn.* 168.

Pulmonaria gallica f. aurea angustifolia. *Ger. Em.* 304.

- γ Hieracium macrocaulon hirsutum folio rotundiore. *R. Syn.* 169.

Hieracium fruticosum folio subrotundo. *Bauh. pin.* 126.

- δ Hieracium λεπτόκαυλον hirsutum folio longiore. *R. Syn.* 169.

- ε Pilosellæ majoris seu Pulmonariæ luteæ species angustifolia. *Bauh. hist. II.* 1034.

Anglis,

Anglis, French or golden Lungwort.

Habitat in sylvis, muris antiquis et aggeribus umbrosis passim. 4. VII.

paludosum 6. **HIERACIUM** caule paniculato, foliis amplexicaulibus dentatis glabris, calycibus hispidis. *Sp. pl.* 803. *Fl. suec.* 702.

Hieracium foliis amplexicaulibus minute dentatis glabris. *Hall. herb.* 748.

Hieracium montanum latifolium glabrum minus. *Bauh. pin.* 129. *Hist. ox.* III. 69. f. 7. t. 5. f. 47.

Hieracium montanum cichorei folio. *R. Syn.* 166.

Anglis, Succory-leaved Hawkweed.

Habitat in pratis humidis et juxta rivulos Cravoniæ montosos abundat. *R. Syn.* 8. VII.

sabaudum. 7. **HIERACIUM** caule erecto multifloro, foliis ovato-lanceolatis dentatis femiamplexicaulibus. *Roy. lugdb.* 124. *Sp. pl.* 804. *Fl. suec.* 703.

Hieracium fruticosum latifolium hirsutum. *Bauh. pin.* 129. *Park.* 802. *R. Syn.* 167.

β Hieracium fruticosum latifolium glabrum. *Park.* 168.

γ Hieracium s. Pilosellæ majoris species humilis, foliis longioribus rarius dentatis plurimis simul, flore singulari nostras. *R. Syn.* 170. *Pluk. ph.* t. 37. f. 3. *Anglis*, broad-leaved bushy Hawkweed.

Habitat in sylvis et sepibus, β in sylvis humidis opacis. 4 VII. VIII.

umbellatum. 8. **HIERACIUM** foliis linearibus subdenticulatis sparsis, floribus subumbellatis. *Sp. pl.* 804. *Fl. suec.* 704.

Hieracium fruticosum angustifolium majus. *Bauh. pin.* 129. *Park.* 801. *R. Syn.* 168.

Hieracium intybaceum. *Ger. Em.* 298.

β Pulmonaria angustifolia glabra. *Pet. h. br.* t. 13. f. 12.

γ Pulmonaria graminea. *Pet. h. br.* t. 13. f. 12.

Anglis, narrow-leaved bushy Hawkweed.

Habitat cum priore. 4. VII. VIII.

CREPIS. 819.

*Recept. nudum. Cal. calyculatus: squamis deciduis.
Pappus plumosus, stipitatus.*

1. CREPIS foliis lanceolatis sagittatis sessilibus lævibus, *testorum.*
inferioribus dentatis. *Sp. pl.* 807. *Fl. suec.* 705.

Hieracium luteum glabrum sive minus hirsutum. *Bauh. hist. II.* 1024. *R. Syn.* 165.

Cichoreum pratense luteum lævius. *Bauh. pin.* 126.
Park. 778.

Hieracium aphacoides. *Ger. Em.* 297.

β Hieracium aphacoides acutum. *Pet. h. br. t.* 12. *f.* 7.

γ Hieracium folio acuto minus. *Pet. h. br. t.* 12. *f.* 3.

δ Hieracium folio obtuso minus. *Pet. h. br. t.* 12. *f.* 4.

ε Hieracium Lactucæ folio. *Pet. h. br. t.* 12. *f.* 5.

Dens leonis ramosus glaber. *Pet. h. br. t.* 14. *f.* 12.

Anglis, smooth Succory Hawkweed.

Habitat in pratis, pascuis, muris, testis ubique. ☉. VI.

—IX.

2. CREPIS foliis pinnato hastatis scabris: basi superne biennis.
dentatis, calycibus muricatis. *Sp. pl.* 807. *Fl. suec.*
706.

Hieracium maximum chondrillæ folio, asperum. *Bauh. pin.* 127. *pr.* 64. *R. Syn.* 166.

Hieracium cichorei folio majus. *Pet. h. br. t.* 12.
f. 10.

Anglis, rough Succory Hawkweed.

Habitat in pratis pascuis et ad sepes. In Cantio frequens. *R. Syn.* 8. VII. VIII.

3. CREPIS foliis pinnatis angulatis, petiolis dentatis. *fætida.*
Roy. lugdb. 126. *Sp. pl.* 807.

Hieracium luteum cichorei sylvestris folio, amygdalas
amaras olens. *Hist. ox. III.* 63. *f.* 7. *t.* 4. *f.* 4.

Hieracium castorei odore. *Pet. h. br. t.* 12. *f.* 8.
monspeliensium. *R. Syn.* 165.

Senesio hirsutus. *Bauh. pin.* 131.

Anglis, stinking Hawkweed.

Habitat in pascuis circa Cantabrigiam et alibi sed rarius. *R. Syn.* On Bansted Downs, by the side of the road

road to Darking in Surry. In Northfleet chalk-pits, Kent. Blackf. spec. bot. ♂. VI. VII.

HYOSERIS. 821.

Recept. nudum. Cal. subæqualis. Pappus pilosus vel obsoletus.

minima. 1. HYOSERIS caule diviso nudo. *Sp. pl. 809. Fl. suec. 707.*

Hyoseris angustifolia. Tabern. ic. 180.

Hyoseris mascula. Ger. Em. 288.

Hieracium minimum Clusii, Hyoseris Tabernæmontani et Gerardi. Park. 791. R. Syn. 173.

Hieracium minus folio subrotundo. Baub. pin. 127.

Anglis, small Swine's Succory.

Habitat in pratis et arvis solo glareoso. Circa Hampton-court copiose observavit, D. Doody. R. Syn. in a sand-pit on the right-hand going to the duke of Argyll's house at Petersham, Surry, Dr. Wilmer. In the common corn-field at Walthamstow, Essex, Dr. Watson. In Tuddington field near Hampton-court. Blackf. spec. bot. ☉. V. VI.

HYPOCHÆRIS. 822.

Recept. paleaceum. Cal. subimbricatus. Pappus plumosus.

radicata. 1. HYPOCHÆRIS foliis dentato-finuatis scabris, caule ramoso nudo, pedunculis squamosis. *It. scan. 225. Sp. pl. 811. Fl. suec. 709.*

Hieracium, dentis leonis folio obtuso, majus. Baub. pin. 127.

Hieracium longius radiculatum. Lob. hist. 120. Ger. Em. 298. Park. 790. R. Syn. 165.

Anglis, Long-rooted Hawkweed.

Habitat in pratis et pascuis. 4. V.—VII.

macculata. 2. HYPOCHÆRIS caule subnudo, ramo subfolitario, foliis ovato-oblongis integris dentatis. *Sp. pl. 810. Fl. suec. 708.*

Hieracium I. latifolium. Clus. hist. II. 139. R. Syn. 167.

Hieracium

Hieracium alpinum latifolium hirsutie incanum, flore magno. Bauh. pin. 128.

Anglis, spotted Hawkweed.

Habitat in pratis montosis. Found on the banks of the devil's ditch near Reche, not far from Newmarket; also on Bernuk heath in Northamptonshire. R. Syn. 4. VII.

3. *HYPOCHÆERIS glabra: calycibus oblongis imbricatis, caule nudo foliis dentato-sinuatis. Sp. pl. 811.*

Hieracium minus dentis leonis folio oblongo glabro. Bauh. pin. 127.

Hieracium parvum in arenosis nascens, seminum papillis densius radiatis. R. Syn. 166.

Hieracium minimum. Col. ecphr. I. 27.

Anglis, smooth Hawkweed.

Habitat in pratis montosis. Found on the gravelly grounds near Middleton in Warwickshire. R. Syn. circa Bristolium frequens. 8. VII.

L A P S A N A. 823.

Recept. nudum. Cal. calyculatus, squamis singulis interioribus canaliculatis.

1. *LAPSANA calycibus fructus angulatis, pedunculis communis, tenuibus ramosissimis. Sp. pl. 811. Fl. suec. 710.*

Lampfana. Ger. Em. 255. R. Syn. 173. vulgaris. Park. 810.

Soncho affinis Lampfana domestica. Bauh. pin. 124.

Anglis, Nipplewort.

Habitat in sepibus et ruderatis. 6. VI. VII.

C I C H O R I U M. 825.

Receptaculum paleaceum. Cal. calyculatus. Pappus margo quinquedentatus.

1. *CICHORIUM caule simplici ramoso, floribus confertis; omnibus sessilibus. Syst. nat. 1197.*

Cichorium caule simplici, foliis dentato-sinuatis. Sp. pl. 813. Fl. suec. 711.

Cichoreum sylvestre. Ger. Em. 284. Park. 776. R. Syn. 172.

Cichoreum sylvestre s. officinarum. Bauh. pin. 125. Anglis,

Anglis, wild Succory.

Habitat *ad agrorum margines*. ♂. VII. VIII.

Pharmac. Cichorei *Herba*, *Radix*.

**** Capitali.**

A R C T I U M. 830.

Cal. globosus, squamis apice hamis inflexis.

- Lappa.* 1. ARCTIUM foliis cordatis inermibus petiolatis. *Sp. pl.* 816. *Fl. suec.* 712.
Lappa major seu Arctium dioscoridis. *Baub. pin.* 198. *R. Syn.* 197.
Bardana major. *Ger. Em.* 809.
Bardana vulgaris major. *Park.* 1222.
 β *Lappa* major capitulo glabro maximo. *R. Syn.* 196.
 γ *Lappa* major capitulis parvis glabris. *R. Syn.* 197.
 δ *Lappa* major montana, capitulis tomentosis s. Arctium Dioscoridis. *Baub. pin.* 198.
 ε *Lappa* major montana, capitulis minoribus, rotundioribus et magis tomentosis. *Tourn.* 450.
 ζ *Lappa* major ex omni parte minor, capitulis parvis eleganter reticulatis. *Pluk. alm.* 205.
Anglis, Burdock, or Clot-burr.
 Habitat *in vis, ruderalis, cultis*. ♂. VII. VIII.

S E R R A T U L A. 831.

Cal. subcylindricus, imbricatus, muticus.

- tinctoria.* 1. SERRATULA foliis pinnatifidis: lacinia terminali maxima. *Sp. pl.* 816. *Fl. suec.* 713.
Serratula. *Baub. pin.* 225. *hist III.* 23. *R. Syn.* 196.
Serratula purpurea. *Ger. Em.* 713.
Serratula vulgaris flore purpureo. *Park.* 474.
 β *Serratula* flore candido. *Boerb. ind. alt.* 144.
Anglis, Saw-wort.
 Habitat *in sylvis et pratis*. 4. VII.
- alpina.* 2. SERRATULA calycibus hirsutis ovatis. *Sp. pl.* 816. *Fl. suec.* 714.
Serratula foliis petiolatis: radicalibus ovato-oblongis dentatis, caulinis integerrimis. *Fl. suec. ed. I.* 661.
 Cirsium

Cirsium foliis triangularibus lunate dentatis, subtus tomentosis, inermibus, in spica congestis. *Hall. helv.* 683. t. 22.

Cirsium humile montanum cynoglossi folio, polyanthemum. *R. Syn.* 193.

Carduo-cirsium minus britannicum, floribus plurimis summo caule congestis. *Pluk. ph* t. 154. f. 3.

β *Cirsium* montanum polyanthemum falicis folio angusto denticulato. *R. Syn.* 193.

Anglis, Mountain Saw-wort.

Habitat in summis rupibus Arvonix, v. g. Clogwyn y Carnedh yscolion duon, &c. On the highest rock of Snowdon, and near Brearcliff, near Brunly, Lancashire. *R. Syn.* 4. VII.

2. SERRATULA foliis dentatis spinosis. *Sp. pl.* 820. *arvensis*.

Fl. suec. 715.

Carduus radice repente, foliis lanceolatis dentatis margine aculeato. *Fl. lapp.* 293.

Carduus vulgarissimus viarum. *Ger. Em.* 1173. *R. Syn.* 194.

Carduus ceanothos five viarum et vinearum repens. *Park.* 959.

Carduus vinearum repens folio Sonchi. *Bauh. pin.* 377.

Anglis, Corn Saw-wort, or Way-thistle.

Habitat in arvis et ad vias. 4. VII.

C A R D U U S. 832.

Cal. ovatus, imbricatus squamis spinosis. *Recept.* pilosum.

* *Foliis decurrentibus.*

1. CARDUUS foliis decurrentibus hastato-pinnatis his- lanceolatis : laciniis divaricatis, calycibus spinosis villosis, tus. caule piloso. *Sp. pl.* 821. *Fl. suec.* 716.

Carduus lanceatus. *Ger. Em.* 1174. *R. Syn.* 195.

Carduus lanceatus latifolius. *Bauh. pin.* 385. f. major. *Park.* 982.

β *Carduus* lanceatus major. *R. Syn.* 195.

γ *Carduus* lanceatus flore et capite minoribus. *Merr. pin.* *R. Syn.* 195.

X

Anglis,

Anglis, Spear-thistle.

Habitat in viarum marginibus ubique. ♂. VII.

- nutans*. 2. **CARDUUS** foliis spinosis semidecurrentibus, floribus cernuis: squamis calycinis superne patentibus. *Sp. pl.* 821. *Fl. suec.* 717.
Cirsium majus singulari capitulo magno. *Bauh. pin.* 377.
Carduus nutans. *Bauh. hist.* III. 56. *R. Syn.* 193.
Carduus moschatus. *Pet. h. br. t.* 21. f. 1.
Anglis, Musk-thistle.
 Habitat in pratis et pascuis siccissimis in pratis inter ericetum nigrum et Charleton in Cantio. ♂. VII.
- acanthoides*. 3. **CARDUUS** foliis sinuatis decurrentibus: margine spinosis, calycibus pedunculatis solitariis erectis villosis. *Sp. pl.* 821. *Fl. suec.* 718.
Carduus acanthoides. *Bauh. hist.* III. 59.
Carduus spinosissimus capitulis minoribus. *P. B. R. Syn.* 194.
 β *Carduus palustris* mitior, bardanæ capitulo in summo caule singulari. *Pluk. alm.* 82.
Anglis, welshed Thistle.
 Habitat ad vias et in aggeribus fossarum. ☉. VI. VII.
- crispus*. 4. **CARDUUS** foliis sinuatis decurrentibus: margine spinosis, floribus aggregatis terminalibus muticis. *Sp. pl.* 821. *Fl. suec.* 719.
Carduus caule crispo. *Bauh. hist.* III. 59. *R. Syn.* 194.
Carduus spinosissimus angustifolius vulgaris. *Bauh. pin.* 385. *Læf. pruss.* 43. t. 5.
Carduus polyanthos. *Park.* 982. primus. *Ger. Em.* 1173.
Anglis, Thistle upon Thistle.
 Habitat ad vias et in aggeribus fossarum. ☉. VI.
- palustris*. 5. **CARDUUS** foliis linearibus dentatis decurrentibus: margine spinosis, floribus racemosis erectis, pedunculis inermibus. *Sp. pl.* 822. *Fl. suec.* 720.

Carduus palustris. *Baub. pin.* 377. *pr.* 156. *Park.* 983. *Pet. h. br. t.* 21. *R. Syn.* 194.
Anglis, Marsh-thistle.
Habitat in pratis palludosis et umbrosis. 4. VII.

6. *CARDUUS* foliis lanceolatis decurrentibus denticulis *dissectus*,
 inermibus, calycibus spinosis. *Sp. pl.* 822.

Cirsium anglicum. *Lob. ic.* 583. *Ger. Em.* 1183.
R. Syn. 193. primum. *Park.* 961.

Cirsium majus singulari capitulo magno, f. incanum
 varie dissectum. *Baub. pin.* 377.

Anglis, English soft or gentle Thistle.

Habitat in palustribus passim. 4. VI. VII.

** *Foliis amplexicaulibus.*

7. *CARDUUS* foliis lanceolatis dentatis amplexicauli- *helenioides*
 bus : spinulis inæqualibus ciliatis, caule inermi.
Sp. pl. 825.

Cirsium foliis longe lanceolatis ferratis : subtus to-
 mentosis. *Hall. helv.* 683. *t.* 21.

Cirsium britannicum Clusii repens. *Baub. hist.* III.
 46. *R. Syn.* 193.

Cirsium aliud Anglicum. *Park.* 961.

Cirsium singulari capitulo squamato, vel incanum al-
 terum. *Baub. pin.* 377.

Anglis, melancholy Thistle.

Habitat in montosis agri Eboracensis, Westmorlandici,
Cumberlandici et Cambrobritannici frequens. On
Hounslow heath. *R. Syn.* 4. VI. VII.

8. *CARDUUS* foliis amplexicaulibus hastato pinnatifi- *marianus*;
 dis spinosis calycibus aphyllis : spinis canaliculatis
 duplicato spinosis. *Sp. pl.* 823.

Carduus squamis calycinis margine apiceque spinosis.
Hort. cliff. 393.

Carduus Mariæ. *Ger. Em.* 989. *R. Syn.* 195. vul-
 garis. *Park.* 979.

Carduus albis macculis notatus vulgaris. *Baub. pin.*
 281.

β *Carduus Mariæ* hirsutus non macculatus. *R. Syn.*
 194.

SYNGENESIA : POLYGAMIA ÆQUALIS.

Anglis, Milk-thistle.

Habitat *ad agrorum margines et in aggeribus fossarum.*

⊙. VII.

Pharmac. Cardui mariæ *Herba, semina.*

eriphorus 9. CARDUUS foliis sessilibus bifariam pinnatifidis : lacinii alternis erectis, calycibus globosis erectis. *Sp. pl.* 823.

Carduus foliis finuatis decurrentibus : denticulis superficieque spinosis, calycibus lanigeris. *Roy. lugdb.* 133.

Carduus eriocephalus. *Ger. Em.* 1152.

Carduus capite rotundo tomentoso. *Bauh. pin.* 382.

Carduus tomentosus corona fratrum dictus. *Park.* 978.

Anglis, Woolly-headed Thistle.

Habitat *in pratis montosis. Circa Bristol copiose.* ♂. VII.

acaulos. 10. CARDUUS acaulis, calyce glabro. *Sp. pl.* 1199. *Fl. suec.* 722.

Carduus acaulos, minore purpureo flore. *Bauh. pin.* 380. *Ger. Em.* 1158. *R. Syn.* 195.

Carduus acaulis septentrionalium. *Park.* 969.

Anglis, dwarf Carline-thistle.

Habitat *in pascuis montosis siccioribus. In ericeto nigro prope Londinum.* ♀. VII.

O N O P O R D U M. 834.

Recept. nudum. *Cal.* squamæ mucronatæ.

Acanthi- 1. ONOPORDUM calycibus squarrosis : squamis subulatis, foliis ovato-oblongis sinuatis. *Sp. pl.* 827. *Fl. suec.* 724.

Onopordum foliis decurrentibus margine spinosis. *Hort. cliff.* 393.

Carduus tomentosus, Acanthium dictus, vulgaris. *R. Syn.* 196.

Acanthium album. *Ger. Em.* 1149.

Acanthium vulgare. *Park.* 979.

Spina alba tomentosa latifolia sylvestris. *Bauh. pin.* 382.

Anglis, Cotton-thistle.

Habitat *in ruderalis.* ♂. VII.

CAR-

CARLINA. 836.

Cal. radiatus squamis marginalibus longis, coloratis.

1. CARLINA caule multifloro corymbofo, floribus ter-*vulgaris.*
minalibus. *Hort. cliff.* 395. *Sp. pl.* 829. *Fl. suec.*
725.

Carlina sylvestris major. *Ger. Em.* 1159.

Carlina sylvestris quibusdam aliis atractylis. *Bauh. hist.*
III. 81. *R. Syn.* 175.

Cnicus sylvestris spinosior. *Bauh. pin.* 378.

Anglis, wild Carline Thistle.

Habitat in pratis et pascuis siccioribus. ♂. VI.

*** *Discoidei.*

BIDENS. 840.

Recept. paleaceum. *Pappus* aristatis erectis, scabris.

Cal. imbricatus. *Cor.* rarius flosculo uno alterne
radiante instruitur.

1. BIDENS foliis trifidis corollis flosculis, calycibus sub-*tripartita.*
foliolis, feminibus erectis. *Sp. pl.* 831. *Fl. suec.* 726.
Verbesina f. Cannabina aquatica flore minus pulchro,
elator et magis frequens. *Bauh. hist. III.* 1073.
R. Syn. 187.

Cannabina aquatica folio tripartito diviso. *Bauh. pin.*
321.

Eupatorium cannabinum fæmineum. *Ger. Em.* 711.

Eupatorium aquaticum duorum generum. *Park.* 595.

Anglis, trifid Water-hemp-agrimony.

Habitat in palustribus et aquosis, ubi aquæ stagnant. ☉:
VII. VIII.

2. BIDENS foliis lanceolatis amplexicaulibus, floribus *cernua.*
cernuis, feminibus erectis. *Sp. pl.* 832. *Fl. suec.*
727.

Verbesina pulchriore flore luteo. *Bauh. hist. II.* 1074.
R. Syn. 187.

Cannabina aquatica folio non diviso. *Bauh. pin.* 321.

Eupatorium cannabinum fæminæ varietas altera. *Ger.*
Em. 711.

Eupatorium aquaticum folio integro. *Park.* 596.

X 3

Anglis,

SYNGENESIA : POLYGAMIA ÆQUALIS.

Anglis, whole-leaved Water-hemp-agrimony.
Habitat in fossis et palustribus frequens. ☉. VII. VIII.

- minima.* 3. *BIDENS* foliis lanceolatis sessilibus, floribus seminibusque erectis.
Verbeina minima. *Cat. giff.* 167. *app.* 66. *R. Syn.* 188. *t.* 7. *f.* 2.
Eupatorium cannabinum palmare et angustifolium.
Nierr. pin.
Anglis, the least Water-hemp-agrimony.
Habitat in palustribus. *In Surrey near to Somerset-bridge, in the fish-pond upon the moor.* *R. Syn.* ☉. VII. VIII.

EUPATORIUM. 842.

Recept. nudum. *Pappus plumosus.* *Cal. imbricatus, oblongus.* *Stylus semibifidus, longus.*

- Cannabi-num.* 1. *EUPATORIUM* foliis digitatis. *Sp. pl.* 838. *Fl. suec.* 728.
Eupatorium cannabinum. *Bauh. pin.* 320. *Park.* 595. *R. Syn.* 179.
Eupatorium cannabinum mas. *Ger. Em.* 711.
Eupatorium cannabinum folio integro, seu non digitato. *R. Syn.* 180.
Anglis, Hemp-agrimony, or Dutch-agrimony.
Habitat ad ripas fluviorum et rivulorum. ☿. VII. VIII.

POLYGAMIA SUPERFLUA.

* *Discoidei.*

TANACETUM. 848.

Recept. nudum. *Pappus submarginatus.* *Cal. imbricatus, hemisphericus.* *Cor. radii obsoleti, trifidi.*

- vulgare.* 1. *TANACETUM* foliis bipinnatis serratis. *Sp. pl.* 848. *Fl. suec.* 730.
Tanacetum. *Ger. Em.* 650. *R. Syn.* 188. *vulgare.* *Park.* 81. *luteum.* *Bauh. pin.* 132.
 β *Tanacetum foliis crispis.* *Bauh. pin.* 132. *Anglis,*

Anglis, common Tanfy.

Habitat in pratis montosis. Circa Kendal in Westmorlandia frequens. 2. VIII.

Pharmac. Tanaceti Herba, Flores, Semina.

ARTEMISIA. 849.

Recept. subvillosum vel nudiusculum. Pappus nullus.

Cal. imbricatus: squamis rotundatis, conniventibus. Cor. radii nulli.

* *Procumbentes ante florescentiam.*

1. ARTEMISIA foliis multifidis linearibus, caulibus campestris procumbentibus virgatis. *Sp. pl.* 846. *Fl. suec.* 732.

Artemisia tenuifolia. *Dod. pempt.* 33.

Abrotanum campestre. *Bauh. pin.* 136. *Ger. Em.* 1106. *Park.* 94. *R. Syn.* 191.

Anglis, Field Southernwood.

Habitat in campis montosis apricis aridis. Near Elden in Suffolk. *R. Syn.* 2. VIII.

2. ARTEMISIA foliis multipartitis tomentosis, racemis maritima. cernuis flosculis fæmineis ternis. *Sp. pl.* 846. *Fl. suec.* 734.

Absinthium marinum album. *Ger. Em.* 1099. *R. Syn.* 188.

Absinthium seriphium belgicum. *Bauh. pin.* 139. *hist. III.* 178.

- β *Absinthium maritimum seriphio belgico simile, latiore folio, odoris grati.* *Pluk. R. Syn.* 188.

- γ *Absinthium maritimum Gallicum.* *Bauh. pin.* 139.

- δ *Absinthium seriphium tenuifolium marinum Narbonense.* *Bauh. hist. III.* 177.

- ε *Absinthium marinum nostras.* *R. Syn.* 189.

Anglis, Sea Wormwood.

Habitat in littoribus maritimis. 2. VIII.

Pharmac. Absinthii marini Herba.

** *Erectæ herbaceæ, foliis compositis.*

3. ARTEMISIA foliis compositis multifidis, floribus Absinthii subglobosis pendulis: receptaculo villosa. *Sp. pl. um.* 848. *Fl. suec.* 735.

Abinthium vulgare. *Park.* 98. *R. Syn.* 188. *majus.*
Baub. hist. III. 168.

Abinthium latifolium seu *Ponticum.* *Ger. Em.* 1096.

Abinthium ponicum f. *romanum officinarum* f. *disco-
ridis.* *Baub. pin.* 138. *Tourn.* 457.

Anglis, common Wormwood.

Habitat in ruderatis et ad vias. 4. VIII.

Pharmac. *Abinthii vulgaris Herba.*

vulgaris. 4. *ARTEMISIA* foliis pinnatifidis planis incis: sub-
tus tomentosis. racemis simplicibus, floribus ova-
tis, radio quinquefloro. *Sp. pl.* 848.

Artemisia vulgaris. *Park.* 90. *R. Syn.* 190. *Ger. Em.*

1103. *Baub. hist.* III. 184. *major.* *Baub. pin.* 137.

Anglis, Mugwort.

Habitat ad sepes et margines agrorum. 4. VIII.

Pharmac. *Artemisæ Herba.*

GNAPHALIUM. 850.

Recept. nudum. *Pappus plumosus.* *Cal.* imbricatus:
squamis marginibus rotundatis, membranaceis, co-
loratis.

* *Argyrocomæ calycibus albis.*

dioicum. 1. *GNAPHALIUM* caule simplicissimo, corymbo sim-
plici terminali, sarmentis procumbentibus. *Sp. pl.*
850. *Fl. suec.* 736.

Gnaphalium montanum album. *Ger. Em.* 640. *R.*
Syn. 181.

Gnaphalium montanum sive *Pes cati.* *Park.* 690.

Gnaphalium montanum flore rotundiore. *Baub. pin.*
263.

Anglis, Mountain Cudweed, or Cat's-foot.

Habitat in pratis montosis aridis. On Newmarket
heath, not far from Bottesham beacon, plentifully;
also on Bernak heath in Lincolnshire; in comitati-
bus Eboracensi, Lancastriensi, Westmorlandico,
frequens. 4. V.

*margari-
taceum.* 2. *GNAPHALIUM* foliis lineari-lanceolatis acumina-
tis alternis, caule superne ramoso, corymbis fasti-
giatis.

giatis. *Hort. cliff.* 401. *Gmel. sib.* II. 107. *Sp. pl.* 850.

Gnaphalium americanum. *Bauh. hist.* III. 162. *Ger.*

Em. 641. *R. Syn.* 182. *latifolium.* *Bauh. pin.* 263.

Argyrocome f. *Gnaphalium americanum.* *Park.* 690.

Anglis, American Cudweed.

Habitat in pratis et pascuis. In prato prope Bocking, vicum agri Efflexiensis, D. Dale; on the banks of Rymny river, for the space of at least twelve miles, Mr. Lhwyd. R. Syn. 4. VIII.

3. *GNAPHALIUM* foliis femiamplexicaulibus ensifor-
mibus repandis obtusis utrinque pubescentibus, flo-
ribus conglomeratis. *Ray. lugdb.* 149. *Sp. pl.* 851.

Gnaphalium majus lato oblongo folio. *Bauh. pin.* 263.

Pluk. ph. t. 31. *f.* 6.

Elichrysum sylvestre latifolium capitulis conglobatis.

Bauh. pin. 264. *R. Syn.* 182.

Anglis, Jersey Cudweed.

Habitat in the isle of Jersey, on dry banks and walls, very common. R. Syn. 4. VII. VIII.

** *Filagines calycibus cinereis.*

4. *GNAPHALIUM* caule simplicissimo, floribus spar-
sifloris. *Fl. lapp.* 298. *suec.* 739. *Sp. pl.* 856.

Gnaphalium anglicum. *Ger. Em.* 639. *R. Syn.* 180.

vulgare majus. *Park.* 685.

Gnaphalium majus, angusto oblongo folio, alterum.

Bauh. pin. 263.

Gnaphalium rectum. *Bauh. hist.* III. 160.

Anglis, upright Cudweed.

Habitat in pascuis et sylvis arenosis. In ericeto Hampsteadiano, et prope Charleton in Cantio. 3. VIII.

5. *GNAPHALIUM* caule ramoso diffuso, floribus con-
fertis terminalibus. *Fl. lapp.* 300. *suec.* 740. *Sp. pl. sum.* 856.

Gnaphalium longifolium humile ramosum capitulis nigris. *R. Syn.* 181.

Gnaphalium medium. *Bauh. pin.* 263.

Filago minor. *Park.* 686.

Anglis,

Anglis, black-headed Cudweed.

Habitat in aquosis præcipue ubi aquæ hyemali tempore stagnarunt. ☉. VIII.

C O N Y Z A. 854.

Recept. nudum. *Pappus* simplex. *Cal.* imbricatus, subrotundus. *Cor.* radii trifidi.

squarrosa. 1. CONYZA foliis lanceolatis acutis, caule annuo corymbofo. *Hort. cliff.* 405. *Roy. lugdb.* 157. *Sp. pl.* 861.

Conyza major vulgaris. *Bauh. pin.* 265.

Baccharis monspeliensium. *Ger. Em.* 792. *Park.* 114.

R. Syn. 179.

Anglis, Plowman's Spikenard.

Habitat in pratis et pascuis montosis et ad vias. ♂. VII. VIII.

** *Radiati.*

E R I G E R O N. 855.

Recept. nudum. *Pappus* pilosus. *Cor.* radii lineares, angustissimi.

acre. 1. ERIGERON pedunculis alternis unifloris. *Sp. pl.* 863. *Fl. suec.* 741.

Conyza cærulea acris. *Ger. Em.* 484. *Bauh. pin.* 265.

Conyza odorata cærulea. *Park.* 126.

After arvensis cæruleus acris. *Tourn.* 481. *R. Syn.* 175.

Anglis, blew-flowered Fleabane.

Habitat in pratis et pascuis siccissimis. 4. VII. VIII.

canadense. *2. ERIGERON caule floribusque paniculatis. *Sp. pl.* 863.

Conyza canadensis annua, acris, alba, linariæ foliis. *Boc. pl. rar. t.* 46. *R. Syn.* 175.

Conyza annua acris alba, elatior, linariæ foliis. *Hist. ox. III.* 115. *f. II. t.* 20. *f.* 29.

Anglis, Canada Fleabane.

Habitat in ruderalis circa Londinum frequens. ☉. VIII.

TUS-

TUSSILAGO. 856.

Recept. nudum. *Pappus simplex.* *Cal.* squamæ æquales, discum æquantes, submembranaceæ.

1. TUSSILAGO scapo imbricato unifloro, foliis subcordatis angulatis denticulatis. *Mat. med.* 391. *Sp. pl.* 865. *Fl. suec.* 743. *Farfara.*

Tussilago. *Ger. Em.* 811. *Park.* 1220. *R. Syn.* 173. vulgaris. *Bauh. pin.* 197.

Anglis, common Coltsfoot.

Habitat in locis humidis. 4. III.

Pharmac. *Tussilaginis sive Farfaræ Herba, Flores.*

2. TUSSILAGO thyrso oblongo, flosculis femineis nudis plurimis. *Sp. pl.* 866. *Roy. lugdb.* 159. *hybrida.*

Petasites major, floribus pediculis longis infidentibus.

R. Syn. 179. *Dill. elth.* 309. t. 230. f. 297.

Anglis, long-stalked Butter-bur.

Habitat in pratis humidis. Near Loughborough in Leicestershire, Mr. Hill. *Blackf. spec. bot.* 4. III.

3. TUSSILAGO thyrso ovato, flosculis omnibus hermaphroditis. *Sp. pl.* 866. *Fl. suec.* 746. *Petasites.*

Petasites. *Ger. Em.* 814. *R. Syn.* 179. vulgaris. *Park.* 419.

Petasites major et vulgaris. *Bauh. pin.* 197.

Anglis, common Butter-bur.

Habitat in pratis et pascuis humidis. 4. III. IV.

Pharmac. *Petasitidis Radix.*

SENECIO. 857.

Recept. nudum. *Pappus simplex.* *Cal.* cylindricus, calyculatus : squamis apice sphacelatis.

* *Floribus flosculosis.*

1. SENECIO corollis nudis, foliis pinnato-sinuatis amplexicaulibus, floribus sparsis. *Roy. lugdb.* 165. *vulgaris.*
Sp. pl. 867. *Fl. suec.* 747.

Senecio foliis pinnatifidis denticulatis : laciniis æqualibus patentissimis, rachi lineari. *Hort. cliff.* 406.

Senecio vulgaris. *Park.* 671. *R. Syn.* 178.

Senecio

Senecio minor vulgaris. Bauh. pin. 131.

Erigeron. Ger. Em. 278.

Anglis, common Groundfel, or Simpson.

Habitat in cultis, ruderatis ubique. ☉. V.

Pharmac. fenecionis Herba.

*** Floribus radiatis: radio revolutis.*

sylvaticus. 2. SENECIO corollis revolutis, foliis pinnatifidis denticulatis, caule corymbofo erecto. Sp. pl. 868. Fl. suec. 748.

Senecio minor, latiore folio five montanus. Bauh. pin. 131. R. Syn. 178. Dill. elth. 258. t. 258. f. 337.

Senecium montanum. Tabern. ic. 169.

Erigerum. Ger. Em. 279.

Jacobæa annua fenecionis folio, parvo flore. Aët. gall. 1720. p. 298. n. 19.

Anglis, Mountain Groundfel.

Habitat in sylvis, sepibus et cricetis frequens. ☉. VII.

viscosus. 3. SENECIO corollis revolutis, foliis pinnatifidis viscidis, squamis calycinis laxis longitudine perianthii. Sp. pl. 868. Fl. suec. 749.

Senecio foliis pinnatifidis sessilibus: laciniis angulosis: superioribus latoribus, florum radiis disco brevioribus. Roy. lugdb. 141.

Senecio incanus pinguis. Bauh. pin. 131.

Senecio hirsutus viscidus major odoratus. Bauh. hist. II. 1092. R. Syn. 178.

Senecio foetidus. Park. 671.

Senecio hirsutus viscidus graveolens. Dill. elth. 347. t. 258. f. 336.

Anglis, Cotton Groundfel, or stinking Groundfel.

Habitat in arenosis passim. ☉. VI.

**** Floribus radiatis: radio patente, fol. pinnatifidis.*

Jacobæa. 4. SENECIO corollis radiantibus, foliis pinnato-lyratis: laciniis lacinulatis, caule erecto. Sp. pl. 870. Fl. suec. 751.

Jacobæa vulgaris. Bauh. hist. II. 1057. R. Syn. 177: major. Park. 668.

Jacobæa.

Jacobæa. *Ger. Em.* 280. vulgaris laciniata. *Bauh. pin.* 131.

Anglis, common Ragwort.

Habitat in pratis, pascuis, montosis, et ad vias. 2.
VII.

- 5 SENECIO corollis radiantibus, foliis dentatis, radi- *aquaticus*.
calibus ovatis, caulinis pinnatifidis; lacinia extima
majore.

Jacobæa latifolia palustris. *R. Syn.* 178.

Jacobæa barbæræ instar laciniata. *Fl. prus.* 129.
t. 35.

Jacobæa latifolia. *Bauh. hist. II.* 1056. *Ger. Em.*
280.

Jacobæa latifolia pannonica prima. *Park.* 668.

Jacobæa alpina laciniata flore Buphthalmi. *Bauh. pin.*
131.

Anglis, Water Ragwort.

Habitat in aquosis. 2. VII. VIII.

6. SENECIO corollis radiantibus, foliis pinnatifidis *erucifoli-*
dentatis incanis, caule erecto. *Sp. pl.* 869. *Fl. us.*
suec. 750.

Jacobæa fenecionis folio incano perennis. *R. Syn.*
177.

Jacobæa altissima, foliis erucæ artemisiæve similibus
et æmulis. *Rupp. jen.* 164.

Anglis, hoary perennial Ragwort.

Habitat in sylvis et ad sepes. 2. VII.

**** *Floribus radiatis: radio patente, fol. indivisis.*

7. SENECIO corollis radiantibus, foliis ensiformibus *paludosus*.
acute ferratis subtus subvillosis, caule stricto. *Sp. pl.*
870. *Fl. suec.* 752.

Solidago caule simplici, corymbo terminali, foliis sub-
tus tomentosis, flosculis fæmineis sedecim. *Lech.*
scan. 68.

Jacobæa foliis longis integris et mucronatis. *Hist. ox.*
III. 110. *f.* 7. *t.* 7. *f.* 22.

Virgæ aureæ five Solidagini angustifoliæ affinis, Lin-
gua avis Dalechampii. *Bauh. hist. II.* 1063. *R. Syn.*
176.

Conyza

Conyza palustris. *Park.* 1231. *ferratifolia*. *Bauh. pin.* 266.

Anglis, Marsh Ragwort, or Bird's-tongue.

Habitat in fossis palustribus earumque aggeribus in insula Eliensi, near Stretham ferry. *R. Syn.* 4. VIII.

sarraceni- 8. **SENECIO** corollis radiantibus, floribus corymbosis, *cus.* foliis lanceolatis ferratis glabriusculis. *Hort. ups.* 266. *Sp. pl.* 871.

Solidago caule simplici, corymbo terminali; pedunculis partialibus alternis nudis longitudine folio. *Hort. cliff.* 410.

Solidago sarraceni. *Fuchs. hist.* 728. *Dod. pempt.* 141.

Virga aurea angustifolia ferrata. *Bauh. pin.* 268.

Virga aurea maxima radice repente. *Bobart. R. Syn.* 177.

Anglis, broad-leaved Ragwort.

Habitat in pratis, pascuis, nemorosis subhumidis. Found plentifully by the side of a small river between Wells and Glastenbury, by Mr. Bobart. It grows as plentifully in the fields at Salkeld as the vulgaris, Mr. Nicholson. *R. Syn. circa* Clapham in comitatu Eboracensi. 4. VII. VIII.

S O L I D A G O. 856.

Recept. nudum. *Pappus simplex*. *Cor.* radii circiter quinque. *Cal.* squamæ imbricatæ, clausæ.

Virgaurea 1. **SOLIDAGO** caule subflexuoso angulato racemis subpaniculatis erectis confertis. *Sp. pl.* 880. *Fl. suec.* 754.

Solidago floribus per caulem simplicem undique sparsis. *Fl. lapp.* 306.

Virga aurea. *Ger. Em.* 429. *R. Syn.* 176. vulgaris. *Park.* 542.

Virga aurea latifolia ferrata. *Bauh. pin.* 268.

Virga aurea vulgaris latifolia. *Bauh. hist. II.* 1062.

β *Virga aurea* IV. s. *angustifolia minus ferrata*. *Bauh. pin.* 268.

γ *Virga aurea vulgari humilior*. *R. Syn.* 176.

Anglis,

Anglis, common Golden Rod.

Habitat in *sylvis*, *sepibus*, *ericetis frequens*. γ in *erico* Hampstediano. α . VIII.

Pharmac. Virgæ aureæ *Herba*.

2. SOLIDAGO foliis lineari-lanceolatis subserratis in-*cambrica*.
canis panicula corymbosa terminali.

Virga aurea montana, folio angusto subincano, flos-
culis conglobatis. *R. Syn.* 177.

Virga aurea Cambriæ. *Pet. h. br. t.* 16. *f.* 11.

Anglis, Welch Golden Rod.

Habitat in *pratis montosis* in *Wallia*. α . VI.

A S T E R. 858.

Recept. nudum. *Pappus simplex*. *Cor.* radii plures
decem. *Cal.* imbricati, squamæ inferiores patulæ.

1. ASTER foliis lanceolatis integerrimis carnosis gla-*Tripolium*
bris, ramis inæqualibus, floribus corymbosis. *Roy.*
lugdb. 168. *Sp. pl.* 872. *Fl. suec.* 753.

Aster maritimus cæruleus *Tripolium dictus*. *R. Syn.*
175.

Tripolium vulgare majus et minus. *Ger. Em.* 413.

Tripolium majus cæruleum et minus. *Bauh. pin.* 267.

Tripolium majus f. vulgare et minus. *Park.* 673.

Anglis, Sea Starwort.

Habitat in *littoribus maritimis*. α . VIII.

I N U L A. 860.

Recept. nudum. *Pappus simplex*. *Antheræ*, basi in
fetas duas desinentes.

1. INULA foliis ovatis rugosis : subtus tomentosis, ca-*Helenium*
lycum squamis ovatis. *Amæn. acad. I.* 410. *Sp.*
pl. 881. *Fl. suec.* 755.

Helenium. *Ger. Em.* 793. *R. Syn.* 176. vulgare.
Bauh. pin. 276.

Helenium f. Enula campana. *Park.* 674. *Bauh. hist.*
III. 108.

Anglis, Elecampane.

Habitat in *pratis et pascuis humidis*. In *Essexia fre-*
quens. α . VII. VIII.

Pharmac. Enulæ campanæ *Radix*.

2. INULA

dysenterica. 2. *INULA* foliis oblongis, caule hirsuto paniculato, squamis calycinis fetaceis. *Sp. pl.* 882. *Fl. suec.*

757.

Conyza media. *Ger. Em.* 482. *R. Syn.* 174.

Conyza media afteris flore luteo, vel tertia Dioscoridis. *Bauh. pin.* 265.

Herba dysenterica. *Hoffm. altd.*

Anglis, middle Fleabane.

Habitat in aquosis et ad ripas fluviorum. 4. VIII.

Pulicaria. 3. *INULA* foliis undulatis amplexicaulibus caule prostrato. *Sp. pl.* 882. *Fl. suec.* 758.

Conyza minor. *Matth. P. II.* 223. *R. Syn.* 174.

Conyza minima. *Ger. Em.* 482.

Conyza minor flore globofo. *Bauh. pin.* 266.

Anglis, small Fleabane.

Habitat in locis ubi per hyemem aquæ stagnarant. 6. VIII. IX.

crithmoides. 4. *INULA* foliis linearibus carnosis tricuspidatis. *Sp. pl.* 883.

After flore terminali foliis linearibus tricuspidatis. *Hort. cliff.* 409. *Roy. lugdb.* 168. *Guett. stamp. II.* 464.

After maritimus flavus, *Crithmum chrysanthemum* dictus. *R. Syn.* 174.

Crithmum chrysanthemum. *Ger. Em.* 533. *Park.* 1287.

Crithmum maritimum flore afteris attici. *Bauh. pin.* 288.

Anglis, Golden Sampire.

Habitat in pratis maritimis passim. 4. VIII.

BELLIS. 864.

Recept. nudum, conicum. *Pappus* nullus. *Cal.* hemisphæricus : squamis æqualibus. *Sem.* obovata.

perennis. 1. *BELLIS* scapo nudo. *Sp. pl.* 886. *Fl. suec.* 761. *Bellis sylvestris minor.* *Bauh. pin.* 261. *R. Syn.* 184. *Bellis minor sylvestris.* *Ger. Em.* 635. *simplex.* *Park.* 530.

Anglis,

Anglis, common Daïsie.

Habitat in pratis et pascuis. 4. III.—IX.

Pharmac. Bellidis minoris, *Herba, Flores, Radix.*

CHRYSA N T H E M U M. 866.

Recept. nudum. *Pappus* nullus. *Cal.* hemisphæricus imbricatus: squamis marginalibus membranaceis.

1. CHRYSA N T H E M U M foliis amplexicaulibus: superne laciniatis; inferne dentato-ferratis. *Sp. pl.* 889. *Fl. suec.* 762. *segetum.*

Chrysanthemum segetum. *Ger. Em.* 743. *R. Syn.* 182. nostras *Park.* 1370.

Bellis lutea, foliis profunde incisfis major. *Baub. pin.* 262.

- β *Chrysanthemum segetum nostras*, folio glauco multiscisso majus, flore minore. *R. Syn.* 183.

- γ *Chrysanthemum segetum nostras*, calendulæ folio glauco, neque secto, neque ferrato. *Pluk. alm.* 102. *Anglis*, Corn Marigold.

Habitat in arvis. ☉. VI. VII.

2. CHRYSA N T H E M U M foliis amplexicaulibus oblongis: superne ferratis: inferne dentatis. *Sp. pl. themum.* 188 *Fl. suec.* 763. *Leucan-*

Leucanthemum vulgare. *Tourn.* 492. *R. Syn.* 184.

Bellis major. *Ger. Em.* 634. *Baub. hist. III.* 114.

Bellis major vulgaris f. sylvestris. *Park.* 528.

Bellis sylvestris caule folioso major. *Baub. pin.* 261.

Anglis, The greater Daïsie, or Ox-eye.

Habitat in pratis et pascuis siccis. 4. V.

Pharmac. Bellidis majoris *Herba, Flores.*

M A T R I C A R I A. 867.

Recept. nudum. *Pappus* nullus. *Cal.* hemisphæricus, imbricatus; marginibus solidis acutiusculis.

1. M A T R I C A R I A foliis compositis planis; foliolis ovatis incisfis, pedunculis ramosis. *Sp. pl.* 890. *Mat. um.* med. 400. *Partheni-*

Matricaria. Ger. Em. 652. R. Syn. 187. vulgaris. Park. 83.

Matricaria vulgaris f. fativa. Bauh. pin. 133. Anglis, Feverfew.

Habitat in ruderatis et ad sepes. ♂. f. 4. VI. Pharmac. Matricariæ Herba.

Chamomilla.

2. **MATRICARIA** receptaculis conicis, radiis patentibus, feminibus nudis, squamis calycinis margine æqualibus. *Fl. suec. 764.*

Chamæmelum. Ger. Em. 754. R. Syn. 184. vulgare. Dod. pempt. 257. Park. 85.

Chamæmelum vulgare, Leucanthemum Dioscoridis. Bauh. pin. 135.

Anglis, Corn Feverfew.

Habitat in arvis. ☉. VI.

Pharmac. Chamomeli vulgaris Herba, Flores.

- suaveolens* 3. **MATRICARIA** receptaculis conicis, radiis deflexis, feminibus nudis, squamis calycinis margine æqualibus. *Fl. suec. 766.*

Matricaria leucanthemos annua suaveolens, chamæmeli folio, ovariis albicantibus. Vaill. art. 556.

Chamæmelum tenue incisum, foliis floris deorsum protensis, umbone extuberante, suavius olens. Cup. cath. alt. 17.

Anglis, sweet-scented Feverfew.

Habitat in arenosis frequens. ☉. V.

- inodora.* 4. **MATRICARIA** receptaculis hemisphæricis, radiis patentibus, feminibus coronato-marginatis, squamis calycinis margine exfoletis. *Fl. suec. 765.*

Chamæmelum inodorum annuum humilius, foliis obfcure virentibus. Hist. ox. III. 36. R. Syn. 186.

Chamæmelum inodorum f. Cotula non foetida. Bauh. hist. III. 120.

Anglis, Field Feverfew.

Habitat juxta vias, plateas, agros, sed rarius; observed by Mr. Rand along the way to Chelsea. R. Syn. ☉. VIII.

5. MATRICARIA receptaculis hemisphæricis, foliis *maritima*. bipinnatis subcarnosis; supra convexis, subtus carinatis. *Sp. pl.* 891.

Chamæmelum maritimum perenne humilius, foliis brevibus crassis, obscure virentibus. *R. Syn.* 186. *t.* 7. *f.* 1.

Anglis, Sea Feverfew.

Habitat in ora *maritima* Sufflexiæ, ubi loco Cockbush vocato, septem miliaribus Ciceſtria diſtante, ſatis copioſe obſervavit D. Dillenius. *R. Syn.* 4. VII.

A N T H E M I S. 870.

Recept. paleaceum. *Pappus* nullus. *Cal.* hemisphæricus, subæqualis. *Flosculi* radii plures quam quinque.

* *Radio* discolore *f. albo*.

1. ANTHEMIS foliis pinnato-compositis linearibus a- *nobilis*, cutis subvillosis. *Sp. pl.* 894.

Matricaria receptaculis conicis, radiis deflexis, feminibus nudis, squamis calycinis margine æqualibus. *Mat. med.* 401.

Chamæmelum nobile seu Leucanthemum odoratius. *Bauh. pin.* 135.

Chamæmelum odoratissimum repens flore simplici. *Bauh. hiſt.* III. 118. *R. Syn.* 185.

Chamæmelum romanum. *Ger. Em.* 755.

Anglis, sweet-scented Camomile.

Habitat in pratis et pascuis apricis. 4. VII. VIII.

Pharmac. Chamæmeli Flores.

2. ANTHEMIS receptaculis conicis; paleis fetaceis, *Cotula*, feminibus nudis. *Fl. suec.* 767. *Sp. pl.* 894.

Anthemis foliis pinnato decompositis; laciniis linearibus caulibus diffusis, radice annua. *Roy. lugdb.* 172.

Chamæmelum foetidum. *Bauh. pin.* 135. *R. Syn.* 185.

Chamæmelum annuum præcox foetidum, semine aureo. *Hiſt. ox.* III. 36.

Anglis, stinking May-weed.

Habitat in arvis et ad vias. ☉. VI. VII.

arvensis. 3. *ANTHEMIS* receptaculis conicis: paleis fetaceis, seminibus coronato-marginatis. *Sp. pl.* 894. *Fl. suec.* 768.

Chamæmelum inodorum. Baub. pin. 135. *R. Syn.* 184.

Bupthalmum album. Pet. h. br. t. 19. *f.* 8.

Anglis, Corn Camomile.

Habitat in arvis et ad vias. ♂. VII.

maritima. 4. *ANTHEMIS* foliis pinnatis denticulatis carnosif punctatis, caule ramoso prostrato, calycibus tomentosis. *Sp. pl.* 893.

Anthemis maritima perennis, foliis crassis et punctatis. Mich. gen. 33.

Chamæmelum maritimum latifolium ramosissimum, flore albo. Pluk. alm. 97. *R. Syn.* 186.

Matricaria maritima. Baub. pin. 134.

Anglis, Sea Camomile.

Habitat in pratis maritimis. ♀. VIII.

*** Radio concolore f. luteo.*

tinctoria. 5. *ANTHEMIS* foliis bipinnatis ferratis subtus tomentosis, caule corymbofo. *Sp. pl.* 896.

Bupthalmum caule ramoso, foliis pinnatifidis: laciniis linearibus dentatis ferratis, floribus pedunculatis. Hort. cliff. 414.

Bupthalmum tanacetii minoris folio. Baub. pin. 134.

Bupthalmum vulgare. Ger. Em. 747. *R. Syn.* 182.

Bupthalmum matthioli, f. vulgare millefolii foliis. Park. 1370.

Anglis, common Ox-eye.

Habitat in pratis apricis, sed rarius. I found this on a bank near the river Tees, not far from Sogburn, in the bishoprick of Durham. R. Syn. 4. VII.

A C H I L L E A. 871.

Recept. paleaceum. Pappus nullus. Cal. ovatus, imbricatus. Flosculi radii circiter quinque.

Millefoli. 1. *ACHILLEA* foliis bipinnatis nudis: laciniis linearibus

ribus dentatis. *Hort. cliff.* 413. *Sp. pl.* 899. *Fl. suec.* 770.

Millefolium vulgare. *Park.* 693. *R. Syn.* 183. album.
Bauh. pin. 140.

Millefolium terrestre vulgare. *Ger. Em.* 1072.

Anglis, common Yarrow, or Milfoil.

Habitat in pratis et pascuis. 4. V. VII.

Pharmac. Millefolii Herba.

2. ACHILLEA foliis lanceolatis acuminatis argute serratis. *Sp. pl.* 898. *Fl. suec.* 771. *Ptarmica.*

Achillea foliis acutis integris minutissime ferratis. *Vaill. act.* 1720. 316.

Ptarmica. *Ger. Em.* 606. *R. Syn.* 183. vulgaris,
Park. 859.

Dracunculus pratensis, ferrato folio. *Bauh. pin.* 98.

Anglis, Sneezewort, or Goose-tongue.

Habitat in pratis et sylvis subhumidis. 4. VIII.

Pharmac. *Ptarmicæ* Herba.

POLYGAMIA FRUSTRANEA.

CENTAUREA. 880.

Recept. fetosum. *Pappus simplex.* *Cor.* radii infundibuliformes, longiores, irregulares.

* *Cyani calycibus lævibus inermibus.*

1. CENTAUREA calycibus ferratis, foliis linearibus *Cyanus.*
integerrimis : infimis dentatis. *Mat. med.* 408. *Sp.*
pl. 911. *Fl. suec.* 776.

Cyanus. *Bauh. hist.* III. 21. *R. Syn.* 198. vulgaris.

Lob. ic. 546. *Ger. Em.* 732. *Park.* 482.

Cyanus fegetum. *Bauh. pin.* 273.

Anglis, Blue-bottles.

Habitat in arvis. 0. VII

Pharmac. *Cyani Herba*, *Florès.*

2. CENTAUREA calycibus ciliatis, foliis pinnatifidis : *Scabiosa.*
laciniis lanceolatis. *Sp. pl.* 913. *Fl. suec.* 773.
Jacea major. *Ger. Em.* 728. *R. Syn.* 198.

Jacea nigra vulgaris laciniata. Park. 470.
Scabiosa major squamatis capitulis. Bauh. pin. 269.
Anglis, Great Knapweed, or Matfellow.
Habitat in arvis et ad margines agrorum. 4. VI. VII.

- Jacea.* 3. *CENTAUREA* calycibus squamosis, foliis lanceolatis: radicalibus fluuato dentatis, ramis angulatis.
Sp. pl. 914. Fl. juv. 774.
Jacea nigra. Ger. Em. 726. R. Syn. 198. vulgaris.
Park. 468.
Jacea nigra pratensis latifolia. Bauh. pin. 271.
 β *Jacea nigra minor, tomentosa, laciniata R. Syn. 199.*
Anglis, common Knapweed, or Matfellow.
Habitat in pratis et pascuis. 4. VII. VIII.

** *Calcitrapæ: calycinis spinis compositis.*

- Calcitrapa.* 4. *CENTAUREA* calycibus subduplicato-spinosis, foliis sessilibus lateralibus pinnatifidis linearibus dentatis, caule piloso. *Hort. upf. 237. Sp. pl. 917.*
Carduus stellatus. Ger. Em. 1166. R. Syn. 196.
Carduus stellaris seu Calcitrapa vulgaris. Park. 989.
Carduus stellatus foliis papaveris erratici. Bauh. pin. 387.
Anglis, Star-thistle.
Habitat in pratis sterilioribus et juxta vias. ①. VII.
Pharmac. Calcitrapæ Herba, Semina.

- falsitialis.* 5. *CENTAUREA* calycibus duplicato-spinosis solitariis, foliis rameis decurrentibus inermibus lanceolatis; caulinis lyrato-pinnatifidis. *Sp. pl. 917. Roy. lugdb. 140.*
Carduus stellatus luteus foliis cyani. Bauh. pin. 387.
R. Syn. 196.
Carduus falsitialis. Ger. Em. 1166. Park. 989.
Anglis, St. Barnaby's Thistle.
Habitat ad sepes. By the hedges not far from Cirencester in Gloucestershire, D. Bobart. R. Syn. ①. VII.

POLYGAMIA NECESSARIA.

O T H O N N A. 888.

Recept. nudum. Pappus subnullus. Cal. monophyllus, multifidus, subcylindricus.

1. OTHONNA foliis lanceolatis dentato-sinuatis. *Sp. palustris.*
pl. 924. Fl. suec. 778.

Solidago foliis inferioribus lanceolatis ferrato-sinuatis, superioribus integris amplexicaulibus. Hort. cliff.
 410.

Conyza foliis laciniatis. Ger. Em. 483. R. Syn. 174.

Conyza aquatica laciniata. Bauh. pin. 266.

Conyza helenitis foliis laciniatis. Park. 126.

Anglis, Marsh Fleabane.

Habitat in palustribus. In the fen-ditches about Marsh and Chatterefs in the isle of Ely; in the way from Norwich to Yarmouth, a little before you come to Oaklybridge; in the ditches about Pillin-mofs, Lancashire; and at Aberavon in Merionethshire, in Wales. R. Syn. 4. VIII.

2. OTHONNA foliis lanceolatis subdenticulatis villosis, *integrifolia.*
floribus umbellatis. Sp. pl. 925.

Solidago foliis lanceolatis sessilibus integris denticulatis floribus umbellatis, involucro setaceo. Gmel. fib. II. 154. t. 71.

Jacobæa montana lanuginosa angustifolia non laciniata. Bauh. pin. 131. Hist. ox. III. 111. f. 7. t. 12. f. 28.

Jacobæa pannonica folio non laciniato. hist. II. 1056. R. Syn. 178.

Jacobæa angustifolia. Ger. Em. 280. pannonica non laciniata. Park. 668.

Anglis, Mountain Ragwort.

Habitat in pratis montosis. On Gogmagog hills, Newmarket heath, &c. R. Syn. 4. VI.

F I L A G O. 891.

Recept. nudum. Pappus nullus. Cal. imbricatus.
Flosculi feminei inter squamas calycis locati.

- maritima.* 1. FILAGO tomentosa, corymbo subramoso, foliis oblongis obtusis crenatis. *Sp. pl.* 927.
 Santolina corymbo terminali subdiviso, foliis oblongis integerrimis obtusis. *Hort. cliff.* 398.
 Gnaphalium maritimum. *Baub. pin.* 263. *R. Syn.* 180.
 Gnaphalium marinum. *Ger. Em.* 640. *f. Colonaria. Park.* 687.
Anglis, Sea Cudweed.
Habitat in littoribus maritimis. Near Abermeney ferry, in the isle of Anglesea, and between Penfans and St. Michael's mount in Cornwall. R. Syn. ☉. VI. VII.
- germanica* 2. FILAGO calycibus pyramidatis pentagonis axillaribus, caule dichotomo erectiusculo. *Syst. nat.* 1235.
 Gnaphalium caule erecto dichotomo, capitulis globosis axillaribus terminalibusque. *Sp. pl.* 857.
 Gnaphalium vulgare majus. *Baub. pin.* 263.
 Gnaphalium minus s. Herba impia. *Park.* 685. *R. Syn.* 180.
 Filago seu Herba impia. *Ger. Em.* 641.
Anglis, common Cudweed.
Habitat in pratis et pascuis sterilioribus et ad vias publicas. ☉. VI. VII.
- montana.* 3. FILAGO floribus conicis terminalibus axillaribusque, caule dichotomo erecto. *Syst. nat.* 1235.
 Filago minor. *Ger. Em.* 641.
 Gnaphalium caule erecto diviso, floribus pyramidatis lateralibus axillaribusque. *Sp. pl.* 857.
 Gnaphalium minimum. *Leb. ic.* 481. *Baub. hist. III.* 159. *R. Syn.* 181.
 Gnaphalium minus repens. *Baub. pin.* 363.
Anglis, the least Cudweed.
Habitat in pratis et pascuis arenosis. ☉. VI. VII.

4. FILAGO floribus subulatis axillaribus, caule dichotomo erecto. *Syst. nat.* 1235. *gallica.*

Gnaphalium caule erecto dichotomo, floribus pyramidatis axillaribus. *Sp. pl.* 857.

Gnaphalium parvum ramosissimum foliis angustissimis polyspermon. *R. Syn.* 181.

Gnaphalium minimum alterum nostras, stoechadis citrinæ foliis tenuissimis. *Pluk. ph. t.* 298. *f. 2.*

Anglis, Corn Cudweed.

Habitat in arvis glareosis. Among corn in sandy ground about Castle-Heveningham in Essex. *R. Syn.*

☉. VI. VII.

M O N O G A M I A.

J A S I O N E. 896.

Cal. communis decemphyllus. *Cor.* pentapetala, regularis. *Caps.* infera bilocularis.

1. JASIONE. *Sp. pl.* 928. *Fl. suec.* 782. *montana.*

Jasione foliis lineari-lanceolatis obsolete serratis. *Hort. cliff.* 426.

Rapunculus scabiosæ capitulo cæruleo. *Baub. pin.* 92. *Park.* 646. *R. Syn.* 278.

Scabiosa minima hirsuta. *Ger. Em.* 723.

Anglis, hairy Sheep's Scabious.

Habitat in ericetis, pratis montosis passim. 4. ☉.

VI. VII.

L O B E L I A. 897.

Col. quinquefidus. *Corolla* monopetala, irregularis.

Caps. infera, bilocularis.

1. LOBELIA foliis linearibus bilocularibus integerri- *Dortman-*
mis caule subnudo. *Sp. pl.* 929. *Fl. suec.* 783. *na.*

Dortmanna palustris, floribus sparsis pendulis. *Rudb. act. ups. an.* 1720. *p. 97. t. 2.*

Gladeolus lacustris Dortmanni. *Clus. cur.* 40. *R. Syn.* 287.

Gladeolus lacustris Clusii, f. Leucojum palustre, flore subcæruleo. *Park.* 1250.

Leucojum

Leucojum palustre, flore subcœruleo. *Bauh. pin.* 41.
Anglis, Water Gladiole.

Habitat in omnibus Cambriæ montibus Dr. Richardson.
R. Syn. in lacubus Westmorlandiæ copiose. 4. VII.
VIII.

V I O L A. 898.

*Cal. pentaphyllus. Cor. pentapetala, irregularis, pone
cornuta. Caps. supera trivalvis, unilocularis.*

* *Acaules.*

odorata. 1. *VIOLA acaulis, foliis cordatis, stolonibus reptantibus. Sp. pl.* 634. *Fl. suec.* 784.

Viola martia purpurea. Bauh. hist. II. 542. *R. Syn.*
364. *flore simplici. Bauh. pin.* 199.

Viola nigra five purpurea. Ger. Em. 850.

Viola simplex martia. Park. 755.

β *Viola martia alba. Bauh. pin.* 199.

Anglis, Purple sweet Violet. β white Violet.

Habitat in sepibus et aggeribus fossarum. 4. III.

Pharmac. Violæ Flores.

birta. 2. *VIOLA acaulis, foliis cordatis piloso-hispidis. Sp. pl.*
934.

Viola martia major hirsuta inodora. Hist. ox. II. 475.
f. 5. t. 35. f. 4. R. Syn. 365.

Viola trachelii folio vulgo. R. hist. 1051.

Anglis, hairy Violet.

Habitat in sylvis et sepibus passim. 4. III.

palustris. 3. *VIOLA acaulis foliis reniformibus. Hall. helv.* 501.
Sp. pl. 934. *Fl. suec.* 786.

Viola palustris rotundifolia glabra. Hist. ox. II. 475.
f. 5. t. 35. f. 5. Plot. ox. 144. *t. 9. f. 2. R. Syn.*
364.

β *Viola rubra striata Eboracensis. Park.* 755.

Anglis, Marsh Violet.

Habitat in pratis humidis, paludibus, cespitosis, turfosis. On the banks of the river Chervil, between Oxford and Water-Eaton. On Oakenshaw-moor and Roadeshall woods. R. Syn. supra Sheriey common prope Croydon; in ericeto Hampstediano prope Londinum. 4. VI.

** *Cau-*

** *Caulescentes.*

4. VIOLA caule demum adscente foliis oblongo-cordatis. *Sp. pl.* 935. *Fl. suec.* 785.
 Viola caulibus adscendentibus floriferis, foliis cordatis. *Hort. cliff.* 427.
 Viola martia inodora sylvestris. *Bauh. pin.* 199. *R. Syn.* 364.
 Viola canina sylvestris. *Ger. Em.* 851.
 Viola sylvestris. *Park.* 755.
 β Viola canina flore albo. *Merr. pin.*
 γ Viola canina minor. *R. Syn.* 364. t. 24. f. 1.
Anglis, Dog's Violet.
 Habitat in ericetis. 4. IV.
5. VIOLA caule subtrifloro, foliis cordatis profunde ferratis. *alpina.*
 Viola martia alpina folio tenello circinato. *R. Syn.* 366.
Anglis, Welch Violet.
 Habitat in rupe Clogwyn y Garnedh, supra lucum Phynonlas; D. Lhwyd. *R. Syn.* 4.
6. VIOLA caule triquetro diffuso, foliis oblongis incisis, stipulis dentato-pinnatis, lacinia suprema oblonga crenata petiolata. *tricolor.*
 Viola caule triquetro diffuso, foliis oblongis incisis stipulis dentatis. *Sp. pl.* 935. *Fl. suec.* 791.
 Viola tricolor. *Ger. Em.* 854. *R. Syn.* 365. major et vulgaris. *Park.* 756.
 Viola tricolor hortensis repens. *Bauh. pin.* 199.
 β Viola bicolor arvensis. *Bauh. pin.* 200.
 γ Viola bicolor arvensis, flore cæruleo et candido. *Tourn.* 421.
Anglis, Pansies, or Heart's-ease, vulgo three Faces under a Hood.
 Habitat in arvis. 5. V.—IX.
7. VIOLA caule subtriquetro diffuso, foliis subcordatis, crenatis, stipulis dentatis radice repente. *lutea.*
 Viola montana lutea grandiflora nostras. *R. Syn.* 365.
 Viola flammea lutea. *Ger. Em.* 851.
 Viola montana lutea grandiflora. *Bauh. pin.* 200.
Anglis,

Anglis, yellow Violet, or Pansies.

Habitat in montosis pascuis tam septentrionalibus, tum Walicis, solo putrido et palustri multoties observavimus. R. Syn. 4. V.—VIII.

IMPATIENS. 899.

Cal. diphyllus. *Cor.* pentapetala, irregularis, nectario cuculato. *Capf.* supera, quinquevalvis.

noli me tangere.

I. IMPATIENS pedunculis multifloris solitariis, foliis ovatis, geniculis caulinis tumentibus. *Sp. pl.* 938.

Fl. suec. 792.

Balsamine lutea, f. Noli me tangere. *Bauh. pin.* 306. R. Syn. 316.

Perficaria filiquosa. *Ger. Em.* 440.

Mercurialis sylvestris, Noli me tangere dicta, five Perficaria filiquosa. *Park.* 296.

Anglis, Quick in the Hand. Touch me not.

Habitat ad in umbrosis humidis et ad rivulos. On the banks of Winandermere, near Ambleside. R. Syn. prope Rydall-hall, et alibi in Westmorlandia. ☉. VIII.

Classis

Classis XX.

GYNANDRIA
DIANDRIA.

ORCHIS. 900.

Nectarium corniforme pone florem.* *Bulbis indivisis.*

1. ORCHIS bulbis indivisis, nectarii labio lanceolato in-*bifolia.* x
tegerimo : cornu longissimo, petalis patentibus.
Sp. pl. 939. *Fl. suec.* 793.
Orchis radicibus conicis, labello lingulato simplicissimo. *Hall. helv.* 266.
Orchis alba bifolia minor, calcari oblongo. *Bauh. pin.* 83. *R. Syn.* 380. *Vaill. parif.* 151. t. 30. f. 7.
Orchis psychodes. *Ger. Em.* 211.
Orchis sphegodes, f. Testiculus vulpinus primus. *Park.* 1351.
β Orchis hermaphroditica bifolia. *Bauh. hist.* II. 772.
Anglis, Butterfly Orchis.
Habitat in *pascuis et dumetis passim.* 4. V. VI.
2. ORCHIS bulbis indivisis, nectarii labio quadrilobo *mascula.*
crenulato : cornu obtuso, petalis dorsalibus reflexis.
Fl. suec. 795.
Orchis morio mas, foliis maculatis. *Bauh. pin.* 81.
Park. 1346. *Vaill. parif.* 151. t. 31. f. 12.
Cynoforchis morio mas. *Ger. Em.* 208.
Anglis, Male Fool-stones.
Habitat in *pratis et pascuis.* 4. V.
Pharmac. Satyrii Radix.
3. ORCHIS bulbis indivisis, nectarii labio quadrifido *Morio.*
crenulato : cornu obtuso, petalis omnibus conni-
ventibus. *Sp. pl.* 940. *Fl. suec.* 794.

Orchis

Orchis radicibus subrotundis, cucullo clauso lineato, labello ampliter trifido crenulato. *Hall. helv.* 266.

Orchis morio fœmina. *Baub. pin.* 82. *Park.* 1347.

R. Syn. 377. *Vaill. paris.* 149. t. 31. f. 13, 14.

Cynoforchis morio fœmina. *Ger. Em.* 208.

Anglis, Female Fool-stones.

Habitat in pratis et pascuis subhumidis. 4. V. VI.

- × *pyramidalis*. 4. ORCHIS bulbis indivisis, nectarii labio trifido æquali integerrimo : cornu longo, petalis sub lanceolatis. *Fl. suec.* 798.

Orchis purpurea spica congesta pyramidalis. *R. Syn.* 377. t. 18.

Anglis, Purple late-flowering Orchis.

Habitat in pascuis siccioribus et cretaceis frequens. 4. VI. VII.

- × *ustulata*. 5. ORCHIS bulbis indivisis, nectarii labio quadrifido punctis scabro : cornu obtuso, petalis distinctis. *Fl. suec.* 797. *Sp. pl.* 941.

Orchis pannonica IV. *Clus. hist.* 268. *R. Syn.* 377.

Cynoforchis militaris pratensis humilior. *Baub. pin.* 81.

Cynoforchis militaris pannonica. *Park.* 1345.

Cynoforchis minor pannonica. *Ger. Em.* 207.

Anglis, little purple-flowered Orchis.

Habitat in pascuis siccioribus, et sterilibus, præsertim cretaceis. 4. V. VI.

- coriophora*. 6. ORCHIS bulbis indivisis, nectarii labio trifido reflexo crenato : cornu brevi, petalis conniventibus. *Sp. pl.* 940.

Orchis odore hirci minor. *Baub. pin.* 82. *Vaill. paris.* 149. t. 30. f. 30, 31, 32. *Hall. it. herc.* 7.

Orchis barbata foetida minor flore albo. *R. Syn.* 376.

Tragorchis minor, flore fuliginoso. *R. hist.* 1213.

Anglis, the lesser Lizard Flower.

Habitat

- purpurea*. 7. ORCHIS bulbis indivisis, nectarii labio barbato quinquelobo lateralibus obtusis, crenulatis, medio brevissimo acuto, petalis confluentibus.

Orchis

Orchis magna latis foliis, galea fusca vel nigricante.

Baub. hist. II. 759. R. Syn. 378. t. 19. f. 2.

Anglis, purple Man-Orchis.

Habitat in collibus cretaceis. At Northfleet near Gravesend, Mr. J. Sherard. *R. Syn. 4. VI.*

8. ORCHIS bulbis indivisis, nectarii labio quinquefido *militaris*.
lineari punctis scabro, petalis confluentibus.

Orchis bulbis indivisis, nectarii labio quinquelobo punctis scabro: cornu obtuso, petalis confluentibus. *Sp. pl. 941. Fl. suec. 796.*

Orchis galea et alis fere cinereis. *Baub. hist. II. 755. R. Syn. 378.*

Cynorchis latifolia hiante cucullo major. *Baub. pin. 80.*

Cynorchis latifolia minor. *Park. 1344.*

Cynorchis major altera. *Ger. Em. 205.*

Anglis, the Man Orchis.

Habitat in pratis et pascuis solo cretaceo. On Cawsham hills, by the Thames-side, not far from Reading in Berkshire. *R. Syn. Near the old chalk-pit near the paper-mill at Harefield, plentifully. Blackf. spec. bot. 4. VI.*

** *Bulbis palmatis.*

9. ORCHIS bulbis subpalmatis rectis, nectarii cornu con-*latifolia.* x
nico: labio trilobo lateralibus reflexis, bracteis flore longioribus. *Sp. pl. 941. Fl. suec. 801.*

Orchis palmata pratensis *latifolia*, longis calcaribus. *Baub. pin. 85. R. Syn. 380. Vaill. parif. 152. t. 31. f. 1, 2, 3, 4, 5.*

Orchis palmata mas f. *Palma Christi* mas. *Park. 1356.*

Palma Christi mas. *Ger. Em. 220.*

Anglis, Male-handed Orchis.

Habitat in pratis et pascuis. 4. V. VI.

10. ORCHIS bulbis palmatis patentibus, nectarii cornu *maculata.* x
germinibus brevioribus: labio plano, petalis dorsali-
bus erectis. *Sp. pl. 942. Fl. suec. 800.*

Orchis

*Orchis palmata pratensis maculata. Baub. pin. 85.
R. Syn. 381.*

Orchis palmata foemina f. palma Christi foemina, foliis maculatis. Park. 1357.

Palma Christi foemina. Ger. Em. 220.

Anglis, Female-handed Orchis.

Habitat in pratis et pascuis humidioribus. 4. VI.

× *conopsea.* 11. ORCHIS bulbis palmatis, nectarii cornu fetaceo germinibus longiore, labio trifido, petalis duobus patentissimis. *Syst. nat. 1243. Sp. pl. 942. Fl. suec. 790.*

Orchis palmata minor calcaribus oblongis. Baub. pin. 85. Rudb. elyf. II. 212. f. 5. Vaill. parif. t. 30. f. 8, 9, 10.

Orchis palmata minor flore rubro. Park. 1358.

Serapias minor nitente flore. Ger. Em. 222.

Anglis, red-handed Orchis.

Habitat in pratis et pascuis. 8. VI.

*** *Bulbis fasciculatis.*

abortiva. 12. ORCHIS bulbis fasciculatis filiformibus, nectarii labio ovato integerrimo. *Sp. pl. 943. Dalib. parif. 275.*

Orchis abortiva violacea. Baub. pin. 86.

Limodorum Austriacum. Clus. pann. 242. R. Syn. 383.

Nidus avis purpureus. Park. 1362.

Nidus avis flore et caule purpuro-violaceo. Ger. Em. 228.

Anglis, purple Bird's-nest.

Habitat in pratis siccioribus. In Southamptoniæ comitatu prope Alton oppidum invenit, D. Goodyer.

R. Syn. 4. V.

SATYRIUM. 901.

Nectarium scrotiforme f. inflato-didymum, pone florem.

irsinum. 1. SATYRIUM bulbis indivisis, foliis lanceolatis, nectarii labio trifido: intermedia lineari obliqua præmorfa. *Sp. pl. 944.*

Orchis

Orchis barbata odore hirci, brevior latioreque folio. *Bauh. pin.* 82.

Orchis barbata foetida. *Bauh. hist. II.* 756. *R. Syn.* 376.

Tragorchis maxima, et T. vulgaris. *Park.* 1348.

Tragorchis maximus et T. mas. *Ger. Em.* 225.

Anglis, the Lizzard Flower, or Goat-stones.

Habitat in cretaceis. Circa Dartford et Crayford in Cantio sed rarius. *R. Syn.* 4. VI. VII.

2. SATYRIUM bulbis palmatis, foliis oblongis obtusis, *viride*, nectarii labio lineari trifido intermedia obsoleta. *Sp. pl.* 944. *Fl. suec.* 304.

Orchis palmata flore viridi. *Bauh. pin.* 86. *pr.* 30.

Orchis palmata minor flore luteo-viridi. *R. Syn.* 381.

Serapias batrachites altera. *Ger. Em.* 224.

Anglis, Frog Satyrion, or Orchis.

Habitat in pascuis siccioribus. 4. V. VI.

3. SATYRIUM bulbis palmatis, foliis oblongis, nectarii labio trifido æquali. *fuscum*.

Anglis, brown Satyrion.

Habitat in prato montano Hellse-fellnab dicto prope Kendal in Westmorlandia. 4. VIII.

4. SATYRIUM bulbis fasciculatis, foliis lanceolatis, *albidum*, nectarii labio trifido acuto : intermedia majore. *Sp. pl.* 944. *Fl. suec.* 806.

Orchis palmata thyrsos specioso, longo, dense stipato, ex viridi albente. *R. Syn.* 382.

Orchis palmata alpina, spica densa albo-viridi. *Hall. herb.* 68.

Pseudo-orchis alpina, flore herbaceo. *Mich. gen.* 80. t. 26.

Anglis, white Satyrion.

Habitat in pratis humidis in Cambria, et circa Malham in comitatu Eboracensi, D. Richardson. *R. Syn.* 4. VI.

OPHRYS 902.

Nectarium subtus subcarinatum.* *Bulbis ramosis.*

- Nidus avis.* 1. OPHRYS bulbis fibroso-fasciculatis, caule vaginato, nectarii labio bifido. *Sp. pl.* 942. *Fl. suec.* 815.
Nidus avis. *Lob. ic.* 195. *Lugd.* 673. *R. Syn.* 382.
Orchis abortiva fusca. *Bauh. pin.* 86.
Orchis abortiva rufa, f. *Nidus avis.* *Park.* 1362.
Satyrion abortivum f. *Nidus avis.* *Ger. Em.* 228.
Anglis, Bird's-nest.
Habitat in sylvis et umbrosis. *In sylva* Charlton wood dicta in Cantio. 4. V.
- spiralis.* 2. OPHRYS bulbis aggregatis oblongis, caule subfolioso, floribus secundis, nectarii labio indiviso. *Sp. pl.* 945.
Orchis spiralis albo odorata. *Bauh. hist. II.* 769. *R. Syn.* 378.
Triorchis *Ger. Em.* 218. *alba odorata minor.* *Bauh. pin.* 87. *Park.* 1354.
Anglis, triple Ladies Traces.
Habitat in pascuis siccioribus. 4. VIII.
- ovata.* 3. OPHRYS bulbo fibroso, caule bifolio, foliis ovatis, nectarii labio bifido. *Sp. pl.* 946. *Fl. suec.* 808.
Ophrys bifolia. *Ger. Em.* 402. *Bauh. pin.* 87.
Bifolium sylvestre vulgare. *Park.* 504. *R. Syn.* 385.
Anglis, common Twayblade.
Habitat in sylvis et dumetis humidis. 4. V. VI.
- cordata.* 4. OPHRYS bulbo fibroso, caule bifolio, foliis cordatis, nectarii labio bifido.
Ophrys bulbo fibroso, caule bifolio, foliis cordatis. *Sp. pl.* 946. *Fl. suec.* 809.
Ophrys minima. *Bauh. pin.* 87. *pr.* 31.
Bifolium minimum. *Bauh. hist. III.* 534. *R. Syn.* 385.
Anglis, the least Twayblade.
Habitat in ericetis humidis, præsertim solo turfofo, in comita-

comitatibus Eboracensi, Lancastriensi et Westmorlandico passim. 4. VII.

5. OPHRYS bulbo fibroso, scapo vaginato, nectarii labio *palustris*.
bio integro.

Bifolium palustre. *Park.* 505. *R. Syn.* 385.

Orchis bifolia minor palustris. *Pluk. ph. t.* 247. *f.* 2.
Anglis, Marsh Twayblade.

Habitat in paludibus. Found by Mr. Dent and Mr. Dale, on the boggy and fenny grounds near Gamlingay in Cambridgeshire; by Parkinson on the low wet grounds between Hatfield and St. Albans; and in divers places in Romney marsh. By Mr. Dubois at Hursthill, Tunbridge-wells. *R. Syn.* 4. VII. VIII.

** *Bulbis rotundis.*

6. OPHRYS bulbo subrotundo, scapo subnudo pentagono, foliorum apicibus scabris. nectarii labio integro. *Sp. pl.* 947. *Fl. suec.* 813. *paludosa*.

Orchis minima bulbosa. *R. Syn.* 375.

Anglis, the least Orchis.

Habitat

7. OPHRYS bulbo globoso, scapo nudo, nectarii labio trifido. *Sp. pl.* 947. *Fl. suec.* 810. *Monorchis*.

Orchis odorata moschata, f. *Monorchis*. *Bauh. pin.* 84. *R. Syn.* 378.

Orchis pusilla odorata. *Park.* 1354.

Anglis, yellow or Musk Orchis.

Habitat in pascuis sterilioribus, sed rarus. 4. VII. †

8. OPHRYS bulbo subrotundo, scapo nudo, foliis lanceolatis, nectarii labio integro, petalis dorsalibus linearibus. *Sp. pl.* 946. *Fl. suec.* 811. *lilifolia*.

Ophrys scapo nudo, foliis radicalibus ovato oblongis dimidii scapi longitudine. *Gron. virg.* 185.

Pseudo-orchis bifolia palustris. *R. Syn.* 382.

Chamæorchis lilifolia. *Bauh. pin.* 84.

Chamæorchis latifolia Zelandica. *Park.* 1354.

Anglis, Dwarf Orchis.

Z. 2

Ha-

† Ophrys Monorchis. Knight's ink on a bank at the edge of a little Coppice being at the upper side of a field, which is on the right hand side of the way from Andover to Newbury, particularly.

GYNANDRIA DIANDRIA.

Habitat in palustribus. On Hinton and Teversham moors near Cambridge. *R. Syn.* 4. VII.

anthropophora. 9. OPHRYS bulbis subrotundis, scapo folioso, nectarii labio lineari tripartito : medio elongato bifido. *Sp. pl.* 948.

Orchis anthropophora oreades. Col. ec. I. 320. R. Syn. 379. fœmina. Park. 1348.

Orchis flore nudi hominis effigiam repræsentans, fœmina. Bauh. pin. 82.

Anglis, green Man Orchis.

Habitat in cretaceis. Found by Mr. Dale in an old gravel-pit at Dalington near Sudbury ; at Northfleet and Greenhithe in Kent, plentifully. *R. Syn.* 4. VI.

muscifera. 10. OPHRYS bulbis subrotundis, scapo folioso, nectarii labio quadrifido.

Orchis myodes galea et alis herbidis. Bauh. hist. II. 767. R. Syn. 379.

Orchis muscæ corpus referens minor, vel galea et alis herbidis. Bauh. pin. 83.

Orchis myodes minor. Park. 1352.

β *Orchis myodes major. Park. 1351.*

Anglis, Fly Orchis.

Habitat in pratis et pascuis passim. 4. VI.

* *apifera.* 11. OPHRYS bulbis subrotundis, scapo folioso, nectarii labio quinquelobo, lobis subtus inflexis.

Orchis fuciflora galea et alispurpurascentibus. Bauh. hist. II. 766. R. Syn. 379.

Orchis f. Testiculus vespinus, 2. Sphegodes. Ger. Em. 212.

Orchis fucum referens major, foliolis superioribus candidis et purpurascentibus. Bauh. pin. 83.

Orchis sphegodes seu fucum referens. Park. 1350.

β *Orchis five Testiculus sphegodes hirsuto flore. Bauh. hist. II. 767. R. Syn. 380.*

Anglis, Bee Orchis.

Habitat in pratis et pascuis siccioribus. 4. VI.

CYPRIPEDIUM. 906.

Nectarium ventricosum, inflatum, cavum.

1. CYPRIPEDIUM radicibus fibrosis, foliis ovato lanceolatis caulinis. *Sp. pl.* 951. *Fl. suec.* 820.
Calceolus Mariæ. *Ger. Em.* 443. *R. Syn.* 385.
Helleborine flore rotundo seu Calceolus. *Bauh. pin.* 187.
Elleborine major seu Calceolus Mariæ. *Park.* 217.
Anglis, Ladies Slipper.
Habitat in sylvis et dumetis, sed rarius. In sylva inter Clapham et Ingleborough sed sparsim. 4.
VII.

SERAPIAS. 903.

Nectarium ovatum, gibbum : labio ovato.

1. SERAPIAS bulbis fibrosis, nectarii labio obtuso cretato, petalis æquali, fructibus pendulis. *latifolia.*
Serapias caule multifolio multifloro. *Hort. cliff.* 429.
Helleborine latifolia montana. *Bauh. pin.* 186. *R. Syn.* 383.
Helleborine. *Dod. pempt.* 384. *Ger. Em.* 442.
Elleborine flore viridante. *Park.* 218.
β Helleborine altera atro-rubente flore. *Bauh. pin.* 186. *R. Syn.* 383.
γ Helleborine palustris nostras. *R. Syn.* 284.
Anglis, broad-leaved Bastard Hellebore.
Habitat in sylvis et sepibus. γ in palustribus. 4. VII.
VIII.
2. SERAPIAS bulbis fibrosis, nectarii labio obtuso petalis brevioribus, fructibus erectis. *longifolia.*
Serapias radicibus fibrosis, nectarii labio obtuso petalis brevioribus, foliis ensiformibus. *It. oel.* 144.
Helleborine flore albo. *Ger. Em.* 442. *R. Syn.* 383. vel *Damafonium montanum latifolium.* *Bauh. pin.* 187.
Elleborine minor flore albo. *Park.* 218.
β Helleborine latifolia, flore albo clauso. *R. Syn.* 384.

γ Helleborine foliis prælongis angustis acutis. *R. Syn.* 384.

δ Helleborine montana angustifolia purpurascens. *Bauh. pin.* 187.

Anglis, white-flowered Bastard Hellebore.

Habitat in sylvis et dumetis. α in the woods near Stoken-Church, Oxfordshire, *R. Syn.* β in a Beachwood, just below the duke of Bedford's seat at Cheyneis, Buckinghamshire. *Blackf. spec. bot.* α, γ, δ, circa Clapham et Ingleton, in comitatu Eboracensi. 4. VIII.

P O L Y A N D R I A.

A R U M. 915.

Spatha monophylla, cucullata. *Spadix* supra nudus, inferne femineus, medio stamineus.

maculatum. 1. ARUM acaule, foliis hastatis integerrimis, spadice clavato. *Hort. cliff.* 8. *Sp. pl.* 966. *Roy. lugdb.* 8. *Arum. Bauh. hist.* II. 783. *R. Syn.* 266. vulgare. *Ger. Em.* 834. maculatum et non maculatum. *Park.* 373.

Arum vulgare maculatum. *Bauh. pin.* 195.

Anglis, Wake-Robin, Cuckow-pint.

Habitat in umbrosis et ad sepes. 4. V.

Pharmac. Ari Radix.

Z O S T E R A. 919.

Spadix linearibus, altero latere fructifero. *Calyx* nullus. *Cor.* nulla. *Stam.* alterna. *Semina* solitaria alterna.

marina. 1. ZOSTERA. *It. W—goth.* 166. i. 4. f. 1. *Sp. pl.* 968.

Alga. Ger. Em. 1569 *R. Syn.* 52. angustifolia vitriatorum. *Bauh. pin.* 364. *hist.* III. 794. *R. Syn.* 53.

Fucus marinus sive alga, marina graminea. *Park.* 1242.

β *Fucus*

- β Fucus f. alga marina graminea angustifolia seminifera
ramosior. *R. Syn.* 52.
γ Fucus marinus seu Alga marina graminea minor.
R. Syn. 52.
δ Potamogeton marinum in utriculis epiphyllispermon
minus. *R. Syn.* 53.
Anglis, Grafs Wrack.
Habitat in littoribus maritimis. 24.

X 4

Classis

Classis XXI.

MONOECIA

MONANDRIA

ZANNICHELLIA. 920.

Masculi. *Cal.* o. *Cor.* o.Feminei. *Cal.* monophyllus. *Cor.* o. *Germina* circiter quatuor. *Sem.* totidem.*palustris.* 2. ZANNICHELLIA. *Fl. lapp.* 321. *succ.* 824. *Sp. pl.* 969.Zannichellia palustris major foliis gramineis acutis. *Mich. gen.* 71. *t.* 34. *f.* 1. 2.Aponogiton aquaticum graminifolium, staminibus singularibus. *Pont. anth.* 117. *R. Syn.* 135.Potamogitoni similis Graminifolia ramosa et ad genicula polyceratos. *Pluk. ph. t.* 102. *f.* 7.*Anglis*, horned Pondweed.

Habitat in fossis et stagnis frequens. ☉. VII.

DIANDRIA

LEMNA. 923.

Masculi. *Cal.* monophyllus. *Cor.* nulla.Fem. *Cal.* monophyllus. *Cor.* nulla. *Stylus* unus. *Capf.* unilocularis.*trifolca.* 1. LEMNA foliis petiolaris lanceolatis. *Fl. lapp.* 370. *succ.* 828. *Sp. pl.* 970.Lenticula aquatica trifolca *Baub. pin.* 362. *R. Syn.* 129.Hederula aquatica. *Ger. Em.* 830.Ranunculus hederaceus aquaticus. *Park.* 1260.*Anglis*,

Anglis, Ivy-leaved Duck's-meat.

Habitat in *fossis et stagnis*. 4. V. VI.

2. LEMNA foliis sessilibus utrinque planiusculis, radi-
cibus solitariis. *Hort. cliff.* 417. *Fl. suec.* 827. *Sp. minor.*
pl. 970.

Lens palustris. *Ger. Em.* 829. *Bauh. hist.* III. 784.

R. Syn. 129. t. 4. f. 1.

Lens palustris vulgaris. *Bauh. pin.* 362.

Lens palustris f. aquatica vulgaris. *Park.* 1262.

Anglis, the least Duck's-meat.

Habitat in *stagnis*. 4. VI.

3. LEMNA foliis sessilibus, radicibus confertis. *Hort. cliff.* 477. *Fl. suec.* 825. *Sp. pl.* 970. *polyrhiza.*

Lenticula palustris major. *Comm. Cat. pl. Holl.* R.

Syn. 129. t. 4. f. 2.

Anglis, greater Duck's-meat.

Habitat in *fossis*. 6. VI. VII.

TRIANDRIA.

TYPHA. 924.

Masc. *Amentum* cylindricum. *Cal.* triphyllus.
Cor. o.

Fem. *Amentum* cylindricum infra masculos. *Cal.*
capillo villosa. *Cor.* o. *Sem.* unicum, infidens
pappo capillari.

1. TYPHA foliis subensiformibus, spica mascula femi-
neaque approximatis. *It. scan.* 168. *Sp. pl.* 971. *latifolia.*
Fl. suec. 831.

Typha. *Ger. Em.* 46. palustris major. *Bauh. pin.*
20. *hist.* II. 539. *R. Syn.* 436.

Typha palustris maxima. *Park.* 1204.

Anglis, great Cat's-tail, or Reed-mace.

Habitat ad *ripas fluviorum, piscinarum, paladium fre-*
quens. 4. VII.

2. TYPHA foliis femicylindricis, spica mascula femine-
aque remotis. *It. scan.* 198. *Fl. suec.* 830. *Sp. pl. lia.*
971.

Ty-

MONOECIA TRIANDRIA.

Typha palustris media. *Bauh. hist. II.* 540. *R. Syn.* 436.

Typha palustris clava gracili. *Bauh. pin.* 20.

Typha minor. *Park.* 1204.

Anglis, narrow-leaved Cat's-tail.

Habitat in fossis, et stagnis. 4. VII.

SPARGANIUM. 925.

Masc. *Amentum* subrotundum. *Cal.* triphyllus.

Cor. o.

Fem. *Amentum* subrotundum. *Cal.* triphyllus. *Cor.*

o. *Stigm.* bifidum. *Drupa* exsucca, disperma.

erectum. 1. SPARGANIUM foliis triquetris erectis: *Syst. nat.* 1260. *Sp. pl.* 971. *Fl. suec.* 831. *lapp.* 345.

Sparganium ramosum. *Park.* 1205. *Ger. Em.* 45.

Bauh. pin. 15. *R. Syn.* 437.

β *Sparganium non ramosum.* *Bauh. pin.* 15.

Anglis, great Bur-reed.

Habitat in fossis, paludibus et ad ripas fluviorum.

4. VII.

natans. 2. SPARGANIUM foliis decumbentibus planis. *Sp. pl.* 971. *Fl. suec.* 832.

Sparganium foliis complanatis natantibus, spicis paucissimis. *Hall. helv.* 260.

Sparganium minimum. *R. hist.* 1910. *syn.* 437.

Anglis, the least Bur-reed.

Habitat in stagnis et fluviis leniter fluentibus, sed rarius. 4. VII.

CAREX. 928.

MASC. *Amentum* imbricatum. *Cal.* monophyllus.

Cor. o.

FEM. *Amentum* imbricatum. *Cal.* monophyllus.

Cor. o. *Nectarium* inflatum, tridentatum. *Stigm.*

tria. *Sem.* triquetrum, intra nectarium..

* *Spica unica simplici.*

dioica. 1. CAREX spica simplici dioica. *Hort. cliff.* 438. *Fl. suec.* 833. *Sp. pl.* 972.

Gramen cyperoides minimum spica simplici casta.

Hist. ex. III. 244. *f. 8. t. 12. f. 22.*

Gramen cyperoides minus ranunculi capitulo longiore. *R. Syn.* 425. *An-*

Anglis, small Carex.

Habitat in paludibus turfosis. 4. VI.

2. CAREX spica simplici androgyna ovata: superne mascula, capsulis imbricatis patulis. *Syst. nat.* 1261. *capitata.*

Gramen cyperoides minimum ranunculi capitulo rotundo. *Hist. ox. III.* 245. f. 8. t. 12. f. 36. R. *Syn.* 425.

Gramen cyperoides spica echinata simplici. *Mer. pin.*

Anglis, round-headed Carex.

Habitat in paludibus turfosis, in Sherley common, prope Croydon. 4. VI.

3. CAREX spica simplici androgyna: superne mascula, capsulis divaricatis retroflexis. *Syst. nat.* 1261. *Sp. pl.* 972. *pulicaris.*

Gramen cyperoides minimum, feminibus deorsum reflexis puliciformibus. R. *Syn.* 424. *Pluk. phyt. t.* 34. f. 10. *Hist. ox. III.* 244. f. 8. t. 12. f. 21.

Anglis, Flea Carex.

Habitat in paludibus limosis. In ericeto Hampstediano et Sherley common prope Croydon. 4. VI.

** *Spicis androgynis.*

4. CAREX racemo composito, spiculis androgynis. *paniculata* *Cent. pl.* 93. *Syst. nat.* 1262.

Gramen cyperoides palustre elatius, spica longiore laxa. R. *Syn.* 422. *Hist. ox. III.* 244. f. 8. t. 12. f. 23.

Anglis, Panicked Carex.

Habitat in palustribus putridis, et aquosis, in sylvula Hampstead wood dicta prope Hampstedium. 4. VI. VII.

5. CAREX spica composita subdisticha, spiculis ovatis imbricatis androgynis, foliolo longiori instructis, culmo triquetro. *disticha.*

Gramini cyperoidi ex monte Ballon simile, spica totali e pluribus spicis composita. R. *Syn.* 423. *Pluk. ph. t.* 34. f. 7.

Gramen cyperoides elegans multifera spica. *Park.* 1172. *Gramen*

- β Gramen cyperoides elegans spica composita molli.

Pluk. ph. t. 34. f. 4.

Anglis, soft Carex.

Habitat in palustribus. 4. V. VI.

divisa.

6. CAREX spica composita spiculis rotundis confertis androgynis, foliolo longiori instructis, culmo subtereti.

Gramen cyperoides palustre majus spica divulsa. *Bauh. pin. 6.*

Gramen cyperoides ex monte Ballon spica divulsa. *Bauh. hist. II. 498. R. Syn. 423.*

Gramen cyperoides parvum. *Ger. Em. 21.*

Gramen cyperoides palustre minus. *Park. 1287.*

Anglis, Marsh Carex.

Habitat in palustribus, sed rarius. In the meadows near the Hithe at Colchester in Essex, Mr. Newton ; by Hithe in Kent, Mr. J. Sherard. R. Syn. 4. VII.

leporina.

7. CAREX spica composita, spiculis ovatis sessilibus approximatis androgynis nudis. *Sp. pl. 973. Fl. suec. 837.*

Carex spicis ternis sessilibus confertis androgynis. *Fl. lapp. 322.*

Gramen cyperoides spica e pluribus spicis brevibus mollibus composita. *R. Syn. 422. Sch. agr. 456. t. 10. f. 15.*

Gramen cyperoides palustre majus spica divisa. *Bauh. pin. 6. Hist. ex. III. 244. f. 8. t. 12. f. 29.*

Anglis, naked Carex.

Habitat in palustribus et aquosis 4. VI. VII.

arenaria.

8. CAREX spica composita, spiculis androgynis : inferioribus remotioribus foliolo longiori instructis, culmo triquetro. *It. oel. 139. Sp. pl. 973. Fl. suec. 835.*

Gramini cyperoides ex monte Ballon simile humilius. in maritimis et arenosis nascens. *R. Syn. 423. Pluk. ph. t. 34. f. 7.*

Gramen

Gramen cyperoides elegans multifera spica. *Park.*

1172.

Anglis, Sea Carex.

Habitat in arenosis maritimis passim. 4. VI. VII.

9. CAREX spica composita disticha nuda, spiculis an-
drogynis oblongis contiguis patentibus, culmo nu-
do. *Cent. pl.* 91. *Syst. nat.* 1262.

Gramen cyperoides palustre elegans spica composita
asperiore. *R. Syn.* 423.

Anglis, rough Carex.

Habitat in palustribus et aquosis sed rarius. 6. VI.

10. CAREX spica supradecomposita inferne laxiore, spi-
culis androgynis glomeratis : superne masculis. *Fl.*
suec. 838. *Sp. pl.* 973.

Gramen cyperoides palustre majus, spica compacta.
Bauh. pin. 6. *R. Syn.* 423. *Hist. ox. III.* 244. f.
8. t. 12. f. 24.

Gramen palustre cyperoides. *Ger. Em.* 21.

Gramen cyperoides palustre majus. *Park.* 1266.

Anglis, great Carex.

Habitat ad ripas fluviorum. 4. VII.

11. CAREX spiculis subrotundis approximatis andro-
gynis capsulis ovatis acutis.

Gramen cyperoides parvum. *Ger. Em.* 18.

Gramen cyperoides spicis minoribus minusque com-
pactis. *Bauh. pin.* 6. pr. 13. *hist. II.* 497.

Gramen cyperoides spicatum minus. *R. Syn.* 424.

Anglis, spiked Carex.

Habitat in aquosis. 4. V. VI.

12. CAREX spiculis subovatis sessilibus remotis andro-
gynis capsulis acutis divergentibus spinosis. *Fl. suec.*
839. *Sp. pl.* 974.

Gramen cyperoides spicatum minimum spica divulsa
aculeata. *R. Syn.* 424.

Gramen nemorosum spicis parvis asperis. *Bauh. pin.*
7. th. 100: *Hist. ox. III.* 344. f. 8. t. 12. f. 26.

Gramen cyperoides echinatum minimum. *Park.* 1272:
Anglis,

Anglis, prickly Carex.

Habitat in palustribus et sylvis humidis. 2. VI.

canescens. 13. CAREX spiculis subrotundis remotis sessilibus obtusis androgynis capsulis ovatis obtusiusculis. *Fl. suec.* 842. *Sp. pl.* 974.

Gramen cyperoides spicatum minus, spica longa divulsa seu interrupta. *R. Syn.* 424.

Gramen cyperoides spicis curtis divulsis. *Fl. pr.* 117. t. 32.

Anglis, grey Carex.

Habitat ad sepes et in sylvis humidis. 2. V.—VIII.

remota. 14. CAREX spicis ovatis subsessilibus remotis androgynis foliis caulinis culmum æquantibus. *Cent. pl.* 92. *Syst. nat.* 1262.

Gramen cyperoides angustifolium, spicis parvis sessilibus infoliorum alis. *R. Syn.* 424. *Hist. ox. III.* 243. f. 8. t. 12. f. 17. *Pluk. ph. t.* 34. f. 3.

Anglis, long-leaved Carex.

Habitat in aggeribus fossarum et sylvis humidis. 2. VIII.

*** Spicis sexu distinctis : femineis sessilibus.

flava. 15. CAREX spicis confertis subsessilibus subrotundis mascula lineari, capsulis acutis recurvis. *Fl. suec.* 843. *Sp. pl.* 975.

Gramen cyperoides aculeatum germanicum vel minus. *Baub. pin.* 7. *Hist. ox. III.* p. 243. f. 8. t. 12. f. 19.

Gramen palustre echinatum. *Baub. hist. II.* 497. *Ger. Em.* 17. *R. Syn.* 421.

Gramen aculeatum Germanicum. *Park.* 1187.

8 Gramen cyperoides echinatum majus. *Pet. conc. gr.* 169.

Anglis, yellow Carex.

Habitat in palustribus et humidioribus pratis. 2. VI.

pilulifera. 16. CAREX spicis terminalibus confertis subrotundis masculina oblonga. *Roy. lugdb.* 75. *Sp. pl.* 976.

Gramen

Gramen cyperoides tenuifolium spicis ad summum
caulem sessilibus globulorum emulis. *Pluk. ph. t.*
9. f. 8. *R. Syn.* 422.

β Gramen cyperoides spicis brevibus congestis folio molli. *R. Syn.* 421.

Anglis, round-headed Carex.

Habitat in pratis et pascuis humidis passim. 4. VI. VII.

17. CAREX spicis tribus ovatis sessilibus alternis ; *saxatilis*.
mascula oblonga. *Fl. suec.* 848. *Sp. pl.* 976.

Gramen cyperoides vernum minimum. *R. Syn.* 421.

Gramen cyperoides spicatum. *Ger. Em.* 22.

Gramen caryophyllatæ foliis, spica divulsa. *Bauh. pin.*
3. th. 46.

Gramen spicatum foliis Caryophylleis. *Park.* 1160.

Anglis, vernal Carex.

Habitat in pratis montosis. 4. IV. V.

**** Spicis sexu distinctis ; femineis pedunculatis.

18. CAREX spicis pedunculatis erectis remotis : femi- *panicea*.
neis linearibus, capsulis obtusiusculis inflatis. *Fl.*
suec. 853. *Sp. pl.* 977.

Gramen cyperoides foliis caryophylleis, spicis e rario-
ribus et tumidioribus granis compositis. *R. Syn.* 418.

Pluk. ph. t. 91. f. 7.

Anglis, Pink Carex.

Habitat in paludibus turfosis. 4. VI. VII.

19. CAREX spicis erectis cylindricis ternis subsessili- *cæspitosa*.
bus : mascula terminali, culmo triquetro. *Fl. suec.*
853. *Sp. pl.* 978.

Gramen cyperoides foliis caryophylleis, spicis erectis
sessilibus, e feminibus confertis compositis. *R. Syn.*
418.

Gramen cyperoides angustifolium, spica spadiceo-vi-
ridi minus. *Bauh. pin.* 6. pr. 13.

Anglis, turf Carex.

Habitat in pratis, pascuis et sylvis humidis. 4. V. VI.

pallefcens. 20. CAREX spicis pendulis : mafcula erecta, femineis ovatis imbricatis; capfulis confertis obtufis. *Fl. fuc.* 852. *Sp. pl.* 977.

Carex spicis tribus petiolatis erectis remotis, mafcula et femineis. *Fl. lapp.* 327.

Gramen cyperoides polyftachyon flavicans, spicis brevibus prope fummitatem caulis. *R. Syn.* 419. *Hift. ox. III. f. 8. t. 12. f. 16.* *Pluk. ph. t. 34. f. 5.* *Anglis*, pale Carex.

Habitat in pratis et pascuis humidis paffim. 24. VI.

diffans. 21. CAREX spicis remotiffimis fubpedunculatis, bractea vaginali, capfulis angulatis mucronatis. *Syft. nat.* 1262.

Gramen cyperoides palufre, spicis tribus fubrotundis vix aculeatis, fpatio distantibus. *Hift. ox. III. t. 12. f. 18.*

Gramen cyperoides spicis parvis longiffime distantibus. *R. Syn.* 420.

Anglis, loofe Carex.

Habitat in udis turfosis frequens. 24. VI.

Pseudo-cyperus. 22. CAREX spicis pendulis, pedunculis geminatis, capfulis biaristatis. *Sp. pl.* 978. *Fl. fuc.* 854.

Gramen cyperoides spica pendula brevior. *Bauh. pin. 6. th* 86. *Hift. Ox. III. 242. f. 8. t. 12. f. 5.* *R. Syn.* 419.

Cyperus f. Pseudo-cyperus spica pendula brevior. *Park.* 1266.

Pseudo-cyperus. *Ger. Em.* 29.

Anglis, bastard Carex.

Habitat ad ripas feffarum frequens. 24. VII.

pendula. 23. CAREX spicis pendulis, mafcula erecta, femineis cylindricis pedunculo longioribus, capfulis acumina-
tis.

Gramen cyperoides spica pendula longior. *Park.* 1267. *R. Syn.* 420.

Gramen spica pendula longior et anguftior. *Bauh. pin. 6. pr.* 13. *hifl. II.* 497.

β Gramen

Gramen cyperoides polytachyon majusculum latifolium, spicis multis, longis, strigosis. *R. Syn.* 419.
Anglis, pendulous Carex.

Habitat in sylvis & sepibus humidis; in sepibus inter Hampstead et Highgate copiose. 4. VI. VII.

22. CAREX spicis pendulis, mascula erecta, femineis *sylvatica* filiformibus pedunculo brevioribus, capsulis acutis distantibus.

Gramen cyperoides sylvarum tenuius spicatum. *Park.*

1171. *Hist. Ox. III.* 243. f. 8. t. 12. f. 9.

Anglis, Wood Carex.

Habitat in sylvis humidis. 4. V. VI.

***** *Spicis sexudistinctis: masculis pluribus.*

23. CAREX spicis masculis pluribus, femineis subseffi- *acuta*. libus, capsulis obtusiusculis. *Fl. suec.* 857. *Sp. pl.* 978?

Gramen cyperoides. *Ger. Em.* 12. latifolium spica rufa f. caule triangulo. *Bauh. pin.* 6.

Gramen cyperoides cum paniculis nigris. *Bauh. hist. II.* 494. *R. Syn.* 417.

Gramen cyperoides majus latifolium. *Park.* 1265.

β Gramen cyperoides majus angustifolium. *Park.* 1265.

γ Gramen cyperoides minus angustifolium. *Park.* 1266.

Anglis, brown Carex.

Habitat in aquosis et ad rivulos. 4. V.

24. CAREX spicis masculis pluribus, femineis pedunculatis, capsulis inflatis acuminatis. *Fl. suec.* 856. *Sp. pl.* 979.

Gramen cyperoides majus præcox, spicis turgidis teretibus flavescentibus. *Hist. Ox. III.* 242. t. 12. f. 6. *R. Syn.* 420.

Gramen cyperoides vesicarium majus. *Cat. Giff.* 63.

β Gramen cyperoides polytachyon majus, spicis teretibus erectis. *R. syn.* 419.

Anglis, greater Bladder Carex.

Habitat in palustribus et aquosis passim. 6. VI.

MONOECIA TRIANDRIA.

- inflata.* 25. CAREX spicis masculis pluribus, femineis sessilibus capsulis ovatis inflatis acuminatis, bracteis duplo longioribus.
Gramen cyperoides palustre, spicis tribus subrotundis vix aculeatis, spatio distantibus. *Hist. Ox. III. t. 12. f. 18. R. Syn. 420.*
Gramen cyperoides vesicarium minus. *Cat. giff. 63. Anglis, lesser Bladder Carex.*
Habitat in palustribus & aquosis in comitatibus Eboracensi & Westmorlandico *passim. 4. VI. VII.*
- hirta.* 26. CAREX spicis remotis, masculis pluribus, femineis subpedunculatis erectis; capsulis hirtis. *Fl. suec. 858.*
Carex spicis remotis, femineis subsessilibus capsulis hirsutis. *Fl. lapp. 335. Sp. pl. 975.*
Gramen cyperoides polystachyon lanuginosum. *R. Syn. 418. Hist. Ox. III. 243. f. 8. t. 12. f. 10.*
Gramen cyperoides Norvegicum parium lanosum. *Park. 1172.*
Anglis, hairy Carex.
Habitat in pratis & pascuis humidis frequens. *4. VI.*

TETRANDRIA.

BETULA.

- Masc. *Cal. monophyllus, subtrifidus, triflorus. Cor. quadripartita.*
Fem. *Cal. monophyllus subtrifidus, biflorus. Strobilus cylindricus. Sem. utrinque membrana alata.*
- alba.* 1. BETULA foliis ovatis acuminatis ferratis. *Hort. cliff. 442. Fl. suec. 859. Sp. pl. 982.*
Betula. Bauh. hist. 1. 148. pin 427. R. Syn. Ger. Em. 1478.
Anglis, the Birch Tree.
Habitat in sylvis & sepibus humidis. *h. VII.*

BUXUS. 934.

- Masc. *Cal. tryphyllus. Petala duo. Germinis rudimentum.*
Fem. *Cal. tetraphyllus. Petala tria. Styli tres. Caps. trirostris, trilocularis. Sem. duo.*

1. BUXUS.

1. BUXUS. *Baub. hist.* 1. 496. *Ger. Em.* 1410. *R. Syn.* *sempervirens*.
 445. *Hort. cliff.* 441. *Sp. pl.* 983.
Buxus arborescens. Baub. pin. 471.
Buxus arbor vulgaris. Park. 1428.
Buxus angustifolia. R. Syn. 445.
Anglis, Box-Tree.
Habitat in sylvis & sepibus sed rarius. 4 IV.
Pharmac. Buxi folia lignum.

U R T I C A. 935.

Masc. Cal. tetraphyllus. Cor. nulla. Nectarium centrale cyathiforme.
Fem. Cal. bivalvis. Cor. o. Sem. unicum nitidum.

1. URTICA foliis oppositis ovatis serratis, amentis pilulifera.
fructiferis globosis. Hort. cliff. 440. *ups.* 282.
Roy. lugdb. 209. *Sp. pl.* 983.
Urtica urens pilulas ferens. Baub. pin. 232.
Urtica romana. Ger. Em. 784. *Park.* 440.
Urtica pilulifera folio profundius urticæ majoris in modum ferrato, semine magno lini. R. Syn. 140.
Anglis, Roman Nettle.
Habitat in ruderatis; plentifully at Great Yarmouth in Norfolk; Aldborough in Suffolk; R. Syn. ☉ VIII.
Pharmac. Urticæ romanæ semina.

2. URTICA foliis oppositis ovalibus. *Sp. pl.* 984. *urens.*
Fl. suec. 862.
Urtica minor. Ger. Em. 704. *R. Syn.* 140.
Urtica urens minor. Baub. pin. 232.
Anglis, lesser Nettle.
Habitat in cultis, ruderatis. ☉ VIII.

3. URTICA foliis oppositis cordatis, racemis geminis, dioica.
Sp. pl. 984. *Fl. suec.* 863. *The Balckirian, a Tartarian nation make cloth of this.*
Urtica urens. Ger. Em. 404. *maxima. Baub. pin.* 232.
Urtica racemifera major perennis. R. Syn. 139.
Anglis, common Nettle.
Habitat in ruderatis & ad sepes. 4. VII.
Pharmac. Urticæ semina, folia, cymæ.

P E N T A N D R I A.

X A N T H I U M. 937.

Masc. *Cal.* communis imbricatus. *Cor.* monopetalæ, quinquefidæ, infundibuliformes. *Receptaculum* paleaceum.

Fem. *Cal.* involucrum biphyllum, biflorum. *Cor.* o. *Drupa* sicca muricata, bifida. *Nucleus* bilocularis.

strumari- 1. XANTHIUM caule inermi. *Sp. pl.* 987. *Fl. suec.* 864.

Xanthium seu Lappa minor. *Baub. hist.* III. 572. *Park.* 1222.

Lappa minor Xanthium Dioscoridis. *Baub. pin.* 198. *Bardana* minor. *Ger. Em.* 800.

Anglis, lesser Burdock.

Habitat in fimetis sed rarius. ☉. VIII. IX.

Pharmac. Strumaræ Herba.

A M A R A N T U S. 941.

Masc. *Cal.* triphyllus. *Cor.* o.

Fem. *Cal.* triphyllus. *Cor.* o. *Styli* tres. *Caps.* unilocularis circumscissa. *Sem.* unicum.

Blitum. 1. AMARANTUS spicis triandris erectiusculis, foliis ovatis retusis. *Syst. nat.* 1268.

Amarantus floribus glomeratis lateralibus, foliis ovatis retusis, caule diffuso. *Sp. pl.* 990. *Fl. suec.* 865.

Amaranthus sylvestris et vulgaris. *Tourn. paris.* 385.

Blitum rubrum minus. *Baub. pin.* 118. *Cam. epit.* 235. *R. Syn.* 157.

β Blitum album minus. *Cam. epit.* 236.

Anglis, the least Blite.

Habitat in ruderalis. ☉. VIII.

P O L Y A N D R I A.

CERATOPHYLLUM. 944.

Masc. *Cal.* multipartitus. *Cor.* o. *Stam.* sedecim ad viginti.

Fem. *Cal.* multipartitus. *Cor.* o. *Pistil.* unicum. *Stylus* o. *Sem.* unicum, nudum.

1. CERATOPHYLLUM. *Fl. suec.* 866. *Sp. pl.* 992. *demersum.*

Hydroceratophyllon folio aspero, quatuor cornibus armato. *Vaill. aët. gal.* 1719. p. 16. t. 2. f. 1. *R. Syn.* 135.

Millefolium aquaticum equisetifolium, seu Equisetum ramosum aquis immersum. *Cat. cant. syn. ed. II.* 280.

β Hydroceratophyllon folio lævi, octo cornibus armato.

Vaill. aët. gal. 1719. p. 16. t. 2. f. 2.

Anglis, horned Pondweed.

Habitat in aquis pigrioribus et fossis ubique. 4. VII.

MYRIOPHYLLUM. 945.

Masc. *Cal.* tetraphyllus. *Cor.* o. *Stam.* octo.

Fem. *Cal.* tetraphyllus. *Cor.* o. *Pist.* quatuor.

Stylus o. *Sem.* quatuor nuda.

1. MYRIOPHYLLUM floribus masculis interrupte spicatum.

spicatis. *Fl. suec.* 867. *Sp. pl.* 992.

Myriophyllum vulgare majus. *Vaill. aët. gal.* 1719. p. 30.

Potamogiton foliis pennatis. *Tourn.* 233. *R. Syn.* 150.

Millefolium aquaticum pennatum spicatum. *Bauh. pin.* 141. pr. 73. *Park.* 1257.

β Millefolium aquaticum pennatum minus, foliolis singularibus latiusculis flosculis subjectis donatum.

Hist. ox. III. 622. f. 15. t. 4. f. 7.

Anglis, spiked Water Millfoil.

Habitat in fossis, lacubus et aquis quietis frequens. 4. VI. VII.

2. MYRIOPHYLLUM floribus omnibus verticillatis. *verticilla-*

Fl. suec. 868. *Sp. pl.* 992.

tum.

*Myriophyllum aquaticum minus. Clus. hist. II. 252.
Ger. Em. I. 616.*

*Myriophyllum aquaticum minus. Vaill. act. gal.
1719. p. 31.*

Pentapterophyllum aquaticum flosculis ad foliorum nodos: Cat. giff. 112. R. Syn. 316.

Millefolium aquaticum minus. Park. 1257. Bauh. hist. II. 783.

Millefolium aquaticum flosculis ad foliorum nodos. Bauh. pin. 141.

Anglis, verticillated Water Millfoil.

Habitat in fossis et aquis stagnantibus passim. 4. VII.

SAGITTARIA. 946.

Masc. Cal. triphyllus. Cor. tripetala. Filamenta fere viginti quatuor.

Fem. Cal. triphyllus. Cor. tripetala. Pist. nulla. Sem. multa, nuda.

sagittifolia. 1. *SAGITTARIA foliis sagittatis acutis. Fl. lapp. 344. succ. 869. Sp. pl. 993.*

Sagitta. Bauh. hist. III. 789. R. Syn. 258. minor latifolia Park. 1247.

Sagitta aquatica minor latifolia. Bauh. pin. 194.

2. *Sagitta aquatica omnium minima. D. Pluk. R. Syn. 258.*

Anglis, Arrow-head.

Habitat in fossis et ad ripas fluviorum. 4. VI.

POTERIUM. 948.

Masc. Cal. tetraphyllus. Cor. quadripartita. Stam. a triginta ad quinquaginta.

Fem. Cal. tetraphyllus. Cor. quadripartita. Pist. duo. Bacc. e tubo corollæ indurato.

sanguiforba. 1. *POTERIUM caulibus subangulosis. Hort. cliff. 446. upf. 288. Sp. pl. 994.*

Poterium inerme, filamentis longissimis. Roy. lugdb. 129. Dalib. parif. 291.

Sanguiforba minor. Bauh. hist. III. 113. R. Syn. 203.

Pimpinella sanguiforba minor hirsuta. Bauh. pin. 160.

Pimpinella vulgaris minor. Park. 582.

Pim-

Pimpinella sylvestris. Ger. Em. 1045.

β *Sanguisorba major* flore spadiceo. *Bauh. hist. III.* 120.
Anglis, Burnet.

Habitat in pratis montosis præsertim solo cretaceo, β in
pratis humidis. 2. VII.

Pharmac. *Pimpinellæ* f. *Sanguisorbæ* Herba.

Q U E R C U S. 949.

Masc. *Cal.* quinquefidus feræ. *Cor.* o. *Stam.* a
quinque ad decem.

Fem. *Cal.* monophyllus, integerrimus, scaber. *Cor.*
o. *Stylus* unus. *Sem.* unicum, ovatum.

1. QUERCUS foliis deciduis oblongis superne latioribus : sinubus acutioribus ; angulis obtusis. *Fl. suec.*
870. *Sp. pl.* 996. *Hall. herb.* 152. *Roy. lugdb.* 80.

Quercus vulgaris. Ger. Em. 1339.

Quercus latifolia. Park. 1385. *R. Syn.* 440.

Quercus cum longo pediculo. *Bauh. pin.* 420.

β *Quercus latifolia* mas quæ brevi pediculo est. *Bauh.*
pin. 419.

Anglis, the common Oak.

Habitat in sylvis et sepibus. 2. IV.

Pharmac. *Quercus Cortex, Folia, Glandes, Cupulæ.*

F A G U S. 951.

Masc. *Cal.* quinquefidus, campanulatus. *Cor.* o.
Stam. duodecim.

Fem. *Cal.* quadridentatus. *Cor.* o. *Styli* tres. *Capf.*
muricata quadrivalvis. *Sem.* duo.

*1. FAGUS foliis lanceolatis acuminato-ferratis subtus nudis. *Hort. cliff.* 447. *ups.* 287. *Roy. lugdb.* 79.
Sp. pl. 997.

Castanea. *Bauh. hist. I.* 121. *Ger. Em.* 1442. *R.*
Syn. 440. *vulgaris.* *Park.* 1400.

Castanea sylvestris. *Bauh. pin.* 419.

β *Castanea sativa.* *Bauh. pin.* 419.

Anglis, Chesnut Tree.

Habitat in sylvis et sepibus in Cantio frequens. 2. V.

Pharmac. *Castanæ Fructus.*

- sylvatica*. 2. FAGUS foliis ovatis obsolete ferratis. *Fl. suec.* 871.
Sp. pl. 998.
 Fagus. *Ger. Em.* 1444. *Bauh. pin.* 419. *Park.* 1403.
R. Syn. 439.
Anglis, Beach Tree.
 Habitat in sylvis et sepibus in Cantio, Suffexia, South-
 amptonia vulgarissima. *R. Syn.* h. V.

C A R P I N U S. 952.

- Masc. *Cal.* monophyllus, squama ciliata. *Cor.* o.
Stam. viginti.
 Fem. *Cal.* monophyllus, squama ciliata. *Cor.* o.
Germina duo. *Styli* singulis duo. *Nux* ovata.
Betulus. 1. CARPINUS squamis strobilorum planis. *Hort. cliff.*
 447. *Sp. pl.* 998.
 Carpinus. *Dod. pempt.* 841.
 Ostrea ulmo similis, fructu in umbilicis foliaceis.
Bauh. pin. 427. *R. Syn.* 451.
 Ostrea five Ostrea. *Park.* 1406.
 Betulus five Carpinus. *Ger. Em.* 1479.
Anglis, the Horn or Hard-beam Tree, Horse or
 Horn-beech Tree.
 Habitat in sylvis. h. V.

C O R Y L U S. 953.

- Masc. *Cal.* monophyllus, trifidus, squamæ formis
 uniflorus. *Cor.* o. *Stam.* octo.
 Fem. *Cal.* diphyllus lacerus. *Cor.* o. *Styli* duo.
Nux ovata.
Avellana. 1. CORYLUS stipulis ovatis obtusis. *Hort. cliff.* 448.
Fl. suec. 873. *Mat. med.* 431. *Sp. pl.* 998.
 Corylus sylvestris. *Ger. Em.* 1479. *Bauh. pin.* 418.
R. Syn. 439.
 Corylus f. Nux Avellana sylvestris. *Park.* 1416.
Anglis, the Hasel-nut Tree.
 Habitat in sylvis et sepibus. h. III.
Pharmac. Coryli f. Avellanæ Fructus.

MONADELPHIA.

PINUS. 956.

Masc. *Cal.* tetraphyllus. *Cor.* o. *Stam.* plurima.
Antheræ nudæ.

Fem. *Cal.* strobili: squama biflora. *Cor.* o. *Pist.*
 unum. *Nux* Ala membranacea excepta.

*1. PINUS foliis geminis: primordialibus solitariis glau- *sylvestris*
 bris. *Fl. suec.* 874. *Sp. pl.* 1000.

Pinus sylvestris. *Bauh. pin.* 491.

Pinus sylvestris foliis brevibus glaucis, conis parvis
 albertibus. *R. Syn.* 442.

Anglis, Scotch Firr.

Habitat in alpibus Scotiis. h. V.

2. PINUS foliis solitariis emarginatis. *Sp. pl.* 1001. *Picea.*

Abies. *Ger. Em.* 1363. *Park.* 1539.

Abies conis sursum spectantibus s. mas. *Bauh. pin.*
 505.

Anglis, the Yew-leaved Firr.

Habitat in alpibus Scotiis?

Pharmac. Abietis Summitates, Resina.

3. PINUS foliis subulatis mucronatis lævibus bifariam *Abies.*
 versus. *Sp. pl.* 1002. *Fl. suec.* 875.

Abies mas Theophrasti. *R. Syn.* 441.

Picea. *Park.* 1538. major. *Ger. Em.* 1354.

Picea major prima, sive *Abies* rubra. *Bauh. pin.* 493.

Anglis, common Firr, or Pitch Tree.

Habitat in alpibus Scotiis?

Pharmac. Abietis Resina.

Classis

Classis XXII.

D I O E C I A
D I A N D R I A.

S A L I X. 976.

Masc. *Amenti* squamæ. *Cor.* o. *Glandula* baseos nectarifera.

Fem. *Amenti* squamæ. *Cor.* o. *Stylus* bifidus.
Capf. unilocularis, bivalvis. *Sem.* papposa.

* *Foliis* glabris *ferratis*.

*herma-
phroditica*

1. SALIX foliis ferratis glabris, floribus hermaphroditis diandris. *Sp. pl.* 1015. *Fl. suec.* 878.

Salix latifolia folio splendente. *R. Syn.* 450.

Anglis, shining Willow.

Habitat in *sepibus* humidis. *About* Afton in Cumberland. *R. Syn.* h. III.

pentandra. 2. SALIX foliis ferratis glabris, flosculis pentandris.
Hort. cliff. 454. *Fl. suec.* 879. *Sp. pl.* 1016.

Salix folio laureo, seu lato glabro odorato. *Ph. br.*

R. Syn. 449.

Anglis, sweet Willow.

Habitat in *montosis* in *comitatibus* Westmorlandico, Cumberlandico, Eboracensi, *frequens.* h. IV.

vitellina. *3. SALIX foliis ferratis ovatis acutis glabris: ferraturis cartilagineis, petiolis calloso-punctatis. *Sp. pl.* 1016.

Salix fativa lutea, folio coenato. *Baub. pin.* 473.

Anglis, yellow Willow.

Habitat in *Salicetis* *passim.* h. IV. V.

4. SALIX foliis ferratis glabris lanceolatis petiolatis, fl- amygdali-
pulis trapeziformibus. *Roy. lugdb.* 83. *Fl. suec. na.*
881.
Salix perficæ folio auriculato. *Hall. helv.* 151.
Salix folio auriculato splendente flexilis. *Cat. cantab.*
R. Syn. 448.
Salix folio Amygdalino utrinque virente aurito. *Bauh.*
pin. 473.
Salix viminalis nigra. *Park.* 1431.
Anglis, Almond-leaved Willow.
Habitat in Salicetis et ad ripas fluviorum. *h.* V.
5. SALIX foliis ferratis glabris ovato-lanceolatis, petio- fragilis.
lis dentato-glandulosis. *Fl. lapp.* 359. *t.* 8. *f.* 13.
suec. 883. *Sp. pl.* 1017.
Salix folio longo latoque splendente fragilis. *R. Syn.*
448.
Salix fragilis. *Bauh. pin.* 474. *Park.* 1431.
Salix spontanea fragilis amygdalino folio auriculata
et non auriculata. *Bauh. hist.* I. 214.
β Salix folio Amygdalino utrinque aurito corticem abji-
ciens. *R. Syn.* 448.
Anglis, Crack Willow.
Habitat in aquosis. *h.* V.
- 6 SALIX foliis subferratis glabris lanceolatis : inferio- purpurea.
ribus oppositis. *Sp. pl.* 1017.
Salix folio longo subluteo non auriculata viminibus
rubris. *R. Syn.* 450.
Anglis, Purple Willow.
Habitat in salicetis sed rarius, ad ripam Thamefis in-
ter Vauxhall et Nine-Elms, prope Londinum. *h.*
IV. V.
7. SALIX foliis ferratis glabris lanceolato-linearibus : Helix.
superioribus oppositis obliquis. *Roy. lugdb.* 83. *Da-*
lib. paris. 299. *Sp. pl.* 1017.
Salix foliis utrinque lævibus : inferne glaucis, squamis
et capsulis tomentosis. *Hall. helv.* 153.
Anglis, Rose Willow.
Habitat in aquosis sed rarius. *h.* V.

herbacea. 8. *SALIX* foliis ferratis glabris orbiculatis. *Fl. lapp.* 355. *t.* 8. *f.* *N. et t.* 7. *f.* 3, 4. *Fl. suec.* 887. *Sp. pl.* 1018.

Salix alpina alni rotundo folio repens. *Bocc. mus. P.* II. *E. I. R. Syn.* 448.

Salix saxatilis minima. *Bauh. pin.* 474.

Anglis, herbaceous Willow.

Habitat in *ascensu montis Snowdon copiose*, D. G. Sherard. *R. Syn. et in montibus Westmorlandicis.* 4. VI.

**** Foliis glabris integerrimis.**

reticulata. 9. *SALIX* foliis integerrimis glabris ovatis obtusis. *Fl. suec.* 888. *Sp. pl.* 1018.

Salix foliis integris ovatis subtus reticulatis. *Fl. lapp.* 359. *t.* 8. *f.* *L. et t.* 7. *f.* 1, 2.

Salix pumila folio rotundo. *Bauh. hist. I.* 12. *Sch. it. alp.* 43. 340. *R. Syn.* 449.

Anglis, round-leaved Willow.

Habitat in *montibus Cambrobritanicis, Eboracensibus, Westmorlandicis passim.* 4. VI.

rubra. 10. *SALIX* foliis integris, glabris lineari-lanceolatis acutis.

Salix minime fragilis, foliis longissimis, utrinque viridibus, non ferratis, D. Sherard. *R. Syn.* 449.

Anglis, red Willow.

Habitat in *Salicetis sed rarius. In the Osterholt between Maidenhead and Windsor; and near Salisbury*, by Mr. Sherard and Mr. Rand. *R. Syn.* 4. IV. V.

Foliis integerrimis, villosis.

arenaria. 11. *SALIX* foliis integris ovatis acutis supra subvillosis, subtus tomentosis. *Fl. suec.* 894. *Sp. pl.* 1019.

Salix foliis integris subtus villosis ovatis acutis. *Hall. helv.* 155. *t.* 5. *f.* 2.

Salix pumila foliis utrinque candicantibus et lanuginosis. *R. Syn.* 447. *t.* 19. *f.* 3.

Anglis, Sand Willow.

Habitat

Habitat in pascuis arenosis. Observed by Sandown-
castle by Mr. J. Sherard. *R. Syn.* 4. VI.

12. SALIX foliis integerrimis lanceolatis utrinque sub-
pilosis caule repente. *Fl. suec.* 896. *Sp. pl.* 1020.

Salix pumila angustifolia inferne lanuginosa. Bauh. hist.

I. 212. *R. Syn.* 447.

Salix pumila brevi angustoque folio incano. Bauh. pin.
474.

Salix humilis angustifolia repens. Park. 1434.

- β *Salix alpina pumila rotundifolia repens, inferne subci-
nerea. Bauh. pin.* 474. *R. Syn.* 448.

Anglis, creeping Willow.

Habitat in paludibus turfosis. 4. V.

13. SALIX foliis integerrimis ovatis subtus villosis niti-
dis. *Fl. lapp.* 364. t. 8. f. 13. *suec.* 897. *Sp. pl.*
1020.

Salix pumila angustifolia prona parte cinerea. Bauh.
hist. 213. *R. Syn.* 447.

Salix pumila linifolia incana. Bauh. pin. 474.

Salix pumila angustifolia recta. Park. 1434.

Chamæitea five Salix pumila. Ger. Em. 1391.

Anglis, brown Willow.

Habitat in ericetis humidis. 4. V. VI.

14. SALIX foliis integerrimis lanceolato linearibus *rosmarini*
strictis sessilibus subtus tomentosis. *Roy. lugdb.* 84. *folio.*
Fl. suec. 898.

Salix pumila Rhamni secundi Clusii folio. R. Syn.
447.

Salix oblongo incano acuto folio. Bauh. pin. 159.

Salix humilis repens angustifolia, Lob. ic. P. 2. 137.

Bauh. hist. I. 214.

Anglis, Rosemary-leaved Willow.

Habitat in montosis udis. 4. V.

15. SALIX foliis integerrimis utrinque villosis obovatis *aurita.*
appendiculatis. *Fl. lapp.* 369. t. 8. f. 17. *suec.* 891.
Sp. pl. 1019.

Salix folio rotundo minore. Dill. cat. giff. 38. *R. Syn.*
450.

β *Salix*

- β *Salix caprea* pumila folio subrotundo, subtus incano.
R. Syn. 450.
Anglis, round-leaved Willow.
Habitat in sylvis et sepibus. Ⅵ. V.

**** *Foliis subserratis, villosis.*

- caprea.* 16. *SALIX* foliis ovatis rugosis subtus tomentosis undatis superne denticulatis. *Roy. lugdb.* 83. *Fl. suec.* 900. *Sp. pl.* 1020.
Salix folio rugoso obiter ferrato subtus lanuginoso, julo crassissimo. *Hall. helv.* 155.
Salix caprea rotundifolia. *Ger. Em.* 1319.
Salix latifolia rotunda. *Bauh. pin.* 474. *Park.* 1432.
R. Syn. 449.
 β *Salix* folio ex rotunditate acuminato. *Bauh. pin.* 474.
Anglis, common Sallow.
Habitat in sylvis et sepibus. Ⅵ. IV. V.
- viminalis.* 17. *SALIX* foliis subintegerrimis lanceolato linearibus longissimis acutis subtus sericeis, ramis virgatis. *Fl. suec.* 901. *Sp. pl.* 1021.
Salix folio longissimo. *Cat. Cant.*
Salix angustis et longissimis foliis crispis subtus albicantibus. *Bauh. hist. I.* 212.
Anglis, the Osier.
Habitat in salicetis. Ⅵ. IV.
- alba.* 18. *SALIX* foliis lanceolatis acuminatis ferratis utrinque pubescentibus: ferraturis infimis glandulosis. *Fl. suec.* 903. *Sp. pl.* 1021.
Salix. *Ger. Em.* 1389. *R. Syn.* 447. arborea angustifolia, alba vulgaris. *Park.* 1430.
Salix vulgaris alba arborescens. *Bauh. pin.* 473.
Anglis, common or white Willow.
Habitat in sylvis et sepibus. Ⅵ. IV.
Pharmac. Salicis Folia, Cortex.

TRIANDRIA.

EMPETRUM. 977.

Masc. *Cal.* tripartitus. *Cor.* tripetala. *Stam.* longa.
Fem. *Cal.* tripartitus. *Cor.* tripetala. *Styli* novem.
Sem. 9-sperma.

1. EMPETRUM procumbens. *Fl. suec.* 904. *Sp. pl.* *nigrum*.

1022.

Empetrum montanum, fructu nigro. *Tourn.* 579. *R.*

Syn. 444.

Erica baccifera procumbens nigra. *Bauh. pin.* 486.

Erica baccifera procumbens. *Ger. Em.* 1383.

Erica baccifera nigra. *Park.* 1485.

Anglis, black-berried Heath, Crow, or Crake-berries.

Habitat in montibus udis Derbienfibus, Eboracenfib.
bus. 4. IV. V.

TETRANDRIA.

VISCUM. 979.

Masc. *Cal.* quadripartitus. *Cor.* o. *Filamenta* o.
Antheræ calyci adnatæ.

Fem. *Cal.* tetraphyllus, superus. *Stylus* o. *Cor.* o.
Bacca monosperma.

1. VISCUM foliis lanceolatis obtusis, caule dichotomo, *album*.

spicis axillaribus. *Fl. suec.* 905. *Sp. pl.* 1023.

Viscum. *Ger. Em.* 1315. *R. Syn.* 464. *vulgare.* *Park.*
1392.

Viscum baccis albis. *Bauh. pin.* 423.

Anglis, white Mistle, or Mistletoe.

Habitat in arboribus, frutex parasiticus. 2. V.

Pharmac. *Visci Folia.*

HIP-

HIPPOPHAE. 980.

Masc. *Cal. bipartitus. Cor. o.*

Fem. *Cal. bifidus. Cor. o. Stylus unicus. Bacca monosperma.*

Rhamnoides. 1. HIPPOPHAE foliis lanceolatis. *Fl. suec. 906. Sp. pl. 1023.*

Rhamnoides fructifera, foliis falcatis, baccis leviter flavescens. Tourn. Cor. 53. R. Syn. 445.

Rhamnus. 2 Clus. hist. 110. Ger. Em. 1334.

Rhamnus falcatis folio angusto, fructu flavescens.

Bauh. pin. 477.

Rhamnus primus Dioscoridis Lobelio, five littoralis.

Park. 1006.

Anglis, Sallow-thorn, or Sea Buck-thorn.

Habitat in littoribus maritimis. h. IV.

MYRICA. 981.

Masc. *Amenti squama lunata. Cor. o.*

Fem. *Amenti squama lunata. Cor. o. Styli duo.*

Bacca monosperma.

Gale. 1. MYRICA foliis lanceolatis subserratis, caule suffruticoso. *Fl. suec. 907. Sp. pl. 1024.*

Gale frutex odoratus septentrionalium, Elæagnus Cordi. Bauh. hist. I. 223. R. Syn. 443.

Rhus myrtifolia Belgica. Bauh. pin. 414.

Rhus sylvestris five Myrtus Brabantica vel Anglica.

Park. 1451.

Myrtus Brabantica f. Elæagnus Cordi. Ger. Em. 1414.

Anglis, Goule, Sweet Willow, Dutch Myrtle. Sweet G.

Habitat in paludibus sed rarius. h. V. On

Bagshot Heath near the XXI mile stone, plentifully.

P E N-

PENTANDRIA.

HUMULUS. 989.

Masc. *Cal.* pentaphyllus. *Cor.* o.

Fem. *Cal.* monophyllus, oblique patens, integer.

Cor. o. *Styli* duo. *Sem.* unicum, intra calycem planum.

1. HUMULUS. *Sp. pl.* 1028. *Fl. suec.* 908.

Lupulus mas et fœmina. *Bauh. hist.* II. 151. *pin.*

298. *R. Syn.* 137.

Lupulus sylvestris. *Park.* 176.

Lupus salictarius. *Ger. Em.* 885.

Anglis, Hops.

Habitat in sepibus. 2. VI.

Pharmac. *Lupuli strobili.*

HEXANDRIA.

TAMUS. 991.

Masc. *Cal.* sexpartitus. *Cor.* o.

Fem. *Cal.* sexpartitus. *Cor.* o. *Stylus* trifidus.

Bacca trilocularis, infera. *Sem.* duo.

1. TAMUS foliis cordatis indivisis. *Sp. pl.* 1028.

communis.

Tamnus racemosa, flore minore luteo pallescente.

Tourn. 103. *R. Syn.* 262.

Bryonia nigra. *Ger. Em.* 869.

Bryonia sylvestris nigra. *Park.* 178.

Bryonia lævis f. *nigra racemosa.* *Bauh. pin.* 297.

Anglis, black Briony.

Habitat ad sepes et in dumetis. 2. VI.

Pharmac. *Bryoniæ nigræ Folia, Radix.*

O C T A N D R I A.

P O P U L U S. 996.

Masc. *Amenti Cal.* lamina lacera. *Cor.* turbinata, obliqua, integra.

Fem. *Amenti Cal. et Cor.* maris. *Stigma* quadrifidum. *Capsf.* bilocularis. *Sem.* multa, papposa.

alba.

1. POPULUS foliis subrotundis dentato-angulatis subtus tomentosis. *Fl. suec.* 911. *Sp. pl.* 1034.

Populus alba. *Ger. Em.* 1486. *Park.* 1400. *R. Syn.* 446. majoribus foliis. *Bauh. pin.* 429.

Anglis, white Poplar.

Habitat in sepibus et ad rivulos. *h.* III.

nigra.

2. POPULUS foliis deltoidibus acuminatis serratis. *Sp. pl.* 1034. *Fl. suec.* 910.

Populus nigra. *Ger. Em.* 1485. *Bauh. pin.* 429. *Park.* 1410. *R. Syn.* 446.

Anglis, black Poplar.

Habitat in umbrosis humidis et ad fluvios. *h.* III.

Pharmac. Populi nigri. *Folia*, *Germinæ*.

tremula.

3. POPULUS foliis subrotundis dentato-angulatis utrinque glabris. *Sp. pl.* 1034. *Fl. suec.* 909.

Populus lybica. *Ger. Em.* 1487. *Park.* 1411. *R. Syn.* 446.

Populus tremula. *Bauh. pin.* 429.

Anglis, trembling Poplar, or Asp.

Habitat in sylvis humidis. *h.* III.

R H O D I O L A. 997.

Masc. *Cal.* quadripartitus. *Cor.* tetrapetala.

Fem. *Cal.* quadripartitus. *Cor.* o. *Nectaria* quatuor. *Pist.* quatuor. *Capsf.* quatuor Polyspermae.

Rosea.

1. RHODIOLA. *Fl. lapp.* 378. *suec.* 912. *Sp. pl.* 1035.

Anacampseros radice rosam spirante major. *Tourn.* 264. *R. Syn.* 262.

Rhodia

DIOECIA ENNEANDRIA.

371

Rhodia radix. Bauh. pin. 286. hist. III. 683. Ger.

Em. 532. Park. 729.

Anglis, Rosewort.

Habitat in montibus Westmorlandicis, Cumberlandicis Eboracensibus et Cambrobritannicis frequens.

4. VI. VII.

ENNEANDRIA.

MERCURIALIS. 998.

Masc. *Cal. tripartitus. Cor. o. Stam. a novem ad duodecem. Antheræ globosæ didymæ.*

Fem. *Cal. tripartitus. Cor. o. Styli duo. Caps. dicocca, bilocularis monosperma.*

1. MERCURIALIS caule simplicissimo, foliis scabris. *perennis,*

Fl. suec. 913. Sp. pl. 1035.

Mercurialis perennis repens Cynocrambe dicta. R. Syn. 138.

Mercurialis montana spicata et Mercurialis montana spicata. Bauh. pin. 122.

Mercurialis sylvestris Cynocrambe dicta vulgaris, mas et foemina. Park. 295.

Cynocrambe mas et foemina. Ger. Em. 333.

Anglis, Dog's Mercury.

Habitat in sylvis et sepibus. 4. IV. V.

2. MERCURIALIS caule brachiato, foliis glabris. *Sp. annua, pl. 1035.*

Mercurialis annua glabra vulgaris. R. Syn. 139.

Mercurialis testiculata f. mas. Bauh. pin. 121.

Mercurialis spicata f. foemina. Bauh. pin. 121.

Mercurialis mas foemina. Ger. Em. 333.

Mercurialis vulgaris mas et foemina. Park. 295.

Anglis, French Mercury.

Habitat in ruderalis, cultis, circa Londinum frequens.

©. IX.

HYDROCHARIS. 999.

Masc. *Spatha diphylla. Cal. trifidus. Cor. tripetala. Filamenta tria interiora stylifera.*

B b 2

Fem.

DIOECIA MONADELPHIA.

Fem. *Cal.* trifidus. *Cor.* tripetala. *Styli* sex.
Capf. sexlocularis polysperma, infera.

- Morfus ranæ.* 1. HYDROCHARIS. *Fl. suec.* 914. *Sp. pl.* 1036.
Nymphæa alba minima. *Bauh. pin.* 193. f. *Morfus ranæ.* *Park.* 1252.
Morfus ranæ. *Ger. Em.* 818.
Stratiotes foliis aloes, femine rotundo. *R. Syn.* 290.
Anglis, Frog-bit.
Habitat in fossis et aquis leniter fluentibus. 4. VI.

MONADELPHIA.

JUNIPERUS. 1005.

Masc. *Amenti* Calyx squamæ. *Cor.* o. *Stam.* tria.
 Fem. *Cal.* tripartitus. *Petala* tres. *Styli* tres.
Bacca trisperma, tribus tuberculis calycis inæqualis.

- communis.* 1. JUNIPERUS foliis ternis patentibus mucronatis, bacca longioribus. *Syst. nat.* 1297. *Fl. suec.* 915. *Sp. pl.* 1040.
Juniperus vulgaris baccis parvis purpureis. *Bauh. hist. I.* 293. *R. Syn.* 444.
Juniperus. *Ger. Em.* 1372. *vulgaris.* *Park.* 1028. *fruticosa.* *Bauh. pin.* 488.
 6 *Juniperus Alpina.* *Bauh. hist. I.* 301.
Anglis, common Juniper.
Habitat in ericetis et montosis. 2. V.
Pharmac. *Juniperi Baccæ, Lignum, Resinâ.*

TAXUS. 1006.

Masc. *Cal.* triphyllus gemmæ. *Cor.* o. *Stamina* multa. *Antheræ* peltatæ octofidæ.
 Fem. *Cal.* triphyllus gemmæ. *Cor.* o. *Stylus* o.
Sem. unicum calyculo baccato, integerrimo.

- baccata.* 1. TAXUS foliis approximatis. *Fl. suec.* 916. *Sp. pl.* 1040.
Taxus. *Ger. Em.* 1370. *Park.* 1412. *Bauh. hist. I.* 241. *pin.* 505. *R. Syn.* 445.
Anglis, Yew Tree.

Habitat

Habitat in montibus Westmorlandicis, Cumberlandicis, Herefordiensibus frequens. \mathfrak{h} . III. IV.

SYNGENESIA.

R U S C U S. 1008.

Masc. *Cal.* hexaphyllus. *Cor.* o. *Nectarium* centrale, ovatum, apice perforatum.

Fem. *Calyx*, *Corolla* et *Nectarium* Maris. *Stylus* unus. *Bacca* trilocularis. *Sem.* duo.

I. RUSCUS foliis supra floriferis nudis. *Sp. pl.* 1041. *aculeatus*.
Dalib. parif. 304.

Ruscus. *Bauh. pin.* 470. *hist. I.* 579. *Park.* 253.

R. Syn. 262.

Ruscus five Bruscus. *Ger. Em.* 907.

Anglis, Knee Holly, or Butcher's Broom.

Habitat in sylvis, dumetis, et ericetis frequens. \mathfrak{h} .
III. IV.

Pharmac. Rusci Radix.

BRYONIA.

Masc. *Cal.* quinquedentatus. *Cor.* quinquepartita.
Filamenta tria.

Fem. *Cal.* dentatus. *Cor.* quinquepartita. *Stylus* trifidus. *Bacca* subglobosa, polysperma.

I. BRYONIA foliis palmatis utrinque calloso-scabris. *alba*.
Fl. suec. 876. *Sp. pl.* 1012.

Bryonia alba. *Ger. Em.* 869. *R. Syn.* 261. *vulgaris*. *Park.* 178.

Bryonia aspera s. alba baccis rubris. *Bauh. pin.* 297.
Anglis, white Briony.

Habitat in sepibus et dumetis. \mathfrak{u} . V.

Pharmac. Bryoniæ Radix.

Classis XXIII.

POLYGAMIA
MONOECIA.

H O L C U S. 1015.

Hermaphroditus. *Calyx*. Gluma biflora. *Cor*. Gluma aristata. *Stamina* tria. *Styli* duo. *Sem.* unicum.

Masc. *Cal.* Gluma bivalvis. *Cor.* o. *Stam.* tria.

lanatus.

1. HOLCUS glumis bifloris villosis : hermaphrodito mutico masculo arista recurva. *Fl. suec.* 917. *Sp. pl.* 1048.

Aira foliis pubescentibus, panicula contracta : flosculo hermaphrodito mutico, masculo arista uncinata calyce brevior. *It. oel.* 122. *Fl. suec. ed. I.* 67.

Gramen pratense paniculatum molle. *Baub. pin.* 2. pr. 3. th. 37. *Sch. agr.* 234. *Park.* 1155.

Gramen miliaceum paniculatum molle. *Pet. conc. gr.* 124. *R. Syn.* 404.

Anglis, Meadow Soft-grass.

Habitat in pratis et pascuis. 4. VI.

mollis.

2. HOLCUS glumis bifloris nudiusculis : hermaphrodito mutico masculo arista geniculata. *Syst. nat.* 1305.

Gramen miliaceum aristatum molle. *Pet. conc. gr.* 125. *R. Syn.* 404.

Gramen caninum longius radicum majus et minus. *Baub. pin.* 1.

Gramen paniculatum molle, radice graminis canini repente. *Hist. ox. III.* 202.

Anglis, creeping Soft-grass.

Habitat in pratis, pascuis et ad sepes. 4. VII.

ÆGILOPS

ÆGILOPS. 1018.

Hermaphrod. *Cal.* Gluma solitaria, bipartita. *Cor.*
Gluma calyci contraria. *Stamina* tria. *Styli* duo.
Sem. unicum.

Masc. *Cal.* et *Cor.* Gluma ut in priore. *Stam.*
tria.

1. ÆGILOPS spica subulata mutica incurva. *Sp. pl. incurva.*
1051.

Nardus spica cylindrica articulata. *Sp. pl.* 53.

Gramen parvum marinum spica loliacea. *Ger. Em.*
78. *R. Syn.* 395.

Gramen loliaceum maritimum spicis articulatis. *Hist.*
ox. III. 182. *f.* 8. *t.* 2. *f.* 8.

Gramen loliaceum minus spica simplici. *Bauh. pin.* 9:
pr. 19.

Phoenix acerosa aculeata. *Park.* 1145.

Anglis, Sea Hard-grass.

Habitat in littoribus maritimis. ☉. VII. VIII.

VALANTIA. 1019.

Hermaphrod. *Cal.* o. *Cor.* quadripartita. *Stam.*
quatuor. *Stylus* bifidus. *Sem.* unicum.

Masc. *Cal.* o. *Cor.* quadripartita. *Stam.* quatuor.
Pist. obsoletum.

1. VALANTIA floribus masculis quadrifidis, peduncu- *cruciata:*
lis diphyllis. *Hort. upf.* 303. *Sp. pl.* 1052.

Gallium foliis quaternis, flosculis in alis confertis.
Hall. helv. 462.

Cruciata. *Ger. Em.* 1123. *R. Syn.* 223.

Cruciata vulgaris. *Park.* 566.

Cruciata hirsuta. *Bauh. pin.* 335.

Anglis, Croswort, or Mugweed.

Habitat in dumetis et ad sepes. ♀. V. VI.

Pharmac. Cruciatæ Herba.

PARIETARIA. 1020.

Hermaphrod. *Cal.* quadrifidus. *Cor.* o. *Stam.* qua-
tuor. *Stylus* unicus. *Sem.* unicum, superum, e-
longatum.

B b 4

Fem.

Fem. *Cal.* quadrifidus. *Cor.* o. *Stylus* unicus. *Sem.* unicum, superum, elongatum.

officinalis. 1. **PARIETARIA** foliis lanceolato-ovatis, pedunculis dichotomis calycibus diphyllis. *Syst. nat.* 1308. *Sp. pl.* 1052.

Parietaria. *Ger. Em.* 331. *Bauh. hist. II.* 976. *R. Syn.* 158.

Parietaria vulgaris. *Park.* 437.

Parietaria officinarum et Dioscoridis. *Bauh. pin.* 121. *Anglis*, Pellitory of the Wall.

Habitat in muris et ruderatis. 4. V. IX.

Pharmac. *Parietariæ Herba.*

A T R I P L E X. 1021.

Hermaphrod. *Cal.* pentaphyllus. *Cor.* o. *Stam.* quinque. *Stylus* bipartitus. *Sem.* unicum, depressum.

Fem. *Cal.* diphyllus. *Cor.* o. *Stylus* bipartitus. *Sem.* unicum compressum.

portulacoides. 1. **ATRIPLEX** caule fruticoso, foliis obovatis. *Fl. suec.* 919. *Sp. pl.* 1035.

Atriplex maritima fruticosa, *Halimus et Portulaca marina dicta*, angustifolia. *R. Syn.* 153.

Halimus seu portulaca marina. *Bauh. pin.* 120.

Halimus vulgaris f. Portulaca marina. *Ger. Em.* 323.

Portulaca marina nostras. *Park.* 724.

Anglis, Sea Purslane.

Habitat in littoribus maritimis. 5. VIII.

laciniata. 2. **ATRIPLEX** caule herbaceo, foliis deltoideo-dentatis subtus argenteis. *Sp. pl.* 1053. *Fl. suec.* 920.

Atriplex maritima. *Bauh. hist. II.* 974. *R. Syn.* 152.

Atriplex maritima laciniata. *Bauh. pin.* 120.

Atriplex marina. *Ger. Em.* 326. *repens.* *Park.* 758.

Anglis, jagged Sea Orache.

Habitat in littoribus maritimis. 6. VIII.

erecta. 3. **ATRIPLEX** caule herbaceo, foliis hastatis dentatis. *Atriplex angustifolia laciniata.* *R. Syn.* 152.

Anglis, Spear-leaved Orache.

Habitat

Habitat in ruderatis. On the entrance into Battersea meadow from Nine-Elms, Mr. Martin. R. Syn. \odot . VIII.

4. ATRIPLEX caule herbaceo, valvulis femineis magnis deltoideis sinuatis. *Fl. suec.* 921. *Sp. pl.* 1053. Atriplex sylvestris, folio hastato seu deltoide. R. Syn. 151.

Atriplex sylvestris annua, folio deltoide triangulari sinuato et mucronato, hastæ cuspidi simili. *Hist. ox.* II. 607. f. 5. t. 32. f. 14.

- β Atriplex maritima perennis folio deltoide seu triangulari minus incano. *Hist. ox.* II. 607. 19.

- γ Atriplex maritima nostras procerior, foliis angulosis incanis, admodum sinuatis. *Pluk. alm.* 60.

- δ Atriplex maritima ad foliorum basin auriculata procumbens et ne vix sinuata. *Pluk. alm.* 61.

Anglis, wild Orache.

Habitat in ruderatis, fimetis et littoribus maritimis. \odot . VIII. IX.

5. ATRIPLEX caule herbaceo patulo, foliis subdeltoideo-lanceolatis calycibus seminum disco dentatis. *Fl. suec.* 922. *Sp. pl.* 1053.

Atriplex caule herbaceo, foliis superioribus lanceolatis integerrimis. *Hort. cliff.* 469. *Roy. lugdb.* 218.

Atriplex sylvestris angustifolia. *Ger. Em.* 326. *Park.* 748. R. Syn. 151.

Atriplex angusto oblongo folio. *Bauh. pin.* 119.

Anglis, narrow-leaved Orache.

Habitat in ruderatis et ad sepes. \odot . VIII.

6. ATRIPLEX caule herbaceo erecto, foliis linearibus serratis.

Atriplex angustifolia maritima dentata. R. Syn. 152.

Atriplex maritima angustifolia. *Bauh. pin.* 152.

Atriplex angustifolia dentata. *Pet. h. br. t.* 7. f. 4.

Anglis, indented Sea Orache.

Habitat in ruderatis et littoribus maritimis. \odot . VIII.

7. ATRIPLEX caule herbaceo erecto, foliis omnibus linearibus line-

linearibus integerrimis. *Roy. lugdb.* 218. *Fl. suec.* 923. *Sp. pl.* 1054.

Atriplex angustifolium et longissimo folio: *H. L. Bat.* 79. *Tourn.* 505. *R. Syn.* 153.

Atriplex graminea maritima. *Pet. h. br. t.* 7. f. 6.

β *Atriplex maritima scopariæ folio.* *D. Dale.* *R. Syn.* 153.

♂ *Atriplex maritima angustifolia, obtusiore folio.* *R. Syn.* 153.

Anglis, Grass-leaved Orache.

Habitat in ruderalis, fimetis et littoribus maritimis.

○. VIII. IX.

*peduncu-
lata.*

8. **ATRIPLEX** caule herbaceo divaricato, foliis lanceolatis obtusis integris calycibus foemineis pedunculatis. *Lent. pl.* 101. *Syst. nat.* 1309.

Atriplex marina femine lato. *Ph. br.* *R. Syn.* 153.

Atriplex maritima Halimus dicta, humilis erecta, femine folliculis membranaceis bivalvibus in latitudinem expansis et utrinque recurvis, longo pediculo insidentibus, clauso. *Pluk. ph. t.* 36. f. 1.

β *Atriplex maritima nostras* *Ocymi minoris folio.* *D. Pluk.* *R. Syn.* 253.

Anglis, Stalked Sea Orache.

Habitat in littoribus maritimis; prope Scirbeck vicum milliare a Bostonia distantem, copiose provenit D. Plukenet. Found in the isle of Thanet, near the ferry-house, in great plenty, by Mr. J. Sherard. R. Syn. ○. VIII. IX. - Near Ramsgate on the beach.

A C E R. 1023.

Hermaphrod. *Cal.* quinquefidus. *Cor.* pentapetala. *Stam.* octo. *Pistil.* unicum. *Caps.* duæ f. tres monospermæ, ala terminatæ.

Masc. *Cal.* quinquefidus. *Cor.* pentapetala. *Stam.* octo.

*Pseudo
platanus.*

1. **ACER** foliis quinquelobis inæqualiter serratis floribus racemosis. *Sp. pl.* 1054.

Acer montanum candidum. *Bauh. pin.* 430.

Acer majus. *Dod. pempt.* 840. *Ger. Em.* 1484. *R. Syn.* 470.

Acer

Acer majus latifolium, *Sycomorus falso dictum*. *Park.*

1425.

Anglis, the greater Maple, or Sycomore-tree.

Habitat in *sepibus et ad domos*. *h. V. VI.*

2. *ACER* foliis lobatis obtusis emarginatis. *Sp. pl.* 1055. *campestre*.

Fl. suec. 925.

Acer minus. *Ger. Em.* 1484. *R. Syn.* 470.

Acer minus et vulgare. *Park.* 1426.

Acer campestre et minus. *Bauh. pin.* 431.

β *Acer campestre et minus*, fructu rubente. *Vaill. bot.*

parif. R. Syn. 470.

Anglis, common Maple.

Habitat in *sepibus et dumetis*. *h. V. VI.*

D I O E C I A.

F R A X I N U S. 1026.

Hermaphrod. *Cal.* quadripartitus. *Cor. o. Stam.*

duo. *Pist.* unicum. *Sem.* unicum lanceolatum.

Fem. *Pist.* unicum. *Sem.* unicum lanceolatum.

I. *FRAXINUS* foliis serratis, floribus apetalis. *Sp. pl. excelsior*.

1057. *Fl. suec.* 926.

Fraxinus. *Ger. Em.* 1472. *R. Syn.* 469.

Fraxinus excelsior. *Bauh. pin.* 416.

Anglis, common Ash-tree.

Habitat in *sylvis et sepibus*. *h. III. IV.*

Classis

*Classis XXIV.*CRYPTOGAMIA
F I L I C E S.

EQUISETUM. 1033.

Spica fructificationibus peltatis, basi dehiscentibus multivalvi.

sylvaticum.

1. EQUISETUM caule spicato, frondibus compositis.
Fl. suec. 927. Sp. pl. 1061.

*Equisetum sylvaticum. Tab. ic. 253. Ger. Em. 1114.
R. Syn. 130.*

Equisetum sylvaticum tenuissimis fetis. Baub. pin. 16.

Equisetum omnium minus tenuifolium. Park. 1201.

- β *Equisetum sylvaticum, procumbens, fetis uno versu dispositis. R. Syn. 131.*

- γ *Equisetum palustre tenuissimis et longissimis fetis.
Baub. pin. 16. pr. 24.*

Anglis, Wood Horse-tail.

Habitat in sylvis et umbrosis humidis. 4. IV. V.

arvense.

2. EQUISETUM scapo fructificante nudo; sterili frondoso. *Sp. pl. 1061. Fl. suec. 928.*

*Equisetum arvense longioribus fetis. Baub. pin. 16.
Park. 1202. R. Syn. 130.*

Equisetum segetale. Ger. Em. 1114.

- β *Equisetum pratense longissimis fetis. Baub. pin. 16.
Anglis, Corn Horse-tail.*

Habitat in arvis humidis. 4. III.

palustre.

3. EQUISETUM caule angulato, frondibus simplicibus. *Fl. suec. 929. Sp. pl. 1061.*

Equisetum palustre brevioribus fetis. Baub. pin. 16.

*Equisetum palustre. Lob. ic. 795. Ger. Em. 1114. R.
Syn. 131. Fl. lapp. 392. minus. Park. 1200.*

Equisetum

- β Equisetum palustre minus polystachion. *Bauh. pin.* 16.
pr. 24. *R. Syn.* 131. *t.* 5. *f.* 3.
Anglis, Marsh Horse-tail.
Habitat in aquosis. 4. VI.
4. EUISETUM caule striato, frondibus subsimplici-*fluuiatile.*
bus. Fl. suec. 930. *Sp. pl.* 1062.
 Equisetum caule non fulcato latissimo, verticillis densissimis. *Hall. helv.* 144.
 Equisetum majus. *Ger. Em.* 1115. *R. Syn.* 130. palustre. *Park.* 1200.
 Equisetum palustre longioribus foliis. *Bauh. pin.* 15.
Anglis, River Horse-tail.
Habitat in aquosis et ad ripas lacuum et fluviorum frequens. 4. V.
5. EUISETUM caule subnudo lævi. *Fl. suec.* 931. *limosum.*
Sp. pl. 1062.
 Equisetum nudum læuius nostras. *R. Syn.* 131. *t.* 5. *f.* 2.
Anglis, smooth Horse-tail.
Habitat in palustribus et aquosis. 4. VI.
6. EUISETUM caule nudo scabro basi subramoso. *byemale.*
Fl. suec. 932. *Sp. pl.* 1062.
 Equisetum nudum. *Ger. Em.* 1113. *R. Syn.* 131.
 Equisetum junceum ramosum. *Park.* 1207.
 Equisetum foliis nudum ramosum. *Bauh. pin.* 16.
 β Equisetum nudum minus variegatum Basiliense. *Bauh. pin.* 16. *pr.* 25. *hist. III.* 730.
 γ Equisetum foliis nudum non ramosum f. junceum.
Bauh. pin. 16.
 Equisetum junceum f. nudum. *Park.* 1201.
 δ Equisetum læve pæne nudum. *Pet. conc. gr.* 238.
Anglis, rough Horse-tail, or Shave-grass.
Habitat in palustribus et aquosis sed rarius. 4. VII.
 VIII.

O P H I O G L O S S U M: 1035.

Spica articulata, disticha: articulis transversim dehiscentibus.

1. OPHI-

- vulgatum* 1. OPHIOGLOSSUM fronde ovata. Fl. suec. 932. Sp. pl. 1062.
 Ophioglossum. Ger. Em. 404. Baub. hist. III. 708.
 R. Syn. 128. f. Lingua serpentina. Park. 506.
 Ophioglossum primum f. vulgatum. Baub. pin. 354.
 Anglis, Adder's Tongue.
 Habitat in pratis et pascuis humidis; in ericeto Hamp-
 stediano. 4. V.
 Pharmac. Ophioglossi herba.

O S M U N D A. 1036.

Spica ramosa: fructificationibus globosis.

* Scapis insidentibus cauli ad basin frondis.

- Lunaria. 1. OSMUNDA scapo caulino solitario, fronde pinnata solitaria. Fl. suec. 934. Sp. pl. 1064.
 Osmunda fronde pinnata caulina, pinnis lunatis. Hort. cliff. 472.
 Lunaria minor. Ger. Em. Park. 507. R. Syn. 128.
 Lunaria racemosa minor. Baub. pin. 354.
 β Lunaria minor foliis dissectis. R. Syn. 129.
 Anglis, Moon-wort.
 Habitat in pratis et pascuis montosis. In comitatibus Westmorlandico, Cumberlandico, Eboracensi, frequens. 4. V.

** Fronde ipsa fructificationes ferente.

- regalis. 2. OSMUNDA fronde bipinnata apice racemifera. Fl. suec. 933. Sp. pl. 1065.
 Osmunda regalis f. Filix florida. Park. 1038.
 Filix ramosa non dentata florida. Baub. pin. 357.
 Filix florida f. Osmunda regalis. Ger. Em. 1131.
 Anglis, flowering Fern, or Osmund Royal.
 Habitat in aqueosis et palustribus putridis passim. 4. VII. VIII.

*** Frondibus aliis foliaceis; aliis fructificantibus.

- Spicant. 3. OSMUNDA frondibus lanceolatis pinnatifidis: laciniiis conniventibus integerrimis parallelis. Fl. suec. 936. Sp. pl. 1066.

Lon-

Lonchitis aspera. *Ger. Em.* 1140. *R. Syn.* 118.

Lonchitis aspera minor. *Bauh. pin.* 359. *Park.* 1042.

Anglis, rough Spleen-wort.

Habitat in nemorosis et ericetis humidis. 4. VII.

4. OSMUNDA frondibus supradecompositis : pinnis al-crispa.
ternis subrotundis incisis. *Sp. pl.* 1067.

Adiantum album crispum alpinum. *Schwenckf. cat.* 10.
R. Syn. 126.

Adiantum album floridum, f. *Filicula petræa crispa*
perelegans. *R. syn. ed. II.* 51. *Pluk. ph. t.* 9. f. 3.
Anglis, Stone Fern.

Habitat in petris et rupibus. In comitatibus Eboracen-
si, et Westmorlandico, frequens. 4. VIII.

ACROSTICHUM. 1037.

Frustrificationes discum totum frondis tegentes.

* *Fronde simplici divisa.*

1. ACROSTICHUM fronde nuda lineari-laciniata. *septentri-*
Fl. lapp. 380. *suec.* 937. *Sp. pl.* 1068. *onale.*

Filix saxatilis corniculata. *Bauh. pin.* 358.

Filix saxatilis Tragi. *Bauh. hist. III.* 755. *Park.*
1045. *R. Syn.* 120.

Anglis, forked, or horned Fern.

Habitat in fissuris rupium. Ad cacumen montis Car-
nedh-Lhewelyn prope Lhan-Lhechyd in agro Arvo-
nienfi invenit, D. Lhwyd. In muris antiquis Lhan-
Dethylæ uno circiter milliari a Lhan-Rhoost aquilo-
nem versus, D. Richardson. *R. Syn. in montibus su-*
pra Ambulside, in comitatu Westmorlandico. 4.
VIII.

** *Fronde sub-bipinnata.*

2. ACROSTICHUM fronde sub-bipinnata : pinnis op-
posito-coadunatis obtusis subtus hirsutis basi inte-
gerrimis. *Fl. suec.* 938. *Sp. pl.* 1071.

Filix alpina, pedicularis rubræ foliis subtus villosis.
D. Lhwyd. *R. Syn.* 118.

Filix pumila, lonchitidis Maranthæ species Cambro-
bri-

britannica; an *Lonchitis aspera ilvensis*. *Pluk. ph. t. 89. f. 8.*

Anglis, Hairy Fern.

Habitat in *fissuris rupium*. It grows on a moist black rock almost at the top of Clogwyn y Garnedh, facing north-west, directly above the lower lake. D. Richardson. *R. Syn.* 24.

Thelypteris.

3. ACROSTICHUM fronde pinnata: pinnis pinnatifidis integerrimis. *Fl. suec.* 939. *Sp. pl.* 1071.

Filix minor palustris repens. *R. Syn.* 122.

Dryopteris. *Lob. ic.* 814. *Ger. Em.* 1135. f. *Filix querna repens.* *Park.* 1041.

Anglis, Marsh Fern.

Habitat in *palustribus cespitosis*. In Westmorlandia *passim.* 24. VIII.

PETERIS. 1038.

Fruetificationes in lineis marginalibus.

aquilina.

1. PETERIS fronde supradecomposita: foliolis pinnatis: pinnis lanceolatis: infimis pinnatifidis; superioribus minoribus. *Hort. cliff.* 473. *Fl. suec.* 940. *Sp. pl.* 1075.

Filix foemina. *Ger. Em.* 1128. *R. Syn.* 124. *vulgaris.* *Park.* 1037.

Filix ramosa major pinnulis non dentatis. *Bauh. pin.* 357.

β *Filicula saxatilis ramosa maritima nostras.* *R. Syn.* 125. *Pluk. ph. t.* 182. *f. 1.*

Anglis, Female Fern, or Brakes.

Habitat in *sylvis et ericetis ubique.* 24. VIII.

ASPLENIUM. 1042.

Fruetificationes in lineolis disci frondis, sparsis.

* *Fronde simplici.*

Scolopendrium.

1. ASPLENIUM frondibus simplicibus cordato-lingulatis integerrimis stipitibus hirsutis. *Sp. pl.* 1079.

Asplenium frondibus ensiformibus integris: basi cordatis inflexis, petiolis hirsutis. *Vir. cliff.* 103. *Hall. belv.* 134.

Phyllitis.

Phyllitis. *Ger. Em.* 1138. *R. Syn.* 116. f. *Lingua*

Cervina vulgaris. Park. 1046.

Lingua Cervina officinarum. Bauh. pin. 353.

Anglis, Hart's-tongue.

Habitat in umbrosis saxosis, passim. 4.

Pharmac. Phyllitidis Folia.

**** Fronde pinnatifida:**

2. **ASPLENIUM** frondibus pinnatifidis: lobis alternis *Ceterach*
confluentibus. *Hort. cliff.* 474. *Hall. helv.* 135.

Dalib. parif. 310.

Asplenium f. *Ceterach. Bauh. hist. III.* 749. *Ger. Em.*

1138. *Park.* 1046. *R. Syn.* 118.

Ceterach officinarum. Bauh. pin. 354.

Anglis, Spleenwort.

Habitat in muris antiquis et fissuris rupium humidis

passim. Circa Bristolium copiose. 4.

Pharmac. Ceterach f. *Asplenii Herba.*

***** Fronde pinnata.**

3. **ASPLENIUM** frondibus pinnatis; pinnis subrotundis *Tricho-*
dis crenatis. Fl. lapp. 388. *suec.* 942. *Sp. pl.* 1080. *manes.*

Trichomanes. Park. 1051. *R. Syn.* 119. f. *Poly-*

trichum officinarum. Bauh. pin. 356.

Trichomanes mas. Ger. Em. 1146.

Anglis, common Maidenhair.

Habitat in saxosis, muris antiquis, et umbrosis fre-
quens. 4.

Pharmac. Trichomanis Herba.

4. **ASPLENIUM** fronde pinnata: pinnis subrotundis *viridi.*
basi truncatis.

Trichomanes costa viridi. D. Lhwyd.

Trichomanes ramosum. Bauh. hist. III. 755. *R. Syn.*

119.

Trichomanes foemina. Ger. Em. 1146.

β *Trichomanes ramosum majus et minus. Bauh. pin.*
356.

Anglis, green Maidenhair.

Habitat in rupibus humidis in comitatibus Eboracensi et Westmorlandico, passim. 4.

- marinum.* 5 **ASPLENIUM** fronde pinnata : pinnis obovatis serratis superne gibbis obtusis basi cuneatis. *Hort. cliff.* 474. *Hall. helv.* 136. *Sp. pl.* 1081.
Filix marina anglica. Park. 1045.
Filicula petræa fœmina f. Chamæfilix marina anglica. Ger. Em. 1143.
Chamæfilix marina anglica. Baub. hist. III. 737. *R. Syn.* 119. *Hist. ox. III.* 575. *t. 3. f. 25.*
Anglis, Sea Maidenhair, or Fern.
Habitat in rupibus maritimis. On the rocks about Prestholm island, near Beaumaris, and at Llandwyn in the isle of Anglesea; about the castle of Hastings in Suffex; in rupibus maritimis in Cumberlandico, 4.

**** *Fronde decomposita.*

- Ruta mu-* 6. **ASPLENIUM** frondibus alternatim decompositis :
 varia. foliolis cuneiformibus crenulatis. *Roy. lugdb.* 498.
Fl. suec. 943. *Sp. pl.* 1081.
Ruta muraria. Baub. pin. 356. *hist. III.* 753. *Ger. Em.* 1144. *R. Syn.* 122. *f. Salvia vitæ. Park.* 1050.
Adiantum album. Tab. ic. 796.
 3 *Adiantum album tenuifolium, Rutæ murariæ accedens. Baub. hist. III.* 753 *R. Syn.* 123. *Pluk. ph. t. 3. f. 3*
Anglis, white Maidenhair.
Habitat in rimis petrarum. 4.
Pharmac. Adianti albi, Herba.
- Adiantum* 7. **ASPLENIUM** frondibus subtripinnatis : foliolis alternis ; pinnis lanceolatis inciso-serratis. *Sp. pl.* 1081.
Asplenium frondibus duplicato-pinnatis ; foliis inferioribus majoribus : foliolis ovatis superne crenatis. Roy lugdb. 498.
Adiantum nigrum officinarum. Baub. hist. III. 742. *R. Syn.* 126.

Adian-

Adiantum nigrum vulgare. Park. 1046.

Adiantum foliis longioribus, pulverulentis, pediculo nigro. Bauh. pin. 356.

Onopteris mas. Ger. Em. 1137.

β *Filix elegans, adianto nigro accedens, segmentis rotundioribus. R. Syn. 127.*

γ *Filix pumila petræa nostras, adianti nigri foliorum æmula, saxorum interveniis prorumpens. Pluk. alm. 91. R. Syn. 127.*

δ *Filix minor longifolia, tarsis raris, pinnulis longis, tenuissimis et oblongis laciniis fimbriatis. D. Sheppard. R. Syn. 127.*

Anglis, black Maidenhair.

Habitat in umbrosis et muris antiquis frequens. 4.

Pharmac. Adianti nigri, Herba.

POLYPODIUM. 1043.

Frustrificationes in punctis subrotundis, sparsis, per discum frondis

* *Fronde pinnatifida lobis coadunatis.*

1. POLYPODIUM frondibus pinnatifidis: pinnis oblongis subserratis, obtusis, radice squamata. *Fl. suec. 944. Sp. pl. 1085.*

Polypodium vulgare. Bauh. pin. 359. Park. 1039.

Polypodium. Ger. Em. 1138. R. Syn. 117.

β *Polypodium murale pinnulis serratis. Pet. bot. hort. 11. pl. 1. R. Syn. 117.*

Anglis, common Polypody.

Habitat in muris antiquis, umbrosis et ad radices arborum. 4.

Pharmac. Polypodii Herba.

2. POLYPODIUM frondibus pinnatis, pinnis lanceolatis lacero-pinnatifidis serratis. *Sp. pl. 1086. Cambri-*

Polypodium Cambrobritannicum, pinnulis ad margines laciniatis. R. Syn. 117.

Polypodium Cambrobritannicum lobis foliorum profunde dentatis. Hist. ox. III. 565. f. 14. t. 2. f. 8.

Filix amplissima, lobis foliorum laciniatis cambrica. Pluk. ph. t. 30. f. 1.

CRYPTOGAMIA FILICES.

Anglis, jagged Polypody.

Habitat on a rock in a wood near Dennys Powys Castle,
not far from Caerdiff in Glamorganshire. *R. Syn.*
4.

** *Fronde pinnata.*

Lonchitis. 3. POLYPODIUM frondibus pinnatis : pinnis lunulatis ciliato-ferratis declinatis, petiolis strigosis. *Sp. pl.* 1088. *Gron. virg.* 123.

Lonchitis aspera major. *Ger. Em.* 1140. *R. Syn.* 118.

Lonchitis aspera. *Bauh. pin.* 359.

Lonchitis aspera major Matthiolo. *Park.* 1042.

Anglis, great Polypody, or Spleenwort.

Habitat in fissuris rupium in Wallia, prope Bingley in agro Eboracensi. 4.

phlegopte- 4. POLYPODIUM frondibus subbipinnatis : foliolis infimis reflexis : paribus pinnula quadrangulari condunatis. *Fl. suec.* 945. *Sp. pl.* 1089.

Filix minor britannica, pediculo pallidiore, alis inferioribus deorsum spectantibus. *Hist. ox.* III. 575. *J. 14. t. 4. f. 17. R. Syn.* 122.

Anglis, Wood Polypody.

Habitat in rimis petrarum humidis et umbrosis in comitatibus Eboracensi et Westmorlandico passim. 4.

fragrans. 5. POLYPODIUM frondibus subbipinnatis lanceolatis : foliolis confertis : lobis obtusis ferratis, stipite paleaceo. *Sp. pl.* 1089.

Dryopteris rubrum idæum spirans. *Amm. euth.* 251.

Anglis, sweet Polypody.

Habitat in rimis petrarum humidis ; prope Kewick in comitatu Westmorlandico. 4.

fontanum. 6. POLYPODIUM fronde subpinnata lanceolata : foliolis subrotundis, stipite lævi. *Sp. pl.* 1089.

Polypodium pinnarum pinnis bidentatis et tridentatis brevissimis. *Hall. helv.* 140.

Filicula fontana minor. *Bauh. pin.* 358.

Filix

Filix saxatilis non ramosa minima. R. cat. eur. Pluk. ph. t. 89. f. 2.

- β *Filix saxatilis omnia minima elegantissima. Hort. reg. paris. Pluk ph. t. 89. f. 3.*

Anglis, Rock Polypody.

Habitat in fissuris rupium et muris antiquis. Supra Hammerham church, D. Bradney; in locis saxosis supra Wybourn in Westmorlandia. 4.

7. **POLYPODIUM** fronde bipinnata pinnulis obtusis *Filix mas.* crenulatis, stipite paleaceo. *Hort. cliff. 475. Sp. pl. 1090. Fl. suec. 946.*

Filix mas vulgaris. Park. 1036. R. Syn. 120.

Filix mas non ramosa dentata. Bauh. pin. 258.

Filix mas non ramosa, pinnulis latis densis minutim dentatis Ger. Em. 1129.

Anglis, Male Polypody, or Fern.

Habitat in sylvis, ericetis, locis saxosis. 4.

Pharmac. Filicis maris Radix.

8. **POLYPODIUM** fronde bipinnata: pinnulis lanceo- *F. femina* latis pinnatifidis acutis. *Fl. lapp. 386. suec. 947. Sp. pl. 1090.*

Filix mas non ramosa, pinnulis angustis raris profunde dentatis. Ger. Em. 1130. Pluk. ph. t. 130. f. 4. Hist. ox. III. 579. f. 14. t. 3. f. 8. R. Syn. 121.

Anglis, Female Polypody.

Habitat in palustribus et umbrosis frequens. 4.

9. **POLYPODIUM** fronde bipinnata: pinnis lanceo- *aculeatum.* lato-ovatis dentato-ciliatis, stipite strigoso.

Polypodium fronde bipinnata: pinnis lanceolatis dentatis, stipite strigoso. Rcy lugdb. 500. Sp. pl. 1090.

Filix mas aculeata major. Bauh pin. 358. pr. 151.

Filix mas non ramosa, pinnulis latis auriculatis spinosis. Ger. Em. 1130. Pluk. ph. t. 179. f. 6. R. Syn. 121.

- β *Filix aculeata major, pinnulis auriculatis crebrioribus, foliis integris angustioribus. R. Syn. 121.*

- γ *Filix mas aculeata nostras, alis expansis muscosa lanugine aspersa. Pluk. ph. t. 180. f. 1.*

CRYPTOGAMIA FILICES.

Anglis, prickly Polypody.
Habitat in umbrosis. 4.

lobatum. 10. POLYPODIUM fronde bipinnata : pinnis lanceolato-ovatis dentato-ciliatis superne basi lobatis, stipite strigoso.

Filix lonchitidis affinis. *R. Syn.* 121. *Pluk. ph. t.* 180. f. 3.

Anglis, lobed Polypody.
Habitat in umbrosis. 4.

gristatum. 11. POLYPODIUM frondibus subbipinnatis : foliolis ovato-oblongis ; pinnis obtusiusculis apice acute ferratis. *Sp. pl.* 1090.

Filix mas ramosa, pinnulis dentatis. *Ger. Em.* 1129. *Pluk. ph.* 181. f. 2. *R. Syn.* 124.

β Filix montana ramosa minor argute denticulata. *D. Lhwyd. R. Syn.* 124.

Anglis, crested Polypody.
Habitat in sylvis et umbrosis humidis. In ericeto Hampstediano. 4.

rhæticum. 12. POLYPODIUM fronde bipinnata ; foliolis pinnisque remotis lanceolatis ferraturis acuminatis. *Sp. pl.* 1091.

Filix rhætica tenuissime denticulata. *Bauh. hist.* III. 740.

Filix fontana major f. Adiantum album filicis folio. *Bauh. pin.* 351.

Filix pumila saxatilis altera. *Clusii* 1043. *R. Syn.* 122.

Anglis, Stone Polypody.
Habitat in montibus saxosis in Westmorlandia passim. 4.

fragile. 13. POLYPODIUM fronde bipinnata : foliolis remotis pinnis subrotundis incis. *Fl. lapp.* 384. *Succ.* 948. *Sp. pl.* 1091.

Filix saxatilis caule tenui fragile. *R. Syn.* 125. *Pluk. ph. t.* 180. f. 5.

β Adiantum nigrum pinnulis cicutarix divisura. *D. Bo- bart. R. Syn.* 126.

Anglis,

Anglis, brittle Polypody.
Habitat in aridis lapidosis. 4.

*** *Fronde supradecomposita.*

14. POLYPODIUM fronde supradecomposita : foliolis *Dryopteris* bipinnatis. *Fl. succ.* 949. *Sp. pl.* 1093. *ris.*

Filix saxatilis ramosa nigris maculis punctata. *Bauh. pin.* 358.

Filix ramosa minor. *Bauh. hist.* III. 741. *R. Syn.* 125.

Filix pumila saxatilis prima Clusii. *Park.* 1043.

Dryopteris Tragi. *Ger. Em.* 1135.

Anglis, branched Polypody.

Habitat in aridis saxosis in comitatibus Eboracensi, Lancastriensi, Westmorlandico et Cumberlandico frequens. 4.

A D I A N T U M. 1044.

Fructificationes in maculis terminalibus, sub replicato margine frondis.

* *Fronde decomposita.*

1. ADIANTUM frondibus decompositis : foliolis alternis : pinnis cuneiformibus lobatis pedicellatis. *Sp. veneris. pl.* 1096.

Adiantum foliis coriandri. *Bauh. pin.* 355.

Adiantum verum, f. *Capillus veneris* verus. *Park.* 1049.

Capillus veneris verus. *Ger. Em.* 1144. *R. Syn.* 123.

Anglis, true Maidenhair.

Habitat in rupibus. Found by Mr. Lhwyd at Barry island, and Porth Kirig in Glamorganshire. *R. Syn.* 4.

Pharmac. *Adianti veri Herba.*

** *Fronde supra decomposita.*

2. ADIANTUM frondibus supradecompositis : foliolis trapeziformibus : pinnis rhombeis incisis utrinque fructificationibus. *Sp. pl.* 1097.

Adiantum majus coriandri folio, adianto vero affine, pediculo pallide rubente. *Sibb. pr.* 7. *R. Syn.* 124.

CRYPTOGAMIA FILICES.

Adiantum vero affinis minor Scotica, folio obtuso, faturate viridi, &c. *Sibb. pr.* 8. *R. Syn.* 124.

Adiantum ramosum pediculis lucidis et nigris. *Plum. fil. amer.* 78. *t.* 95.

Anglis, shining Maidenhair.

Habitat in *Scotia*. *R. Syn.*

TRICHOMANES. 1045.

Frustrificationes solitariae, stylo setaceo terminatae, margini ipsi frondis insertae.

pyxidiferum. 1. TRICHOMANES frondibus subbipinnatis: pinnis alternis lobatis linearibus. *Sp. pl.* 1098.

Filix pyxidifera. *Plum. fil.* 74. *t.* 50.

Filix humilis repens, foliis pellucidis et splendentibus caule alato. *D. Richardson. R. Syn.* 127. *t.* 3. *f.* 3, 4.

Dareæ tunbrigensis facie. *Pet. fil.* 104. *t.* 13. *f.* 13. *Anglis*, Cup Trichomanes.

Habitat in *petris et rupibus aqua humentibus*. Found by *Dr. Richardson at Belbank, scarce half a mile from Bingley, at the head of a remarkable spring.* *R. Syn.* 4.

tunbrigense. 2. TRICHOMANES frondibus pinnatis: pinnis lobatis oblongis crenulatis. *Sp. pl.* 1098.

Adiantum petræum perpuffilum anglicum, foliis bifidis trifidisque. *R. Syn.* 123.

Adiantum radicosum humis parsum, f. Filicula pellucida nostras, coriandri foliis mollicellis, globuliferum. *Pluk. ph. t.* 3. *f.* 5, 6.

Darea tunbrigensis minor. *Pet. mus.* 761.

Anglis, Tunbridge Trichomanes.

Habitat in *fissuris rupium humidis*. Found upon the *moist rocks in Wales, and near Settle, by Dr. Richardson; near Tunbridge, by Mr. Dare; at Cockbush, six miles from Chichester, by Mr. Rand, and Mr. Sherard. R. Syn. in montibus Westmorlandicis passim.* 4. VIII.

PILULARIA. 1047.

Flores *Masculi* ad latus frondis.

Fructificationes *Femineæ* ad radicem, globosa quadri-
locularis.

1. PILULARIA. *Fl. suec.* 950. *Sp. pl.* 1100. *globulifera*

Pilularia palustris juncifolia. *Vaill. parif.* 159. *A.* 15. *ra.*

f. 6. *Dill. musc.* 538. *t.* 79. *f.* 1.

Graminifolia palustris repens, vasculis granorum pi-
peris æmulis. *R. Syn.* 136.

Gramen piperinum. *Merr. pin.*

Anglis, Pepper-grass.

Habitat in inundatis præsertim solo arenoso, prope fon-
tes medicatos *Strethamenses*, Dr. Wilmer; in ericeto
Hounsflemano, prope *Whitton*. *R. Syn. in ericeto*
Hampstediano. 4. VI. IX.

ISOETES. 1048.

Flores *Masculi* *Anthera* intra basin frondis.

Flores *Feminei* *Capsula* bilocularis, intra basin frondis.

1. ISOETIS. *It. scan.* 417. *t.* 419. *Fl. suec.* 951. *Sp. lacustris.*
pl. 1100.

Calamaria folio longiore et graciliore. *Dill. musc.* 541.
t. 80. *f.* 2.

Subularia vulgaris erecta, folio rigidissimo. *R. Syn.*
306.

β *Subularia repens* folio minus rigido. *R. Syn.* 306.

δ *Subularia fragilis*, folio longiore et tenuiore. *R. Syn.*
307.

Anglis, Quillwort.

Habitat in fundo lacus exigui prope summitatem montis
Snowdon Phynon-vrech dicti. *R. Syn.*

MUSCI.

LYCOPODIUM. 1049.

Anthera bivalvis, sessilis; *Calyptra* nulla.

1. LYCOPODIUM foliis sparsis filamentosis, spicis clavatum.
teretibus pedunculatis geminis. *Fl. suec.* 952. *Sp.*
pl. 1101. Ly-

CRYPTOGAMIA MUSCI.

Lycopodium vulgare pilosum anfragosum et repens.
Dill. musc. 441. t. 58. f. 1.

Lycopodium. Tabern. ic. 814. R. Syn. 107.

Muscus terrestris clavatus. Baub pin. 360.

Muscus clavatus s. Lycopodium. Ger. Em. 1562.
Park. 1307.

Anglis, common Club-moss.

Habitat in ericetis et montibus Westmorlandicis, Cum-
berlandicis, Eboracensibus, &c. copiose ut et in erice-
to Hounsflemano. 4. VII.

Pharmac. Musci clavati Herba Lycopodii semina.

inunda-
tum.

2. **LYCOPODIUM** foliis sparsis integerrimis, spicis
 terminalibus foliosis. *Sp. pl. 1102. Fl. succ. 954.*

Lycopodium palustre repens, Clava singulari. Vail.
parif. 123. t. 16. f. 11. Dill. musc. 452. t. 62.
f. 7.

Muscus terrestris repens, clavis singularibus foliosis e-
rectis. Cat. pl. ang. R. Syn. 108.

Anglis, Marsh Club-moss.

Habitat in ericetis humidis passim. 4. VII.

annoti-
num.

3. **LYCOPODIUM** foliis sparsis quinquefariis subser-
 ratis furculis annotino-articulatis, spicis terminali-
 bus glabris erectis. *Sp. pl. 1103. Fl. succ. 956.*

Lycopodium elatius, juniperinum, clavis singularibus
sine pediculis. Dill. musc. 455. t. 63. f. 9. R. Syn.
107.

Muscus clavatus, juniperipis foliis reflexis, clavis sin-
gularibus sine pediculis. Hist. ox. III. 624. f. 15.
t. 5. f. 3.

Anglis, Welsh Club-moss.

Habitat in Alpibus Arvonice sed rarius; copiosum obser-
vavimus in monte Rhiwr Glyder supra lacum Lhyn
y own, prope ecclesiam S. Perisii, nec rarius in de-
pressis ejusdem montis y Glyder, qua rupem y Tri-
van spectat. D. Lhwyd. R. Syn. 4. VII.

alpinum.

4. **LYCOPODIUM** foliis quadrifariam imbricatis acu-
 tis, furculis erectis dichotomis, spicis sessilibus te-
 retibus. *Sp. pl. 1104. Fl. succ. 957.*

Ly-

- Lycopodium sabinae* facie. *Dill. musc.* 445. t. 58.
f. 2. R. Syn. 108.
Muscus clavatus cupressiformis. *Park.* 1309.
Muscus clavatus foliis Cupressi. *Bauh. pin.* 360.
Ger. Em. 1562.
Anglis, Mountain Club-moss.
Habitat in montibus; in comitatibus Eboracensi, Cum-
berlandico, Westmorlandico, et in montibus Wal-
licis. 4. VIII. IX.
5. *LYCOPODIUM* foliis sparsis octifariis, furculo di- *Selago.*
 chotomo erecto fastigiato, floribus sparsis. *Sp. pl.*
 1102. *Fl. suec.* 955.
Selago vulgaris abietis rubrae facie. *Dill. musc.* 435.
 t. 56. *f. 1.*
Selago foliis et facie abietis. *Fl. jen.* 330. *R. Syn.*
 106.
Chamaepeuce. *Turn. part.* 1. p. 129.
Muscus erectus ramosus saturate viridis. *Bauh. pin.*
 360.
Anglis, Firr Club-moss.
Habitat in ericetis montosis. On Snowdon and other
high Mountains in Wales; in comitatibus Eboracensi,
Westmorlandico, Cumberlandico, Lancastriensi,
frequens. 4. VIII.
6. *LYCOPODIUM* foliis sparsis ciliatis lanceolatis *Selaginoi-*
 spicis solitariis terminalibus foliosis. *Sp. pl.* 1101. *des.*
Fl. suec. 953.
Selaginoides foliis spinosis. *R. Syn.* 106. *Dill. musc.*
 460. t. 88. *f. 1.*
Anglis, Prickly Club-moss.
Habitat in locis turfosis, in comitatibus Eboracensi,
Lancastriensi, Westmorlandico, Cumberlandico,
frequens. 4. VIII.

S P H A G N U M. 1051.

Anthera operculata: ore laevi Calyptra nulla.

1. *SPHAGNUM* ramis deflexis. *Fl. lapp.* 415. *suec. palustre.*
 959. *Sp. pl.* 1106.
Sphagnum palustre molle deflexum, squamis cymbi-
formibus. *Dill. musc.* 240. t. 32. *f. 1.*

Sphag-

Sphagnum cauliferum et ramosum palustre molle candidans, reflexis ramulis, foliolis latioribus. Cat. giff. 229. R. Syn. 104.

Muscus palustris albicans terrestri similis, capitulis erectis brevibus, pediculis etiam brevissimis insidentibus. R. Syn. ed. II. 37.

β *Sphagnum palustre molle deflexum, squamis capillaceis. Dill. musc. 243. t. 32. f. 2*

Sphagnum cauliferum et ramosum palustre molle candidans, reflexis ramulis, foliolis angustioribus. Cat. giff. 229. R. Syn. 104.

Anglis, Grey Bog-moss.

Habitat in paludibus turfosis. 4. VII. VIII.

alpinum. 2. *SPHAGNUM subramosum erectum. Sp. pl. 1106.*

Sphagnum subulatum viridissimum, capsulis ovatis. Dill. musc. 245. t. 32. f. 2.

Anglis, Green Bog-moss

Habitat in paludibus turfosis in montibus Cader Idæris et Snowdon copiose. Dill. IX. X.

arborescens. 3. *SPHAGNUM ramosum repens, antheris lateralibus secundis.*

Sphagnum heteromallum polycephalum. Dill. musc. 248. t. 32. f. 6.

Sphagnum cauliferum et ramosum minus hirsutum, capitulis crebris pilosis per ramulorum longitudinem adnascentibus. R. Syn. 105.

Muscus trichoides, capitulis apodibus per longitudinem caulium adnascentibus. Buddl. hort. succ. V. 1. f. 29.

Anglis, Creeping Bog-moss.

Habitat in arborum truncis. In Saltu Enfieldensi. Dill. musc. XI. XII.

PHASCUM. 1052.

Anthera operculata: ore ciliata. Calyptra nulla.

acaule. 1. *PHASCUM acaule, anthera sessili, foliis ovatis imbricatis.*

Phascum acaule anthera sessili. Sp. pl. 1106. Fl. suec. 960.

Sphagnum

Sphagnum acaulon bulbiforme, majus. Dill. musc.
251. t. 32. f. 11.

Sphagnum acaulon, foliis in bulbi formam congestis, majus. Cat. giff. 230. R. Syn. 105.

β *Sphagnum acaulon bulbiforme, minus. Dill. musc.*
252. t. 32. f. 12.

Sphagnum acaulon, foliis in bulbi formam congestis, minus. Cat. giff. 230. R. Syn. 105.

Anglis, Oval Phascum.

Habitat in aggeribus fossarum præsertim solo arenoso.

⊙. I. II.

2. PHASCUM acaule, anthera sessili, foliis subulatis *subulatum* patentibus.

Sphagnum acaulon trichoides. Cat. giff. 229. R. Syn.
105. *Dill. musc.* 251. t. 32. f. 10.

β *Sphagnum acaulon maximum, foliis in centro ciliari-*
bus. Hall. helv. 97. t. 3. f. 3. Dill. musc. 253.
t. 32. f. 13.

Anglis, subulated Phascum.

Habitat in ericetis solo arenoso, β et in pratis arenosis,
circa Snowdon in Arvoniam, Dill. prope Clapham in
comitatu Surriensi. ⊙. III. IV.

3 PHASCUM acaule, anthera pedunculata.

Bryum ampullaceum, foliis serpylli pellucidis collo *peduncu-*
crassiore. Dill. musc. 344. t. 44. f. 4. *latum.*

Bryum erectis gigartinis capitulis foliis serpylli pelli-
cidis obtusis. R. Syn. 93. t. 3. f. 2.

Anglis, Upright Phascum.

Habitat in ericetis humidis. In pascuis Emottmore
dictis, in Lancastriensis comitatus finibus, Dill.
musc. in ericetis montosis Moors dictis in comitatu
Westmorlandico passim. V.

4. PHASCUM caulescens repens. *Sp. pl.* 1107. *repens.*

Sphagnum vermiculare repens, capsulis intra folio-
rum squamosorum alas minimis. Dill. musc. 550.
t. 85. f. 16.

Anglis, Creeping Phascum.

Habitat in arborum truncis in comitatu Eboracensi, D.
Richardson. 4. V.

F O N-

F O N T I N A L I S. 1053.

Anthera operculata. *Calyptra* sessilis, perichætio inclusa.

antipyretica. 1. FONTINALIS foliis complicato-carinatis trifariis acutis, antheris lateralibus. *Fl. suec.* 961. *Sp. pl.* 1107.

Fontinalis triangularis major complicata e foliorum alis capsulifera. *Dill. musc.* 254. t. 33. f. 1.

Fontinalis major foliis triangularibus complicatis capitulis in foliorum alis sessilibus. *R. Syn.* 79.

Musculus aquaticus viticulis longis minus ramosis lucidis, foliis acutis triangularibus cinctis. *Hist. ox.* III. 626. S. 15. t. 6. f. 32.

Anglis, Greater Water-moss.

Habitat in fluviis leniter fluentibus. 4. V.

minor. 2. FONTINALIS foliis ovatis concavis trifariis acutis, antheris terminalibus. *Sp. pl.* 1107.

Fontinalis triangularis minor carinata, e cymis capsuliferis. *Dill. musc.* 257. t. 33. f. 2.

Fontinalis minor foliis triangularibus minus complicatis, capitulis in summis ramulis sessilibus. *R. Syn.* 79.

Anglis, Lesser Water-moss.

Habitat ad ripam Thamesis circa Londinum frequens. 4. IX. X.

squamosa. 3. FONTINALIS foliis imbricatis subulato-lanceolatis capitulis lateralibus. *Sp. pl.* 1108.

Fontinalis squamosa tenuis sericea atrovirens. *Dill. musc.* 250. t. 33. f. 3.

Fontinalis minor lucens. *Baub. hist.* III. 778.

Anglis, Scaly Water-moss.

Habitat in rivulis prope Llanberris in Arvoniam, et in rivulis montosis in Westmorlandia. 4. VIII. IX.

pennata. 4. FONTINALIS foliis bifariis patentibus, capsulis lateralibus. *Sp. pl.* 1108.

Sphagnum pennatum undulatum, vagina squamosa. *Dill. musc.* 250. t. 32. f. 9.

Sphagnum

Sphagnum cauliferum et ramosum, foliis crispis, crebris per caulem capitulis. *Hall. helv.* 96. t. 3. f. 2.

β Sphagnum pennatum undulatum, vagina pilosa. *Dill. musc.* 249. t. 32. f. 8.

Sphagnum pennatum planum. *Dill. musc.* 249. t. 32. f. 7.

Anglis, Feathered Water-moss.

Habitat in arborum truncis in sylvis inter Troutbeck et Ambulfide in Westmorlandia. 4. VIII. IX.

S P L A C H N U M. 1054.

Anthera insidens apophysi maximæ colorata. *Calyptra* caduca.

Stella feminea in distincto individus.

1. SPLACHNUM umbraculo ampullaceo clauso. *Sp. ampullaceum.* pl. 1108. *Amen. acad.* II. 280. *Fl. suec.* 965.

Bryum ampullaceum, foliis thymi pellucidis, collo strictiore. *Dill. musc.* 343. t. 44. f. 3.

Bryum erectis gigartinis capitulis, foliis serpilli pellucidis acutis. *R. Syn.* 93.

Muscus aureus capillaris minor, capitulis geminatis erectis, mutuo incumbitu adnatis. *Hist. ex.* III. p. 629. f. 15. t. 16. f. 10.

Anglis, Common Splachnum.

Habitat in paludibus turfosis. About Hitchin-ferry, near Southampton, and in the Bogs by West-Wickham and Addington, near Croydon, Dr. Sherard. *R. Syn.* III. IV.

P O L Y T R I C H U M. 1055.

Anthera operculata, apophysi minimæ insidens. *Calyptra* villosa.

Stella feminea in distincto individuo.

1. POLYTRICHUM caule simplici, anthera parallelepipedæ. *Sp. pl.* 1109. *Fl. suec.* 966.

Polytrichum quadrangulare vulgare, Yuceæ foliis serratis. *Dill. musc.* 420. t. 54. f. 1.

Polytrichum vulgare et majus capsula quadrangulari. *Cat. giff.* *R. Syn.* 90.

Poly-

Polytrichum aureum majus. *Bauh. pin.* 359. *Park.* 1052.

Muscus capillaris f. *Adiantum aureum majus.* *Ger. Em.* 1559.

β *Polytrichum quadrangulare*, juniperi foliis brevioribus et rigidioribus. *Dill. musc.* 424. t. 54. f. 2.

Polytrichum montanum et minus, capsula quadrangulari. *Cat. giff. R. Syn.* 90.

Polytrichum aureum medium. *Bauh. pin.* 356.

γ *Polytrichum quadrangulare majus*, juniperi foliis pilosis. *Dill. musc.* 426. t. 54. f. 3.

Anglis, Common *Polytrichum*; or, Great golden Maidenhair.

Habitat in sylvis ericetis, uliginosis et sterilissimis. 4. V.

subrotundum. 2. **POLYTRICHUM** caule simplici, anthera subrotunda.

Polytrichum nanum, capsulis subrotundis galeritis, aloes folio non ferrato. *Dill. musc.* 428. t. 55. f. 6.

Polytrichum minus, capsulis subrotundis, calyptra quasi lacera coronatis. *Cat. giff. R. Syn.* 91.

Mnium calyptra villosa. *Sp. pl.* 1112.

Muscus coronatus rigidus minor et humilior capitulis villosis brevioribus. *Hist. ox. III.* 630. f. 15. t. 7. f. 7.

β *Polytrichum parvum*, aloës folio ferrato, capsulis oblongis. *Dill. musc.* 429. t. 55. f. 7.

Anglis, dwarf *Polytrichum*.

Habitat in ericetis. X—III.

urnigerum.

3. **POLYTRICHUM** caule ramosissimo, pedunculis axillaribus. *Sp. pl.* 1109. *Fl. suec.* 967.

Polytrichum ramosum fetis ex alis urnigeris. *Dill. musc.* 427. t. 55. f. 5.

Muscus ramosus erectus, calyptra villosa. *Vaill. parif.* 131. t. 28. f. 13.

β *Polytrichum alpinum ramosum*, capsulis e summitate ellipticis. *Dill. musc.* 427. t. 55. f. 4.

Anglis, branched *Polytrichum*.

Habitat ad radicem montis Caderideris prope Dolgelle in Mervinia. *Dill.* VI.

4. **POLYTRICHUM** caule ramoso, antheris subseffilibus terminalibus, calyptris sursum pilosis. *striatum*.
Polytrichum calyptra striata sursumve pilosa. Fl. suec. ed. I. 869. Dalib. parif. 314.
Polytrichum bryi ruralis facie, capsulis seffilibus majus. Dill. musc. 430. t. 55. f. 8.
Polytrichum capsulis subrotundis pediculis brevissimis insidentibus, calyptra striata, arboreum ramosum, majus. Cat. giff. 222. R. Syn. 91.
Muscus apocarpos arboreus ramosus. R. Hist. III. 40. Vaill. parif. 129. t. 25. f. 5, 6.
 β *Polytrichum bryi ruralis facie, capsulis seffilibus minus. Dill. musc. 430. t. 55. f. 9.*
Polytrichum capsulis subrotundis, pediculis brevissimis insidentibus, calyptra striata, arboreum et terrestre minus ramosum et breve. Cat. giff. 222. R. Syn. 91.
 γ *Polytrichum capsulis seffilibus, foliis brevibus, rectis, carinatis. Dill. musc. 432. t. 55. f. 10.*
 δ *Polytrichum capillaceum crispum, calyptris acutis pilosissimis. Dill. musc. 433. t. 55. f. 11.*
Polytrichum capsulis oblongo-rotundis, calyptris pilosissimis. R. Syn. 91.
Anglis, Tree Polytrichum.
Habitat in arboribus et muris passim. δ in sylvis. II. III.

M N I U M. 1056.

- Anthera operculata, Calyptra lævis.*
Fem. capitulum nudum, pulverulentum, remotum.
 1. **MNIUM** foliis distichis integerrimis. Sp. pl. 1114. *trichomanis*.
Fl. suec. 985.
Mnium trichomanis facie, foliolis integris. R. Syn. 79. Dill. musc. 236. t. 31. f. 5.
Anglis, Fern Mnium.
Habitat in locis udis et umbrosis circa Eltham, Woolwich, et Hampstead. R. Syn. II. III.
 2. **MNIUM** foliis distichis bifidis. Sp. pl. 1114. *fiissum*.
Mnium trichomanis facie, foliolis bifidis. R. Syn. 79. Dill. musc. 237. t. 31. f. 6.
 D d *An-*

Anglis, Forked Mnium.

Habitat in umbrosis passim. III. IV.

jungermannia.

3. MNIMUM foliis distichis, pinnis subtus auriculatis. *Sp. pl.* 1114. *Fl. suec.* 984

Mnium caule procumbente simplici, foliis imbricatis integerrimis alternis antice appendiculatis. *Fl. suec. ed. I.* 914.

Lichenastrum alpinum purpureum, foliis auritis et cochleariformibus. *Dill. musc.* 479. *t.* 69 *f.* 1.

Lichenastrum trichomanis facie, prælongum, foliis concavis unam partem spectantibus. *R. Syn.* 112.

Muscus trichomanoides purpureus, alpinis rivulis innatans. *R. Syn. ed. I.* 18. *ed. II.* 40.

Anglis, purple Mnium

Habitat in rivulis et palustribus; in montibus Cambriæ altissimis, Cader-Idæris, Snowdon et Glyder. *Dill.*

pellucidum.

4. MNIMUM caule simplici, foliis ovatis. *Sp. pl.* 1109. *Fl. suec.* 968.

Mnium serpilli foliis tenuibus et pellucidis. *Dill. musc.* 232. *t.* 31. *f.* 2.

Mnium minus non ramosum, angustioribus et pellucidis foliis. *R. Syn.* 78.

Muscus coronatus minimus, capillaceis foliis, capitulis oblongis. *Vaill. parif.* 130. *t.* 24. *f.* 7.

5. Mnium minimum non ramosum angustissimis et pellucidis foliis. *R. Syn.* 78.

Anglis, transparent Mnium.

Habitat in umbrosis udis; in ericetis juxta Woolwich. *R. Syn.* III.

palustre.

5. MNIMUM caule dichotomo, foliis subulatis. *Sp. pl.* 1110. *Fl. suec.* 970.

Mnium majus, ramis longioribus bifurcatis. *R. Syn.* 78. *Dill. musc.* 233. *t.* 31. *f.* 3.

Muscus palustris flagellis erectis luteolis, raro divisus capitulis oblongis adianti. *Hist. ox. III.* 629. *f.* 15. *t.* 6. *f.* 9.

Anglis, Marsh Mnium.

Habitat

Habitat in paludibus turfosis. In comitatu Westmorlandico copiose, in ericeto Hampstediano. 4. VI. VII.

6. MNIMUM caule ramoso erecto, pedunculis femineis *ramosum.*
axillaribus.

Mnium majus, minus ramosum, capitulis pulverulentis crebrioribus. *R. Syn.* 78. *Dill. musc.* 235. t. 31. f. 4.

Muscus pulverulentis (forsitan variis) capitulis Vernon. *Buddl. hort. succ. Vol. II. f.* 32.

Anglis, branched Mnium.

Habitat in paludibus; in the New-forest, Hampshire, D. Vernon.

7. MNIMUM caule ramoso, androgynum. *Fl. suec.* 971. *androgynum.*
Sp. pl. 1110.

Mnium perangustis et brevibus foliis. *Cat. giff.* 214.
R. Syn. 78. *Dill. musc.* 230. t. 31. f. 1.

Muscus coronatus humilis minimus, foliolis pediculis et capitulis erectis. *Hist. ox.* III. 631. f. 15. t. 7. f. 20.

Anglis, upright Mnium.

Habitat in sylvis et umbrosis udis. In ericeto Hampstediano. III.

B R Y U M. 1057.

Anthera operculata. *Calyptra* lævis. *Filamentum* e tuberculo terminali ortum.

Stella feminea in distincto individuo.

* *Antheris sessilibus.*

1. BRYUM antheris sessilibus terminalibus, calyptra *apocarpum.*
minima. *Sp. pl.* 1115. *Fl. suec.* 986.

Sphagnum nodosum hirsutum incanum. *Dill. musc.* 246. t. 32. f. 5.

Sphagnum cauliferum et ramosum saxatile hirsutum incanum, capitulis virentibus. *Cat. giff.* 229. *R. Syn.* 105.

Anglis, hairy Bryum.

Habitat in saxis et arboribus. In comitatibus Eboracensi et Westmorlandico, frequens. XII.

** *Antheris pedunculatis erectis.*

pomiforme 2. BRYUM antheris erectis sphæricis. *Fl. suec.* 988.

Dalib. parif. 317. *Sp. pl.* 1115.

Bryum capillaceum, capsulis sphæricis. *Dill. musc.* 339. *t.* 44. *f.* 1. *Fl. lapp.* 400.

Bryum trichoides virescens, erectis majusculis capitulis maliformibus. *Cat. giff.* 224. *R. Syn.* 97.

Muscus trichoides minimus sericeus capillaceus, capitulis sphæricis. *Hist. ox. III.* 628. *f.* 15. *t.* 6. *f.* 6.

Anglis, Round-headed Bryum.

Habitat in ericetis. On Hampstead Heath, about Woolwich and Shooter's Hill. *R. Syn.* III.

fontarum. 3. BRYUM antheris obliquis subrotundis, caule simplici geniculis inflexo.

Bryum palustre, scapis teretibus stellatis, capsulis magnis subrotundis. *Dill. musc.* 340. *t.* 44. *f.* 2.

Mnium caule simplici geniculis inflexo. *Fl. lapp.* 414. *suec.* 969. *Sp. pl.* 1110.

Muscus palustris adianto aureo affinis scapis tenuibus, foliolis brevibus. *R. Syn. ed. I.* 19. *III.* 98.

Muscus trichoides palustris erectus coronatus, capitulis sphæricis amplioribus. *Hist. ox. III.* 628. *f.* 15. *t.* 6. *f.* 8.

Anglis, Marsh Bryum.

Habitat ad fontes et in paludibus turfosis. VII. VIII.

pyriforme. 4. BRYUM antheris erectis ovatis, calyptra subulata, surculis acaulibus, foliis ovatis muticis. *Sp. pl.* 1116. *Fl. suec.* 989.

Bryum serpillifolium pellucidum, capsulis pyriformibus. *Dill. musc.* 345. *t.* 44. *f.* 6.

Bryum parvum erectis pyriformibus majusculis capitulis, foliolis serpylli pellucidis. *Cat. giff.* 223. *R. Syn.* 93.

Muscus coronatus humilis stellaris, foliis latiusculis, capitulis pyriformibus erectis turgidiusculis. *Hist. ox. III.* 631. *f.* 15. *t.* 7. *f.* 16.

Anglis, Pear-headed Bryum.

Habitat in pratis arenosis et ad aggeris. III.

5. BRY-

5. BRYUM anthera erecta oblonga minori, calyptris *extincto-*
laxis æqualibus. *Fl. suec.* 990. *Dalib. parif.* 325. *rium.*
Sp. pl. 1116.
Bryum calyptra extincorii forma minus. *Dill. musc.*
349. *t.* 45. *f.* 8.
Bryum erectis capitulis, calyptra laxa conica, foliis
serpilli pellucidis angustioribus. *Cat. giff.* 223. *R.*
Syn. 92.
Muscus capillaceus minimus, calyptra longa conoide
nitida. *Vaill. parif.* 137. *t.* 26. *f.* 1.
β Bryum calyptra extincorii figura, majus et ramosum.
Dill. musc. 350. *t.* 45. *f.* 9.
Anglis, Conic Bryum.
Habitat in pratis arenosis, sed rarius. II.
6. BRYUM antheris erectis subulatis, furculis acauli- *subulatum*
bus *Sp. pl.* 1116. *Fl. suec.* 991
Bryum capulis longis subulatis. *Dill. musc.* 350. *t.*
45. *f.* 10.
Bryum erectis longis et acutis falcatis capitulis, calyp-
tra subfusca, foliis serpilli pellucidis. *Cat. giff.* 223.
R. Syn. 92.
Muscus capillaris, corniculis longissimis incurvis. *R.*
Syn. ed. II. 29
Anglis, subulated Bryum.
Habitat in aggeribus terrenes humidiusculis et sylvis. III.
7. BRYUM antheris erectiusculis, foliis piliferis recur- *rurale.*
vis. *Fl. suec.* 992. *Sp. pl.* 1116. *Dalib. parif.* 320.
It. scan. 118.
Bryum caule erecto, foliis reflexis seta terminatis, ca-
pitulis falcatis. *Fl. lapp.* 397.
Bryum rurale unguiculatum hirsutum elatius et ramo-
sius. *Dill. musc.* 352. *t.* 45. *f.* 12.
Bryum majus erectis falcatis capitulis, foliis latiusculis
extantibus, in pilum canescentem desinentibus. *R.*
Syn. 94.
Muscus erectus hirsutus capitulis longis acutis, ungui-
culis avicularum similibus vulgarissimus. *Hist. ox.*
III. 628. *f.* 15. *t.* 6. *f.* 1.

Anglis, Grey Bryum.

Habitat in muris, tectis, et ad arborum truncos. III.

murale. 8. BRYUM antheris erectis, foliis piliferis rectiusculis, furculis simplicibus cespitosis. *Sp. pl.* 1117. *Fl. suec.* 993

Bryum tegulare humile pilosum et incanum. *Dill. musc.* 355. t. 45. f. 14.

Bryum minus erectis minus falcatis capitulis, foliis latiusculis congestis in pilum canescentem definentibus. *R. Syn.* 94.

β Bryum humile, pilis carens, viride et pellucidum. *Dill. musc.* 356. t. 45. f. 15.

Anglis. Wall Bryum

Habitat in muris, tectis et locis arenosis. III.

parium. 9. BRYUM antheris erectiusculis, pedunculis aggregatis, foliis secundis recurvatis, caule declinato. *Fl. suec.* 995. *Sp. pl.* 1117.

Bryum reclinatum, foliis falcatis, scoparum effigie. *Dill. musc.* 357. t. 46. f. 16.

Bryum erectis capitulis angustifolium, caule reclinato. *Cat. giff.* 222. *R. Syn.* 95.

Muscus coronatus medius, foliis tenuissimis pallidis longioribus, capitulis erectis acutis. *Hist. ox.* III. 630. f. 15. t. 7. f. 11.

Anglis, Broom Bryum.

Habitat in ericetis. III. IV.

undulatum. 10. BRYUM antheris erectiusculis, pedunculis subsolitariis, foliis lanceolatis carinatis undulatis patentibus serratis. *Sp. pl.* 1117. *Fl. suec.* 996.

Bryum phyllitidis folio rugoso acuto, capsulis incurvis. *Dill. musc.* 360. t. 46. f. 18.

Bryum capitulis oblongis rubentibus, foliis oblongis angustis pellucidis rugosis. *Cat. giff.* 222. *R. Syn.* 95.

Anglis, curled Bryum.

Habitat in aggeribus umbrosis et sylvis. XII, I.

11. BRYUM antheris erectiusculis, operculo arcuato, *glaucum*.
foliis erectis, imbricatis, furculis ramosis. *Sp. pl.*

1118. *Fl. suec.* 994

Bryum albidum et glaucum fragile majus, foliis erectis fetis brevibus. *Dill. musc.* 362. t. 46. f. 20.

Bryum trichoides erectis capitulis, albidum fragile.
Cat. giff. 225. *R. Syn.* 97.

Muscus trichoides montanus albidus fragilis. *Hist. ox.*
III 630. f. 15. t. 6. f. 22.

Anglis, white Bryum.

Habitat in ericetis montosis. X, XI.

12. BRYUM antheris erectiusculis, foliis recurvis acutis caule hirsuto. *Sp. pl.* 1118. *Fl. suec.* 997. *pellucidum*.

Bryum palustre pellucidum, capsulis et foliis brevibus recurvis. *Dill. musc.* 364. t. 46. f. 23.

Bryum erectis capitulis subrotundis fuscis, foliis minoribus pellucidis rugosis. *R. Syn.* 96.

β Bryum erectis capitulis brevibus, foliis reflexis.
Dill. musc. 365. t. 46. f. 24.

Anglis, pellucid Bryum

Habitat in palustribus et umbrosis humidis.

- 13 BRYUM antheris erectis, operculo aciformi, foliis *aciculare*.
erectis subsecundis. *Sp. pl.* 1118.

Bryum montanum hemiheterophyllum, operculis acutis. *Dill. musc.* 366. t. 46. f. 25.

Bryum hypnoides erectum montanum, erectis capitulis acutis. *R. Syn.* 94.

Anglis, Mountain Bryum.

Habitat in montosis in Occidentalibus Eboracensis comitatus, D. Richardson; prope Llanberris in Arvoniam, D. Dillenius. VIII.

- 14 BRYUM antheris erectis, foliis setaceis, pedunculis *flexuosum*.
flexuosis. *Sp. pl.* 1118. *Fl. suec.* 998.

Bryum pilosum molle, fetis intortis. *Dill. musc.* 373.
t. 47. f. 33.

Bryum trichoides capitulis erectis, pediculis intortis tenuibus virentibus. *Cat. giff.* 225. *R. Syn.* 97.

Anglis, soft Bryum.

Habitat in rupibus circa Llanberris. *Dill.*

*hetero-
mallum.*

15. BRYUM antheris erectis, foliis setaceis secundis.
Sp. pl. 1118. *Fl. suec.* 999.
 Bryum heteromallum. *Dill. musc.* 375. *t.* 47. *f.* 37.
 Bryum trichoides reclinatis cauliculis, capitulis erectis
 acutis. *R. Syn.* 96.
 β Bryum pilosum interrupte falcatum. *Dill. musc.* 376.
t. 47. *f.* 38.
Anglis, Heath Bryum.
 Habitat in ericetis. X.

tortuosum.

16. BRYUM antheris erectis, foliis setaceis imberbibus :
 arefactione retortis. *Sp. pl.* 1119. *Fl. suec.* 1000.
 Bryum alpinum crinium crispiorum simile, capsula
 barbata. *Hall. belv.* 110 *t.* 4. *f.* 2.
 Bryum cirratum, setis et capsulis longioribus. *Dill.*
musc. 377. *t.* 48. *f.* 40.
 Bryum trichoides longifolium crassiusculis cauliculis,
 capitulis erectis aduncis acutis. *R. Syn.* 98.
Anglis, twisted Bryum.
 Habitat in montosis; in comitatu Eboracensi, D. Rich-
 ardson, in Cambria. *Dill. musc.* VIII. X.

*truncatu-
lum.*

17. BRYUM antheris erectis subrotundis, operculo
 mucronato. *Sp. pl.* 1119 *Fl. suec.* 1001.
 Bryum exiguum, creberrimis capsulis rufis. *Dill.*
musc. 347. *t.* 45. *f.* 7.
 Bryum exiguum erectis parvis subrotundis creberrimis
 capitulis rufis, foliolis serpilli angustis pellucidis.
Cat. giff. 223. *R. Syn.* 94.
 Muscus coronatus minimus, foliis et capitulis oblon-
 gis in pediculis brevissimis. *Hist. ox.* III. 631. *f.*
 15. *t.* 7. *f.* 18.
 β Bryum parvum erectis subrotundis majusculis capitu-
 lis subfuscis, foliis serpilli pellucidis. *Cat. giff.* 223.
R. Syn. 93.
Anglis, brown Bryum.
 Habitat in pratis et pascuis. II.

viridulum.

18. BRYUM antheris erectis ovatis, foliis lanceolatis
 acuminatis imbricato-patulis. *Fl. suec.* 1002. *Sp.*
pl. 1119.

Bryum

Bryum capillaceum breve, pallide et læte virens, capsulis ovatis. *Dill. musc.* 380. t. 48. f. 43.

Bryum trichoides exile, erectis capitulis in pediculis brevissimis. *Cat. giff.* 224. *R. Syn.* 97.

Anglis, green *Bryum*.

Habitat in aggerribus fossarum et ad margines sepium arenosos. XII, II.

19. BRYUM antheris erectis, foliis arefactione revolutis.

Bryum cirratum et stellatum, tenuioribus foliis. *Dill. musc.* 379. t. 48. f. 42.

Bryum trichoides exile, erectis capitulis in pediculis longioribus rubris. *Cat. giff.* 224. *R. Syn.* 97.

Mnium foliis arefactione revolutis. *Sp. pl.* 1111.

Muscus coronatus minimus, capillaceis foliis, capitulis oblongis. *Hist. ox.* III 631. f. 15. t. 7. f. 19.

β *Bryum trichoides* erectis sublongis capitulis, extremitatibus per siccitatem stellatis. *R. Syn.* 98.

Bryum cirratum, setis et capsulis brevioribus et pluribus. *Dill. musc.* 378. t. 48. f. 41.

γ *Bryum perangustis* et brevissimis foliis, extremitatibus stellatis. *Cat. giff.* 226. *R. Syn.* 98.

Anglis, starry *Bryum*.

Habitat in sylvis et ad sepes. III.

20. BRYUM antheris erectis ore non ciliato foliis carinatis patulis.

Bryum tenue imberbe et pallidum, foliis crebrioribus. *Dill. musc.* 382. t. 48. f. 46.

Bryum perangustis foliis et cauliculis, foliis crebrioribus et circa extremitates magis congestis, capitulis erectis, ad summitem magis egredientibus. *Cat. giff.* 225. *R. Syn.* 99.

Anglis, Beardless *Bryum*.

Habitat in locis arenosis gramineis et ad sepes. III.

21. BRYUM antheris erectis, operculis filiformibus setaceum. longitudine capitulorum.

Bryum stellare nitidum pallidum, capsulis tenuissimis. *Dill. musc.* 381. t. 48. f. 44.

Bryum

Bryum trichoides exile pallidum, erectis capitulis e furculis annotinis egredientibus. *R. Syn.* 96

Minium antheris erectis, operculis filiformibus longitudine capitulorum. *Sp. pl.* 1111 *Fl. suec.* 974.

β *Bryum stellare* lacustre, foliis rubris capillaceis. *Dill. musc.* 382. t. 48. f. 45.

Anglis, pale *Bryum*.

Habitat in aggeribus fossarum, sepibus et muris. III.

unguiculatum.

22. *BRYUM* antheris erectis oblongis, pedunculis axillaribus, foliis erectis acutis carinatis.

Bryum unguiculatum et *barbatum*, furculis in summitate crassioribus. *Dill. musc.* 383. t. 48. f. 47.

Bryum angustis viridibus foliis, capitulis erectis, brevibus pediculis insidentibus, calyptra falcata vel avium unguiculos referente *R. Syn.* 96

Muscus capillaris, capitulis erectis lageniformibus. *Buxb. cent. II.* 9. t. 2. f. 9.

Anglis,

Habitat in muris et locis arenosis. III.

Obs. *Caule erecto subramoso. Foliis luteo-viridibus. Pedunculis brevibus ex alis foliorum enascentibus.*

paludosum 23. *BRYUM* acaulon, foliolis fetaceis, antheris obtusissimis patulis. *Sp. pl.* 1119. *Fl. suec.* 1004.

Bryum trichoides acaulon palustre minimum, fetis et capsulis brevissimis. *Dill. musc.* 387. t. 49. f. 53.

Anglis, dwarf *Bryum*.

Habitat in palustribus in comitatu Eboracensi. D. Richardson. III.

Hypnoides 24. *BRYUM* antheris erectis, caule erectiusculo, ramis lateralibus brevibus fertilibus. *Fl. lapp.* 396. *suec.* 1003. *Sp. pl.* 1119.

Bryum hypnoides, hirsute canescens, vulgare. *Dill. musc.* 368 t. 47. f. 27.

Bryum trichoides erectis capitulis, lanuginosum. *Cat. giff.* 224. *R. Syn.* 97.

β *Bryum hypnoides*, hirsute virecens fasciculare alpinum. *Dill. musc.* 370. t. 47 f. 28

γ *Bryum hypnoides* alpinum, fetis et capsulis exiguis. *Dill. musc.* 370. t. 47. f. 29.

♂ *Bryum*

- ♂ *Bryum hypnoides alpinum*, operculis obtusis. *Dill. musc.* 271. t. 47. f. 30.
- ε *Bryum hypnoides*, ericæ facie, capsulis barbatis alpinum. *Dill. musc.* 371. t. 47. f. 31.
Bryum hypnoides capitulis plurimis erectis, non lanuginosum. *R. Syn.* 478.
- ζ *Bryum hypnoides polycephalon lanuginosum alpinum*. *Dill. musc.* 372. t. 47. f. 32.
Bryum hypnoides capitulis plurimis erectis, lanuginosum. *R. Syn.* 97.
Muscus terrestris vulgari similis lanuginosus. *R. Syn. ed. I.* 18. *ed. II.* 37. *Hist. ox. III.* 625. f. 15. t. 5. f. 7.
- η *Bryum hypnoides aquaticum calyptris nigris acutis*. *Dill. musc.* 367. t. 46. f. 26.
Bryum hypnoides repens aquaticum, erectis capitulis acutis. *R. Syn.* 94.
Anglis, Woolly *Bryum*.
Habitat in montosis in comitatibus Eboracensi, Westmorlandico et Cumberlandico frequens, in monte Snowdon dicto in Cambria copiose: *Dill.* XII.
25. **BRYUM** antheris erectis, pedunculis arefactione verticillatoccontortis foliis piliferis, furculis fastigiatis. *Sp. pl. tum.* 1120.
Bryum pilosum verticillatum. *Dill. musc.* 374. t. 47. f. 35.
Brium trichoides brevifolium angustis cauliculis, capitulis erectis parvis et minus aduncis. *R. Syn.* 98.
- β *Bryum pilosum*, sphagni subulati facie. *Dill. musc.* 374. t. 47. f. 34.
Anglis, verticillated *Bryum*.
Habitat in montosis in comitatn Eboracensi, D. Richardson. *R. Syn. in rupibus maritimis prope Bangor.* D. Brewer. *Dill.*
26. **BRYUM** antheris erectis subrotundis, axillaribus, palustre. foliis subulatis distantibus.
Bryum palustre æstivum, conservæ facie. *Dill. musc.* 375. t. 47. f. 36.
Bryum angustissimis foliis crebrioribus, capitulis erectis

tis brevibus, pediculis e furculis novis et longis enascentibus. *R. Syn.* 99.

Anglis, Water Bryum.

Habitat in palustribus in comitatu Eboracenfi. *D. Richardson.* III.

purpureum.

27. BRYUM antheris obliquis, caule dichotomo, axillis pedunculiferis, foliis carinatis.

Bryum tenue stellatum, setis purpureis. *Dill. musc.* 386. t. 49. f. 51.

Bryum perangustis foliis et cauliculis, foliis crebrioribus et circa summitates magis congestis, capitulis erectis e furculis annotinis egredientibus. *Cat. giff.* 226. *R. Syn.* 99.

β Bryum unguiculatum et barbatum, tenuius et stellatum. *Dill. musc.* 384. t. 48. f. 48.

γ Bryum tenue barbatum, foliis angustioribus et rarioribus. *Dill. musc.* 385. t. 48. f. 49.

Bryum perangustis foliis et cauliculis foliis rarioribus cinctis, capitulis erectis, e furculis annotinis egredientibus. *Cat. giff.* 225. *R. Syn.* 99.

δ Bryum polytrichoides palustre, setis longioribus rubris sericeis. *Dill. musc.* 387. t. 49. f. 52.

Bryum perangustis crebrioribus foliis, capitulis erectis longiusculis, pediculis e furculis annotinis nascentibus. *R. Syn.* 99.

Anglis, purple Bryum.

Habitat in muris ericetis et pascuis glareosis. III. IV.

*** *Antheris nutantibus.*

aureum. 28. BRYUM antheris pendulis turbinatis, stipite filiformi, foliis ensiformibus nitidis.

Bryum trichoides aureum, capitulis pyriformibus nutantibus. *Cat. Nottingh. app. Dill. musc.* 391. t. 50. f. 60.

Minium antheris pendulis turbinatis, stipite filiformi, floribus femineis setiferis. *Sp. pl.* 1112.

Anglis, golden Bryum.

Habitat in rupibus Parcì Nottinghamensis. *D. Car. Deering.* IV. V.

29. BRY-

29. BRYUM antheris pendulis, furculis cylindricis imbricatis lævibus. *Sp. pl.* 1120. *Fl. suec.* 1008. *argenteum*
 Bryum pendulum julaceum argenteum et fericeum.
Dill. musc. 392. t. 50. f. 62.
 Bryum capitulis subrotundis reflexis, cauliculis teretibus argenteis. *Cat. giff.* 226. *R. Syn.* 100.
 Muscus minimus e viridi argenteus, capitulis oblongis cernuis. *Hist. ox.* III. 629. f. 15 t. 6. f. 17.
 β Bryum pendulum, furculis teretibus viridibus. *Dill. musc.* 394. t. 50. f. 63.
Anglis, silver Bryum.
 Habitat *in tectis, muris et apricis.* XII.
30. BRYUM antheris subrotundis, pedunculis reflexis, pulvinatoliis piliferis. *Fl. suec.* 1009. *Sp. pl.* 1121. *Datum.*
lib. parif 321.
 Bryum orbiculare pulvinatum hirsutiae canescens, capsulis immerfis. *Dill. musc.* 395. t. 50. f. 65.
 Bryum trichoides hirsutiae canescens, capitulis subrotundis reflexis, in perbreuibis pediculis. *Cat. giff.* 226. *R. Syn.* 100.
 Muscus trichoides hirsutus, capitulis oblongis reflexis, pediculis brevibus insidentibus. *R. Syn. ed. I.* 243.
Hist. ox. III. 629. f. 15. t. 6. f. 21.
Anglis, grey Bryum.
 Habitat *in muris et tectis.* IV.
31. BRYUM anthera oblonga nutante, foliis subulatis. *rubrum.*
 caule simplicissimo, medio pedunculifero.
 Bryum trichoides, capsulis rubris cernuis. *Dill. musc.* 390. t. 50. f. 59.
Anglis, red Bryum.
 Habitat *in locis gramineis sed varius.*
- 32 BRYUM antheris pendulis, foliis lanceolatis acuminato-setaceis, pedunculis longissimis. *Sp. pl.* 1121. *um.*
Fl. suec. 1010.
 Bryum pendulum ovatum caespitium et pilosum, seta bicolori. *Dill. musc.* 396. t. 50. f. 66.

Bryum

Bryum trichoides capitulis reflexis, pediculis ima medietate rubris, summa luteo-virentibus. *Cat. giff.* 226. *R. Syn.* 100.

Muscus trichoides capitulo parvo reflexo, pediculo ima medietate rubro, summa luteo-viridi. *Hist. ox.* III. 629. *f.* 15. *t.* 6. *f.* 15.

Anglis, matted *Bryum*.

Habitat in muris antiquis et glareosis. III.

capillare. 33. *BRYUM* antheris pendulis, foliis ovatis fetiferis carinatis, pedunculis longissimis. *Sp. pl.* 1121.

Bryum foliis latiusculis congestis, capsulis longis nutantibus. *Dill. musc.* 398. *t.* 50. *f.* 67.

Bryum capitulis reflexis, foliolis latiusculis congestis. *Cat. giff.* 227. *R. Syn.* 100.

Muscus capillaris foliolis latiusculis congestis, capitulis oblongis reflexis. *Hist. ox.* III. 629. *f.* 15. *t.* 6. *f.* 19.

β *Bryum* capitulis reflexis foliis congestis latioribus et pellucidis. *R. Syn.* 101.

Anglis, capillary *Bryum*.

Habitat in sylvis et ericetis solo glareoso. III.

annotinum 34. *BRYUM* antheris oblongis nutantibus, foliis ovatis acuminatis pellucidis, pedunculis subradicalibus.

Bryum annotinum lanceolatum pellucidum, capsulis oblongis pendulis. *Dill. musc.* 399. *t.* 50. *f.* 68.

Mnium foliis ovatis acuminatis pellucidis, pedunculis radicalibus, anthera nutante. *Sp. pl.* 1111.

Anglis, long-leaved *Bryum*.

Habitat in sylvis et umbrosis humidis. III.

carneum. 35. *BRYUM* antheris pendulis subglobosis, foliis acutis alternis. *Sp. pl.* 1122. *Fl. suec.* 1011.

Bryum lanceolatum pellucidum capsulis rotundis pendulis, carneis. *Dill. musc.* 400. *t.* 5. *f.* 69.

Bryum nitidum foliis serpilli pellucidis angustioribus reflexis capitulis subrotundis, carnei coloris. *R. Syn.* 102.

β *Bryum* nitidum perangustis serpillinis pellucidis foliis, reflexis capitulis subrotundis virentibus. *R. Syn.* 102.

Anglis,

Anglis, shining Bryum.

Habitat in locis udis ; ad fossulas pratorum inter Blackheath et Eltham, et prope Darking in Surriensi comitatu. Dill. musc. III.

36. BRYUM antheris oblongis pendulis, calyptra re-crudum. curvata, foliis ovatis acutis pellucidis.

Bryum pendulum hornum molle, foliis et lanceolatis et gramineis. Dill. musc. 401. t. 50. f. 70.

Mnium antheris pendulis, calyptra recurvata, foliis pellucidis. Sp. pl. 1112.

Anglis, Spear-leaved Bryum.

Habitat in locis udis in Cambra. Dill. musc. III.

37. BRYUM anthera oblonga pendula: foliis ovatis a-*alpinum*. cuminatis carinatis, furculis ramosis, axillis pedunculiferis.

Bryum hypnoides pendulum sericeum, coma insigni atro-rubente. Dill. musc. 394. t. 50. f. 64.

Anglis, Alpine Bryum.

Habitat prope Castellum quoddam dirutum, non longe a Llanberris, in rupibus tenui terra tectis. Dill. musc.

38. BRYUM antheris oblongis nutantibus, pedunculo *hornum*. curvato, foliolis ovatis margine scabris.

Bryum stellare hornum sylvarum, capsulis magnis nutantibus. Dill. musc. 402. t. 51. f. 71.

Bryum nitidum capitulis majoribus reflexis, calyptra imum virgente, pediculis oblongis e cauliculis novis egredientibus. R. Syn. 102.

Mnium antheris pendulis, pedunculo curvato, furculo simplici, foliolis margine scabris. Sp. pl. 1112. Fl. suec. 977.

- β Bryum nitidum foliis serpilli angustioribus, medium. Cat. giff. 228. R. Syn. 103.

Muscus polytrichoides humilior, foliis brevioribus raris, pallide viridantibus et vix pellucidis. Hist. ox. III. 630. f. 15. t. 6. f. 3.

Anglis, rough Bryum.

Habitat in sylvis et ericetis. IV. V.

- trique-
trum.* 39. BRYUM antheris oblongis pendulis, foliis trifariis
subulato-lanceolatis carinatis, furculis ramosis.
Bryum annotinum palustre, capsulis ventricosus pen-
dulis. *Dill. musc.* 404. *t.* 51. *f.* 72.
Bryum nitidum foliis serpilli angustioribus, majus.
Cat. giff. 228. *R. Syn.* 102.
Mnium foliis trifariis patulis subulato-lanceolatis ca-
rinatis. *Sp. pl.* 1114.
Muscus capillaris major et elatior, capitulis longis et
obtusis deorsum reflexis et veluti pendulis præaltis
pediculis rubris. *Hist. ox.* III. 629. *f.* 15. *t.* 6.
f. 20.
β Bryum lanceolatum bimum, fetis et capsulis longis
pendulis. *Dill. musc.* 405. *t.* 51. *f.* 73.
Bryum nitidum foliis serpilli pellucidis angustis, capi-
tulis tumidis nutantibus, præaltis pediculis e fur-
culis annotinis egredientibus. *R. Syn.* 102.
γ Bryum palustre complicatum rubens, capsulis turbina-
tis pendulis. *Dill. musc.* 406. *t.* 51. *f.* 74.
Bryum nitidum rubens capitulis reflexis, foliis angus-
tis pellucidis, cauliculis proliferis. *R. Syn.* 102.
Anglis, triangular Bryum.
Habitat in paludibus turfosis. IV.
- lygme-
tricum.* 40. BRYUM anthera nutante, calyptra reflexa tetrago-
na.
Bryum bulbiforme aureum, calyptra quadrangulati,
capsulis piriformibus nutantibus. *Dill. musc.* 407.
t. 52. *f.* 75.
Bryum aureum, capitulis reflexis piriformibus, ca-
lyptra quadrangulati, foliis in bulbi formam con-
gestis. *R. Syn.* 101.
Muscus capillaris, s. adiantum aureum minus. *Ger.*
Em. 1559.
Polytrichon aureum minus. *Park.* 1052.
Anglis, yellow Bryum.
Habitat in sylvis, ericetis, hortis et ad semitas. IV.
- serpyllif-
lium.* 41. BRYUM antheris pendulis, pedunculis aggregatis,
foliis patentibus pellucidis.

Mnium pedunculis aggregatis, foliis obovatis integerrimis obtusis punctatis. *Sp. pl.* 1113. punctatum.

Bryum pendulum terpillifolio rotundiore pellucido, capsulis ovatis. *Dill. musc.* 416. *t.* 53. *f.* 81.

Bryum nitidum serpilli rotundis et latoribus foliis pellucidis. *R. Syn.* 103.

Muscus polytrichoides palustris major, serpilli latiori folio pellucido. *Hist. ox. III.* 627. *f.* 15 *t.* 6. *f.* 39, 40.

β Mnium pedunculis aggregatis foliis alternis acutis serratis. *Sp. pl.* 1113. cuspidatum.

Bryum pendulum, foliis variis pellucidis capsulis ovatis. *Dill. musc.* 413. *t.* 53. *f.* 79.

γ Bryum pendulum, serpilli folio longiore pellucido, capsulis oblongis cuspidatis. *Dill. musc.* 416. *t.* 53. *f.* 80.

δ Mnium pedunculis aggregatis foliis rosaceo-congestis lanceolatis acutis. *Sp. pl.* 1113. proliferum.

Bryum stellare roseum majus capsulis ovatis pendulis. *Dill. musc.* 411. *t.* 52. *f.* 77.

Bryum roseum majus, foliis oblongis. *R. Syn.* 92.

Muscus coronatus humilis, foliolis latoribus stellatim nascentibus donatus. *Hist. ox. III.* 630. *f.* 15. *t.* 7. *f.* 9.

ε Bryum stellare roseum pendulum minus, capsulis et fetis longioribus. *Dill. musc.* 412. *t.* 52. *f.* 78.

Bryum roseum minus, foliis subrotundis. *R. Syn.* 92.

Muscus parvus stellaris. *Baub. pin.* 361. *Ger. Em.* 1563. *Park.* 1308.

ζ Mnium pedunculis aggregatis, foliis oblongis undulatis. *Sp. pl.* 1113. undulatum.

Bryum dendroides polycephalon, phyllitidis folio undulato pellucido, capsulis ovatis pendulis. *Dill. musc.* 410. *t.* 52. *f.* 76.

Bryum nitidum foliis oblongis undulatis, capitulis cernuis arbusculam referens. *Cat. giff.* 227. *R. Syn.* 103.

Muscus ad polytrichoidem accedens ramosus, foliis longis lucidis et veluti crispis. *Hist. ox. III.* 630. *f.* 15 *t.* 6. *f.* 1.

Anglis, Thyme-leaved Bryum.

Habitat in sylvis et ericetis humidiusculis. III, IV.

HYPNUM. 1058.

Anthera operculata. Calyptra lævis. Filamentum laterale ortum e perichætio.

Fem. Pistilli solitarii, ex alis foliorum, nudæ plumulæ.

* Frondibus pinnatis.

bryoides. 1. HYPNUM fronde simplicissima pinnata lanceolata apice pedunculifera. *Sp. pl.* 1123. *Fl. suec.* 1012.

Hypnum taxiforme exiguum, versus summitatem capsuliferum. *Dill. musc.* 262. t. 34. f. 1.

Hypnum repens filicifolium non ramosum, pediculis brevibus versus foliorum summitatem egredientibus. *R. Syn.* 88.

Muscus polytrichoides exiguus, capitulis in extremis cauliculis s. foliis, subrotundis erectis. *Hist. ex.* III. 629. f. 15. t. 6. f. 11.

Anglis, Fern Hypnum.

Habitat in umbrosis et ad sæpes. II.

taxifolium 2. HYPNUM fronde simplicissima pinnata lanceolata basi pedunculifera. *Sp. pl.* 1122. *Fl. suec.* 1013.

Hypnum taxiforme minus, basi capsulifera. *Dill. musc.* 263 t. 34. f. 2.

Hypnum repens filicifolium non ramosum, pediculis brevioribus ad radicem egredientibus. *R. Syn.* 88.

Muscus pennatus, capitulis adianti. *Vaill. parif.* 136. t. 24. f. 11.

Anglis, Yew-leaved Hypnum.

Habitat in sylvis et umbrosis. II.

denticulatum. 3. HYPNUM fronde pinnata simplici : pinnis duplicatis, basi pedunculifera. *Sp. pl.* 1122. *Fl. suec.* 1014.

Hypnum denticulatim pennatum, pinnulis duplicatis recurvis. *Dill. musc.* 266. t. 34 f. 5.

Hypnum repens filicifolium non ramosum, pediculis et capitulis longioribus ad radicem egredientibus, foliolis utrinque duplicatis. *Cat. gift.* 218. *R. Syn.* 88.

Muscus

Muscus pennatus capitulis adianti. Hist. ox. III.
626. f. 15. t. 6. f. 36.

Anglis, Feathered Hypnum.

Habitat in sylvis et umbris humidis. IV.

4. HYPNUM fronde pinnata ramosa procumbens: folioli-
olis acutis, medio pedunculifera. *sylvaticum.*

Hypnum denticulatim pennatum, pinnulis simplicibus
rectioribus. Dill. musc. 267. t. 34. f. 6

Hypnum repens filicifolium ramosum, pedunculis et
capitulis longioribus e foliorum alis egredientibus,
foliolis utrinque simplicibus. R. Syn. 88.

Anglis, Wood Hypnum.

Habitat in sylvis ad arborum radices. IV.

5. HYPNUM fronde pinnata ramosa erecta medio pe-
dunculifera. *Sp. pl. 1123. adiantoides.*

Hypnum taxiforme palustre ramosum majus et erec-
tum. Dill. musc. 264. t. 34. f. 3.

Hypnum erectum filicifolium ramosum, pinnulis acu-
tis. R. Syn. 87.

Muscus taxiformis ramosus. Vaill. parif. 136. t. 28.
f. 5.

Anglis, upright Hypnum.

Habitat in paludosis. III.

6. HYPNUM fronde pinnata ramosa: foliolis imbrica-
tis acutis complicatis compressis. *Sp. pl. 1123. Fl. tum.*
succ. 1015.

Hypnum pennatum compressum et splendens, capsu-
lis ovatis. Dill. musc. 268. t. 34. f. 7.

Hypnum repens filicifolium ramosum, ramulis appres-
sis et magis complanatis. Cat. giff. 218. R. Syn.
87.

- β *Hypnum pennatum trichomanoides splendens ramo-*
sum. Dill. musc. 269. t. 34. f. 8.

Hypnum erectum filicifolium ramosum pinnulis obtu-
sis. R. Syn. 87.

Anglis, Flat Hypnum.

Habitat ad truncos arborum in sylvula Hampstead
Wood dicta. III. IV.

** *Surculis vagis.*

- lucens.* 7. HYPNUM furculis ramosis, frondibus subpinnatis foliolis punctatis *Sp. pl.* 1124.
 Hypnum pennatum aquaticum lucens, longis latisque foliis. *Dill. musc.* 270. t. 34. f. 10.
 Hypnum repens filicifolium ramosum, foliolis majoribus magisque crebris. *R. Syn.* 88.
Anglis, shining Hypnum.
 Habitat in umbrosis udis et ad fossas sed rarius. III.
- undulatum.* 8. HYPNUM furculis ramosis, frondibus subpinnatis, foliolis undulatis complicatis. *Sp. pl.* 1124.
 Hypnum pennatum undulatum, lycopodii instar sparsum. *Dill. musc.* 271. t. 36. f. 11.
 Hypnum repens crispum. cauliculi compressis, lycopodii in morem per terram sparsis. *K. Syn.* 88.
Anglis, waved Hypnum.
 Habitat in sylvis, in Bishop's-Wood, Hampstead, Mr. Buddle. *R. Syn.* V.
- crispum.* 9. HYPNUM furculis ramosis, frondibus subpinnatis : foliolis undulatis planis. *Sp. pl.* 1124.
 Hypnum pennatum undulatum crispum, setis et capulis brevibus. *Dill. musc.* 273. t. 36. f. 12.
 Hypnum repens crispum, ramulis compressis, filicinorum more dispositis. *Cat. giff.* 217. *R. Syn.* 89.
Anglis curled Hypnum.
 Habitat ad ripam Thamesis prope Gravesend, D. Doody. *R. Syn.* in rupe S. Vincentii prope Bristolium. *Dill.* III.
- triquetrum.* 10. HYPNUM ramis vagis recurvis, foliis ovatis recurvatis patulis. *Sp. pl.* 1124. *Fl. succ.* 1016.
 Hypnum vulgare triangulum maximum et pallidum. *Dill. musc.* 293. t. 38. f. 28.
 Hypnum repens triangularibus majoribus et pallidioribus foliis. *Cat. giff.* 219. *R. Syn.* 80.
Anglis, triangular Hypnum.
 Habitat in sylvis et sepibus. II.

11. HYPNUM furculis vagis ramosissimis, ramis filiformibus, antheris obliquis. *filiciforme.*

Hypnum sericeum ramosius et tenuius capsulis acuminatis. *Dill. musc. 327. t. 42. f. 62.*

Hypnum repens trichoide terrestre, priori viridius et minus, capitulis cernuis minus tumidis. *Cat. giff. 216. R. Syn. 84.*

Usnea seu Muscus cranio humano innatus. *Dood. R. Syn. ed. II. 36.*

Anglis, green Hypnum.

Habitat in truncis arborum. II.

12. HYPNUM ramis vagis subrepentibus, foliolis ovatis mucronatis imbricatis. *Sp. pl. 1124. rutabulum.*

Hypnum dentatum vulgatissimum, operculis obtusis. *Dill. musc. 295. t. 38. f. 29.*

Hypnum repens triangularibus angustioribus foliis. *R. Syn. 80.*

Muscus denticulatus minor sericeus nostras, capitulis adianti. *Hist. ox. III. 626. f. 15. t. 6. f. 35.*

β Hypnum palustre erectum breve, foliis brevibus angustis tenuibus confertis. *Cat. giff. 219. R. Syn. 83.*

γ Hypnum vulgare dentatum, operculis cuspidatis. *Dill. musc. 297. t. 38. f. 30.*

Hypnum repens, triangularibus minoribus foliis pediculis et capitulis brevioribus et tumidioribus, minus. *R. Syn. 80.*

Anglis, Toothed Hypnum.

Habitat in ericetis, sylvis et sepibus. XII.

13. HYPNUM furculis vagis procumbentibus, foliis subulatis, antheris obliquis. *lutescens.*

Hypnum sericeum, furculis longioribus et rectioribus, capsulis incurvis. *Dill. musc. 325. t. 42. f. 60.*

Anglis, yellow Hypnum.

Habitat in collibus cretaceis inter Northfleet et Gravesend. *Dill I, II.*

*** *Surculis pennatis.*

- filicinum.* 14. HYPNUM *surculis pennatis*: ramulis distantibus, foliis imbricatis incurvis acutis secundis. *Sp. pl.* 1125. *Fl. suec.* 1017.
 Hypnum repens *filicinum crispum.* *R. Syn.* 85.
Dill. musc. 282. *t.* 36. *f.* 19.
 β Hypnum repens *filicinum*, *trichoides palustre.* *R. Syn.* 83. *Dill. musc.* 286. *t.* 35. *f.* 21.
 γ Hypnum repens *trichoides palustre vernalis fuscum* capitulis oblongis incurvis. *R. Syn.* 85.
Anglis, Bog Hypnum.
Habitat in palustribus. III.
- proliferum.* 15. HYPNUM *surculis proliferis plano-pinnatis*, pedunculis aggregatis. *Sp. pl.* 1125. *Fl. lapp.* 405. *suec.* 1018.
 Hypnum *filicinum*, *tamarisci foliis minoribus non splendentibus.* *Dill. musc.* 276. *t.* 35. *f.* 14.
 Hypnum repens *filicinum minus*, *luteo-virens.* *Cat. giff.* 217. *R. Syn.* 86.
Anglis, Proliferous Hypnum.
Habitat in ericetis et sylvis. II.
- parietinum.* 16. HYPNUM *surculis plano-pinnatis continuatis*, pedunculis aggregatis. *Fl. lapp.* 406. *suec.* 1020. *Sp. pl.* 1125.
 Hypnum *filicinum*, *tamarisci foliis majoribus splendentibus.* *Dill. musc.* 274. *t.* 35. *f.* 13.
 Hypnum repens *filicinum*, *veluti spicatum.* *Cat. giff.* 217. *R. Syn.* 86.
 Muscus *filicinus major.* *Ger. Em.* 1561. *Park.* 1309. *Bauh. pin.* 360.
Anglis, Wall Hypnum.
Habitat in sylvis et umbris. III.
- prælongum.* 17. HYPNUM *surculis subpinnatis decumbentibus*: ramulis remotis, foliis ovatis, antheris cernuis. *Sp. pl.* 1125. *Fl. suec.* 1021.

Hypnum filicinum, triangularibus parvis foliis, prælongum. *Cat. giff.* 219. *R. Syn.* 80. *Dill. musc.* 278. *t.* 35. *f.* 15.

Anglis, Trailing Hypnum.

Habitat in arborum truncis, lignis putrescentibus et terris. II, III.

18. HYPNUM furculis pinnatis repentibus, ramis confertis foliis imbricatis subulatis, antheris erectis. *plumosum.*

Hypnum repens filicinum plumosum. *R. Syn.* 86. *Dill. musc.* 280. *t.* 35. *f.* 16.

Muscus sericeus quidam, forte arboreus. *Baub. hist.* III. 765.

Anglis, Winged Hypnum.

Habitat in muris lateritiis circa Londinum, et ad arborum truncos et radices in sylva Enfieldensi. II.

19. HYPNUM furculis pinnatis : ramulis approximatis : apicibus recurvis. *Fl. suec.* 1022. *Sp. pl.* *crista castrensis.* II 125.

Hypnum filicinum, cristam castrensem repræsentans. *Dill. musc.* 284. *t.* 36. *f.* 20.

Muscus filicifolius luteus, folio crasso et undulato. *D. Richardson.* *R. Syn.* 86. *sub num.* 32.

Muscus terrestris repens subflavus, foliolis crispis minoribus, ramulisque densius confertis. *Hist. ox.* III. 625. *f.* 15. *t.* 5. *f.* 12.

Muscus cristam castrensem repræsentans, flavescens nemorosus ramosus castubicus. *Hist. ox.* III. 625. *f.* 15. *t.* 5. *f.* 8.

Anglis, crested Hypnum.

Habitat in collibus cretaceis inter Gravesend et Northfleet in Cantio. *Dill.* II.

20. HYPNUM furculis pinnatis compressis : foliis acutis incurvis, antheris erectiusculis ovatis. *compressum.*

Hypnum filicinum sericeum, molle et pallidum mucronibus aduncis. *Dill. musc.* 286. *t.* 36. *f.* 22.

Hypnum arboreum erectum, fruticuli specie ramulis compressis. *Cat. giff.* 220.

Anglis, Shrubby Hypnum.

Habitat in arborum truncis in sylva prope Tottenham non longe a Londino. *Dill.* IV.

abietinum.

21. HYPNUM furculis pinnatis teretiusculis remotis inæqualibus. *Fl. lapp.* 408. *suec.* 1023. *Sp. pl.* 1126.

Hypnum lutescens, alis subulatis tenacibus. *Dill. musc.* 280. *t.* 35. *f.* 17.

Hypnum repens filicinum trichoides montanum, ramulis teretibus lutescentibus non divisis. *Cat. giff.* 218. *R. Syn.* 86.

Muscus terrestris furculis filamentosis tenacibus abietinis, semel tantum divisis. *Hist. ox. III.* 626. *f.* 15. *t.* 5. *f.* 22.

Anglis, Firr Hypnum.

Habitat in montosis. In comitatu Oxoniensi.

**** *Foliis reflexis*

cupressiforme.

22. HYPNUM furculo subpinnato, foliis secundis recurvis apice subulatis. *Fl. suec.* 1024. *Sp. pl.* 1126.

Hypnum crispum cupressiforme, foliis aduncis. *Dill. musc.* 287. *t.* 37. *f.* 23.

Hypnum repens crispum cupressiforme. *R. Syn.* 89.

- β Hypnum repens trichoides terrestre, foliolis uno versu dispositis. *R. Syn.* 85.

Anglis, Cypress Hypnum.

Habitat in sylvis ad radices arborum. II.

aduncum.

23. HYPNUM furculo erectiusculo subramoso, foliis secundis recurvis subulatis ramis recurvatis. *Sp. pl.* 1126. *Fl. suec.* 1025.

Hypnum palustre erectum summitatibus aduncis. *R. Syn.* 82. *Dill. musc.* 292. *t.* 37. *f.* 26.

Anglis, hooked Hypnum.

Habitat in palustribus. V.

scorpioides

24. HYPNUM ramis vagis procumbentibus recurvis, foliis secundis acuminatis. *Sp. pl.* 1127. *Fl. suec.* 1026

Hypnum scorpioides palustre magnum, Lycopodii instar sparium. *Dill. musc.* 290. *t.* 37. *f.* 25.

Anglis, Scorpion Hypnum.

Habitat

Habitat in paludibus. Prope Bishop's-Castle in agro Salopiensi, et circa Pentir, tribus milliaribus a monte Glyder in Wallia. Dill.

25. HYPNUM furculis repentibus, ramis vagis tereti viticulis, foliis patulis acuminatis. *Sp. pl.* 1127. *sum.*

Hypnum subhirsutum, viticulis gracilibus erectis, capsulis teretibus. *Dill. musc.* 307. *t.* 39. *f.* 43.

Hypnum repens trichoides arboreum majus, capsulis et furculis erectis, minus ramosis. *Cat. giff.* 216. *R. Syn.* 85.

Muscus montanus, flagellis longis tenuibus cupressinis, ut plurimum indivisis. *Hist. ox.* III. 624. *f.* 15. *t.* 5. *f.* 7.

Anglis, Mountain Hypnum.

Habitat in arborum stipitibus et ericetis. In collibus cretaceis inter Northfleet et Gravesend in Cantio. *Dill.* III.

26. HYPNUM ramis vagis, foliis ovatis apice sub-squarrolatis undique reflexis. *Fl. suec.* 1727. *Sp. pl.* *sum.* 1127.

Hypnum repens, triangularibus reflexis foliis, majus. *R. Syn.* 82. *Dill. musc.* 303. *t.* 39. *f.* 38.

Muscus terrestris erectior, foliis reflexis. *Hist. ox.* III. 624. *f.* 15. *t.* 5. *f.* 2.

- β Hypnum repens, triangularibus reflexi foliis, minus. *R. Syn.* 82. *Dill. musc.* 304. *t.* 39. *f.* 39.

Anglis, common Hypnum.

Habitat in pratis humidis. IV.

27. HYPNUM furculis repentibus, ramis vagis erectis, foliis secundis, antheris subrotundis. *Sp. pl.* 1127.

Hypnum loreum montanum, capsulis subrotundis. *Dill. musc.* 305. *t.* 39. *f.* 40.

Hypnum repens, furculis magis erectis, foliis reflexis longioribus cinctis, operculo capituli magno. *R. Syn.* 82.

Muscus terrestris vulgaris. *Ger. Em.* 1370. *alter. Park.* 1306.

Muscus

Muscus repens major, foliis et flagellis longis et tenuibus donatus. *Hist. ox. III. 626. f. 15. t. 5. f. 24.*

Anglis, Ground Hypnum.

Habitat in montosis prope Glocestriam. D. Cole.
Dill.

***** *Surculis dendroidibus.*

dendroides 28. HYPNUM furculo erecto, ramis fasciculatis terminalibus simpliciusculis, antheris erectis. *Fl. suec. 1028. Sp. pl. 1128.*

Hypnum dendroides sericeum, setis et capsulis longioribus erectis. *Dill. musc. 313. t. 40. f. 48.*

Hypnum erectum arbusculam referens, ramulis subrotundis, confertim nascentibus. *R. Syn. 81.*

Muscus dendroides elatior, ramulis crebris minus furculosis capitulis pedunculis brevibus insidentibus. *Hist. ox. III. 626. f. 15. t. 5. f. 31.*

Anglis, Tree Hypnum.

Habitat in sylvis subhumidis. III.

alopecurum. 29. HYPNUM furculo erecto, ramis fasciculatis terminalibus subdivisis antheris nutantibus. *Sp. pl. 1128. Fl. suec. 1029.*

Hypnum dendroides obscurius, setis et capsulis brevioribus nutantibus. *Dill. musc. 315. t. 41. f. 49.*

Hypnum palustre erectum arbusculam referens, ramulis subrotundis. *Cat. giff. 220. R. Syn. 81.*

Muscus dendroides sylvarum erectus ramulis kali æmulis, radice repente. *R. Syn. ed. II. 32. Hist. ox. III. 626. f. 15. t. 5. f. 30.*

Anglis, Fox-tail Hypnum.

Habitat in sylvis et nemorosis humidiusculis. IV.

***** *Surculis teretiusculis.*

curtipendulum. 30. HYPNUM furculis vagis teretibus, foliis ovatis acutis patulis, antheris pendulis. *Fl. suec. 1030. Sp. pl. 1128.*

Hypnum dentatum curtipendulum, viticulis rigidis. *Dill. musc. 333. t. 43. f. 69.*

Hypnum

Hypnum arboreum repens, capsulis reflexis, brevibus pediculis insidentibus. *Cat. giff.* 220. *R. Syn.* 89.

Anglis, pendulous Hypnum.

Habitat ad radices arborum et in locis saxosis. II.

31. HYPNUM furculis pinnato-sparfis subulatis, foliis purum. ovatis obtusis conniventibus. *Fl. suec.* 1031. *Sp. pl.* 1128.

Hypnum ramis inæqualibus sparfis, foliis ovatis imbricatis, capsulis obliquis. *Fl. lapp.* 410.

Hypnum cupressiforme vulgare, foliis obtusis. *Dill. musc.* 309. t. 40. f. 45.

Hypnum terrestre erectum, ramulis teretibus, foliis inter rotunda et acuta medio modo se habentibus. *Cat. giff.* 220. *R. Syn.* 81.

- β Hypnum cupressiforme tenuius et compressius. *Dill. musc.* 312. t. 40. f. 47.

Hypnum longum erectum, foliis angustis caulibus ap-
pressis. *R. Syn.* 83.

Anglis, Meadow Hypnum.

Habitat in sylvis et pascuis. II.

32. HYPNUM furculis ramisque vagis teretibus erecti-illecebrum. usculis obtusis. *Sp. pl.* 1129. *Fl. suec.* 1032.

Hypnum cupressiforme rotundius vel Illecebræ æmulum. *Dill. musc.* 311. t. 40. f. 46.

Hypnum terrestre erectum, ramulis teretibus, foliis subrotundis albo-virentibus cinctis. *Cat. giff.* 220. *R. Syn.* 81.

Muscus terrestris, furculis kali geniculati aut illecebræ æmulis, foliolis subrotundis, squamatim incumben-
tibus. *R. Syn. ed II.* 37. *Hist. ox. III.* 626.

Anglis, geniculated Hypnum.

Habitat in ericetis. III, IV.

- 33 HYPNUM furculis teretibus ramosis, foliis acutis riparium. patulis distantibus. *Sp. pl.* 1129. *Fl. suec.* 1033.

Hypnum aquaticum, flagellis et teretibus et pennatis. *Dill. musc.* 308. t. 40. f. 44.

Hypnum ramosum fluitans pennatum. *R. Syn.* 81.

Anglis, floating Hypnum.

Habitat

Habitat in fluvio prope Hackney, et ad ripam Thamensis supra Londinum. Dill. XII. I.

- cuspidatum.* 34. HYPNUM furculis vagis: apice foliis convolutis acuminato. *Sp. pl.* 1129. *Fl. suec.* 1035.
 Hypnum palustre extremitatibus cuspidatis et pungentibus. *Dill. musc.* 300. *t.* 39. *f.* 34.
 Hypnum repens palustre, foliis triangularibus per caules expansis extremitatibus convolutis et acuminatis. *Cat. giff.* 219. *R. Syn.* 82.
Anglis, pointed Hypnum.
 Habitat in palustribus. III, IV.

***** *Surculis confertis.*

- sericeum.* 35. HYPNUM furculo repente, ramis confertis erectis, foliis subulatis, antheris erectis. *Sp. pl.* 1129. *Fl. lapp.* 412. *suec.* 1036.
 Hypnum vulgare sericeum recurvum, capsulis erectis cuspidatis. *Dill. musc.* 323. *t.* 42. *f.* 59.
 Hypnum repens trichoides terrestre, luteo-virens vulgare majus, capitulis erectis. *Cat. giff.* 215. *R. Syn.* 84.
 Muscus terrestris luteo-viridis, sericeus, repens. *R. Syn. ed. II.* 38. *Hist. ex. III.* 626. *f.* 15. *t.* 5. *f.* 25.
Anglis, filky Hypnum.
 Habitat in muris, truncis arborum et pascuis. XII, I.
- velutinum.* 36. HYPNUM furculo repente, ramis confertis erectis, foliis subulatis, antheris subnutantibus. *Fl. suec.* 1037. *Sp. pl.* 1129.
 Hypnum velutinum, capsulis ovatis cernuis. *Dill. musc.* 326. *t.* 42. *f.* 61.
 Hypnum repens trichoides terrestre viridius minus, capitulis tumidioribus cernuis. *Cat. giff.* 216. *R. Syn.* 84.
 a Hypnum repens trichoides terrestre minimum et breve, capitulis tumidioribus cernuis. *R. Syn.* 84.
Anglis, Velvet Hypnum.
 Habitat in umbrosis et ad sepes. II.

37. HYPNUM furculo repente, ramis confertis erectis, *palustre*.
foliis ovatis, antheris erectiusculis.
Hypnum heterophyllum aquaticum polycephalum, repens. *Dill. musc.* 293. *t.* 37. *f.* 27.
Hypnum repens filicifolium ramosum, ramulis surrectis et minus complanatis. *Cat. giff.* 218. *R. Syn.* 87.
Anglis, Marsh Hypnum.
Habitat *ad ripam* Thamesis *prope* Batterseam. I, II.
38. HYPNUM furculis repentibus, ramis filiformibus, *serpens*.
foliis oblitteratis. *Sp. pl.* 1130.
Hypnum trichoides serpens, setis et capsulis longis erectis. *Dill. musc.* 329. *t.* 42. *f.* 64.
Hypnum repens trichoides terrestre minimum, capitulis majusculis, oblongis erectis. *Cat. giff.* 216. *R. Syn.* 85.
Muscus terrestris omnium minimus, capitulis majusculis oblongis erectis. *Hist. ex. III.* 626. *f.* 15. *t.* 5. *f.* 14. 21.
Anglis, creeping Hypnum.
Habitat *in racemosis et ad radices arborum*. IV.
39. HYPNUM furculis erectis ramosis incurvatis. *Sp. sciuroides*.
pl. 1130.
Hypnum arboreum sciuroides. *Dill. musc.* 319. *t.* 41. *f.* 54.
Hypnum trichoides erectum, ramulis recurvis, obscuri coloris. *R. Syn.* 83.
Anglis, Squirrel-tail Hypnum.
Habitat *in arborum truncis*. II, IV.
40. HYPNUM furculis ramosissimis : ramis subulatis u- *myosuroi-*
trinque attenuatis teretibus. *Sp. pl.* 1130. *des.*
Hypnum myosuroides tenuius, capsulis nutantibus.
Dill. musc. 317. *t.* 41. *f.* 51.
Hypnum polyanthon, triangularibus angustis foliis.
R. Syn. 83.
Muscus trichoides humilis ramosus, capitulis oblongis tumidiusculis donatus. *Hist. ex. III.* 628. *f.* 15. *t.* 6. *f.* 3.

β Hyp-

- β *Hypnum myosuroides crassius*, capsulis erectis. *Dill. musc.* 316. t. 41. f. 50.
Hypnum repens triangularibus angustis foliis, ramulis subrotundis. *R. Syn.* 83.
 γ *Hypnum myosuroides brevius et crassius*, capsulis cernuis. *Dill. musc.* 318. t. 41. f. 52.
 δ *Hypnum myosuroides sericeum tenuius*, capsulis erectis. *Dill. musc.* 318. t. 41. f. 53.
Anglis, Mouse-tail Hypnum.
Habitat in sylvis locis saxosis et muris. II.
- ornithopodes.* 41. *HYPNUM furculis repentibus, ramis fasciculatis teretibus erectiusculis, antheris erectis ovatis.*
Hypnum gracile ornithopoides. *Dill. musc.* 320. t. 41. f. 55.
Anglis, Bird's-foot Hypnum.
Habitat in Fagis in saltu Enfieldienfi copiose; ad saxa per campos elatos Grey-Weathers dictos. S. Brewer. IV.
- clavellatum.* 42. *HYPNUM repens, ramulis erectis confertissimis, antheris incurvis, operculis inflexis.* *Sp. pl.* 1130.
Hypnum clavellatum parvum repens, setis brevibus. *Dill. musc.* 551. t. 81. f. 17.
Anglis, Club Hypnum.
Habitat in arborum truncis et lignis putridis passim. XII.

A L G Æ.

J U N G E R M A N N I A.

Masc. pedunculatus, nudus. *Anthera* quadrivalvis.
Fem. sessilis, nudus, feminibus subrotundis.

* *Frondibus pinnatis secundis.*

- asplenoides.* 1. *JUNGERMANNIA frondibus simpliciter pinnatis: foliolis ovatis subciliatis.* *Sp. pl.* 1131.
Lichenastrum asplenii facie, pinnis laxioribus. *Dill. musc.* 482. t. 69. f. 5.
 β *Lichenastrum asplenii facie, pinnis confertioribus.* *Dill. musc.* 483. t. 69. f. 6.

Liche-

Lichenastrum trichomanis facie, capitulis e foliorum summitate enascentibus, majus. *R. Syn.* 112.

Muscus trichomanoides, foliis rotundioribus pellucidis, squamatim conjuncte sibi incumbentibus. *Hist. ox.* III. 627. *f.* 15. *t.* 6. *f.* 42.

Anglis, Spleenwort: *Jungermannia*.

Habitat in sylvis et umbrosis humidis. II.

2. *JUNGERMANNIA* frondibus simpliciter pinnatis: *viticulosa*. foliolis subulatis. *Sp. pl.* 1131.

Jungermannia terrestris, viticulis longis, foliis perexiguis densissimis ex rotunditate acuminatis. *Mich. gen.* 8. *t.* 5. *f.* 4.

Lichenastrum trichomanis facie e basi et medio florens. *Dill. musc.* 484. *t.* 69. *f.* 7.

Lichenastrum capitulis nudis, trichomanis facie, foliolis densius congestis majus. *Cat. giff.* 211. *R. Syn.* 111.

Anglis, straggling *Jungermannia*.

Habitat in sylvis et umbrosis in comitatu Eboracensi, D. Richardson; circa Slingfoldiam in Suffexia. *Dill.* IV.

3. *JUNGERMANNIA* fronde pinnata simplici basi flo- *multiflora*. riera, foliolis acutis.

Lichenastrum multiflorum exile, foliis angustissimis. *Dill. musc.* 481. *t.* 69. *f.* 4.

Anglis, Many-flowered *Jungermannia*.

Habitat in sylvula collis Shooter's-hill dicti prope Londinum. *Dill. musc.* III.

4. *JUNGERMANNIA* frondibus simpliciter pinnatis: *polyanthos*. foliolis integerrimis imbricatis convexis. *Sp. pl.* 1131.

Jungermannia palustris repens, foliis densissimis ex rotunditate acuminatis. *Mich. gen.* 8. *t.* 5. *f.* 5.

Lichenastrum trichomanis facie, polyanthemum breve et repens. *Dill. musc.* 486. *t.* 70. *f.* 9.

Anglis, imbricated *Jungermannia*.

Habitat in palustribus; in Lead-hill in Surreja. IV.

5. JUN-

lanceolata. 5. JUNGERMANNIA frondibus simpliciter pinnatis lanceolatis apice floriferis, foliolis integerrimis. *Sp. pl.* 1131

Jungermannia palustris minima repens, foliis subrotundis densissimis læte virentibus. *Mich. ged.* 8. t. 5. f. 6.

Lichenastrum trichomanis facie minus, ab extremitate florens. *Dill. musc.* 486. t. 70. f. 10.

Lichenastrum capitulis nudis, trichomanis facie, foliolis densius congestis, minus. *Cat. giff.* 212. *R. Syn.* 112.

Anglis, Spear-leaved Jungermannia.

Habitat in umbris humidis. II.

bidentata. 6. JUNGERMANNIA frondibus simpliciter pinnatis apice floriferis, foliolis bidentatis. *Sp. pl.* 1132.

Jungermannia major repens, foliis bifidis. *Mich. gen.* 8. t. 5. f. 12.

Lichenastrum pinnulis acutioribus et concavis, bifidis, majus. *Dill. musc.* 487. t. 70. f. 11.

Lichenastrum trichomanis facie, foliolis bifidis, majus. *R. Syn.* 113.

Muscus lichenoides, foliis pinnatis bifidis, minor. *R. Syn. Hist. ox.* III. 627. f. 15. t. 6. f. 47.

Anglis, Bifid Jungermannia.

Habitat in sylvis et ericetis humidiusculis. X, XI.

bicuspidata. 7. JUNGERMANNIA frondibus simpliciter pinnatis medio floriferis, foliolis bidentatis. *Sp. pl.* 1132.

Jungermannia minima repens, foliis bifidis, vagina florum cylindracea. *Mich. gen.* 9. t. 6. f. 17.

Lichenastrum pinnulis acutissime bifidis, minimum. *Dill. musc.* 488. t. 70. f. 13.

Lichenastrum trichomanis facie, foliolis bifidis. *Cat. giff.* 213. *R. Syn.* 113

Anglis, forked Jungermannia

Habitat in umbris humidis. III.

** Frondibus

8. JUNGERMANNIA fronde pinnata ramosa apice *quinque-*
 florifera, foliolis apice *quinqüedentatis*.

Lichenastrum multifidum majus, ab extremitate flo-
 rens. *Dill. musc.* 494. t. 71. f. 23.

Lichenastrum trichomanis facie, foliolis multifidis, ca-
 pitulis e summis ramulis nascentibus. *R. Syn.* 113.

Anglis, Toothed Lichenastrum.

Habitat in comitatu Eboracensi. *D. Richardson.* IV.

** *Frondibus pinnatis: pinnulis auriculatis superne*
tectis.

9. JUNGERMANNIA frondibus supra bipinnatis apice *undulato*,
 floriferis, foliolis subrotundis integerrimis. *Sp. pl.*
 1131.

Lichenastrum pinnulis auriculatis majoribus et non cre-
 natis. *Dill. musc.* 490. t. 71. f. 17.

Anglis, Eared Jungermannia.

Habitat in umbrosis. III.

10. JUNGERMANNIA frondibus supra bipinnatis a- *nemorea*,
 pice floriferis, foliolis ciliatis. *Sp. pl.* 1132.

Lichenastrum auriculatum; pinnis minoribus crenatis.

Dill. musc. 490. t. 71. f. 18.

Anglis, Wood Lichenastrum.

Habitat in sylvis udis in comitatu Westmorlandico.
 III.

11. JUNGERMANNIA frondibus supra bipinnatis in- *resupinata*
 ferne floriferis: foliolis crenulatis imbricatis rotun-
 dis. *Sp. pl.* 1132. *Fl. suec.* 1038.

Jungermannia furculosa resupinata, foliolis pinnato-
 imbricatis deflexis rotundis integris: auricula late-
 rali. *Fl. suec. ed. I.* 926.

Lichenastrum auriculatum, pinnulis rotundis crispum.

Dill. musc. 491. t. 71. f. 19.

Lichenastrum trichomanis facie, capitulis foliorum
 summitate enascentibus, medium. *R. Syn.* 112.

Anglis, curled Jungermannia.

Habitat in fissuris rupium. IV.

- albicans.* 12. JUNGERMANNIA frondibus supra bipinnatis apice floriferis: foliolis linearibus recurvatis. *Sp. pl.* 1133. *Fl. suec.* 1039.
 Lichenastrum auriculatum, pinnulis angustis, planis, recurvis. *Dill. musc.* 492. *t.* 71. *f.* 20.
 Lichenastrum trichomanis facie, capitulis e foliorum summitate enascentibus minus. *R. Syn.* 112.
Anglis, white Jungermannia.
 Habitat in *sylvis humidis*. III.
- trilobata.* 13. JUNGERMANNIA frondibus subtus bipinnatis, foliolis quadratis obsolete trilobis. *Sp. pl.* 1133. *Fl. suec.* 1040.
 Jungermannia foliis latiusculis obtusis undulatis et veluti angulosis. *Mich. gen.* 7. *t.* 5. *f.* 10.
 Lichenastrum pinnulis obtuse trifidis nervo geniculato. *Dill. musc.* 493. *t.* 71. *f.* 22.
Anglis, Trifid Jungermannia.
 Habitat in *Cambria*. *Dill. musc.*
- reptans.* 14. JUNGERMANNIA frondibus subtus bipinnatis apice reptantibus, foliolis quadridentatis. *Sp. pl.* 1133.
 Lichenastrum multifidum exiguum ad basin florens per siccitatem imbricatum. *Dill. musc.* 494. *t.* 71. *f.* 24.
 Lichenastrum trichomanis facie, foliolis multifidis, capitulis ex imis cauliculis nascentibus. *R. Syn.* 113.
Anglis, creeping Jungermannia.
 Habitat in *umbrosis udis*. IV.

*** *Frondibus imbricatis.*

- complanata.* 15. JUNGERMANNIA furculis repentibus: foliolis inferne auriculatis duplicato-imbricatis, ramis æqualibus. *Sp. pl.* 1133. *Fl. suec.* 1041.
 Lichenastrum imbricatum majus, squamis compressis et planis. *Dill. musc.* 496. *t.* 72. *f.* 26.
 Lichenastrum imbricatum majus. *R. Syn.* 111.
Anglis, Flat Jungermannia.
 Habitat ad *arborum truncos frequens*. IV.

16. JUNGERMANNIA furculis repentibus, foliolis *dilatata*, inferne auriculatis duplicato-imbricatis, ramis apice latioribus. *Sp. pl.* 1133. *Fl. suec.* 1042.
 Lichenastrum imbricatum minus, squamis convexo-concavis. *Dill. musc.* 497. t. 72. f. 27.
 Lichenastrum imbricatum, minus. *R. Syn.* 111.
Anglis, scaly Jungermannia.
 Habitat ad truncos arborum. III, IV.
17. JUNGERMANNIA foliis imbricatis serie duplici: *tamariscifolia* superioribus subrotundis convexis obtusis quadruplo *folia* majoribus. *Sp. pl.* 1134. *Fl. suec.* 1043.
 Lichenastrum imbricatum tamarisci narbonensis facie. *Dill. musc.* 499. t. 72. f. 31.
Anglis, Tamarisk Jungermannia.
 Habitat in rupibus et truncis arborum comitatu Eboracensi *passim*. III.
18. JUNGERMANNIA furculis procumbentibus sub-*platyphylla* tus imbricatis, foliolis cordatis acutis. *Sp. pl.* 1134. *la*.
 Lichenastrum arboris vitæ facie, foliis minus rotundis. *Dill. musc.* 501. t. 72. f. 32.
 Muscus muralis floridus, foliis subrotundis creberrime imbricatis dispositis. *Hist. ox.* III. 627. f. 15. t. 6. f. 44.
Anglis, Wall Jungermannia.
 Habitat ad truncos arborum et muros. III, IV.
19. JUNGERMANNIA furculis repentibus, foliolis *ciliaris* duplicato-imbricatis, inferne auriculatis ciliatis.
Sp. pl. 1134. *Fl. suec.* 1044.
 Lichenastrum filicinum pulchrum villosum. *Dill. musc.* 503. t. 73. f. 35.
 Lichenastrum filicinum crispum. *R. Syn.* 111.
Anglis, Fern Jungermannia.
 Habitat in ericetis udis et ad rivulos; in comitatu Eboracensi, D. Richardson. Found by Mr. Dandridge in a small Current of Water, which runs through a Wood called Old-fall, between Highgate and Musclevill. *R. Syn.* IV.

- varia.* 20. JUNGERMANNIA furculis erectiusculis bifariam imbricatis, foliolis bipartitis. *Sp. pl.* 1135.
 Lichenastrum foliis variis. *R. Syn.* 113. *Dill. musc.* 505. *t.* 73. *f.* 36.
Anglis, upright Jungermannia.
 Habitat in sylvis et umbrosis humidis. III.

**** Surculis undique imbricatis foliolis sparsis.

- julacea.* 21. JUNGERMANNIA furculis teretibus, foliolis undique imbricatis, floribus pedunculatis. *Sp. pl.* 1135.
 Lichenastrum alpinum Bryi juliacei argentei facie, *Dill. musc.* 506. *t.* 73. *f.* 38.
Anglis, round Jungermannia.
 Habitat in rupibus udis, in montibus Snowdon et Glyder. *Dill.* IX.

- rufesbris.* 22. JUNGERMANNIA furculis teretibus, foliolis subulatis secundis. *Sp. pl.* 1135. *Fl. suec.* 1045.
 Lichenastrum alpinum nigricans, foliis capillaceis reflexis. *Dill. musc.* 507. *t.* 73. *f.* 40.
Anglis, Rock Jungermannia.
 Habitat in rupibus humentibus in montibus Snowdon et Glyder in Arvoniam. *Dill.* IV. V.

- trichophylla.* 23. JUNGERMANNIA furculis teretibus foliolis capillaceis æqualibus. *Sp. pl.* 1135. *Fl. suec.* 1046.
 Lichenastrum trichoides minimum, in extremitate florens. *Dill. musc.* 505. *t.* 73. *f.* 37.
Anglis, hairy Jungermannia.
 Habitat in ericetis turfosis prope North Bierly in comitatu Eboracensi, *D. Richardson.*

- alpina.* 24. JUNGERMANNIA furculis teretibus, foliolis ovatis patentibus, calycibus imbricatis. *Sp. pl.* 1135.
 Lichenastrum alpinum atro-rubens teres, calycibus squamosis. *Dill. musc.* 506. *t.* 73. *f.* 39.
Anglis, Mountain Jungermannia.
 Habitat in palustribus montosis; in montibus Snowdon Llanberis et Cader Idæris. *Dill.* IX. X.

***** Acaule

***** *Acaules frondibus simplicibus.*

25. JUNGERMANNIA acaulis, foliolis frondi innato. *epiphylla.*
Sp. pl. 1135. *Fl. suec.* 1047.
 Lichenastrum capitulis rotundis, e foliorum medio
 enascentibus. *R. Syn.* 110. *Dill. musc.* 508. t.
 74. f. 41.
 Lichen petræus, cauliculo calceato. *Bauh. pin.* 362.
 Lichen alter minor, folio calceato. *Park.* 1314.
Anglis, broad-leaved Jungermannia.
 Habitat in umbrosis udis et ad rivulos. III.
26. JUNGERMANNIA acaulis, fronde oblonga sinu- *pinguis.*
 ata pingui. *Sp. pl.* 1136. *Fl. suec.* 1048.
 Lichenastrum capitulis oblongis, juxta foliorum divi-
 suras enascentibus. *R. Syn.* 110. *Dill. musc.* 509.
 t. 74. f. 42.
Anglis, jagged Jungermannia.
 Habitat in paludibus. IV.
27. JUNGERMANNIA acaulis, fronde bipinnatifida. *multifida.*
Sp. pl. 1136. *Fl. suec.* 1049.
 Lichenastrum ambrosiæ divisura. *R. Syn.* 111. *Dill.*
musc. 511. t. 74. f. 43.
Anglis, dwarf Jungermannia.
 Habitat in sylvis et umbrosis humidis. In sylvula Cane-
 wood dicta prope Hampstedium. *D. Doody.* prope
 Charlton in Cantio. *Dill.* IV.
28. JUNGERMANNIA acaulis, fronde lineari ramo- *furcata.*
 fa : extremitatibus furcatis obtusiusculis. *Sp. pl.*
 1136.
 Lichenastrum tenuifolium furcatum, thecis globosis
 pilosis. *Dill. musc.* 512. t. 74. f. 45.
 Ulva saxatilis furcata, latiusculis et tenerioribus seg-
 mentis. *R. Syn.* 63.
Anglis, Globe Jungermannia.
 Habitat in sylvis. IV.
29. JUNGERMANNIA acaulis, fronde subpinnatif- *pusilla.*
 da, lobis imbricatis, perichætio plicato. *Sp. pl.*
 1136. *Fl. suec.* 1080.

CRYPTOGAMIA ALGÆ.

Lichenastrum exiguum, capitulis nigris lucidis e cotylis parvis nascentibus. *Dill. musc.* 513. t. 74. f. 46.

Anglis, shining *Jungermannia*.

Habitat in ericetis; in ericetis prope Woolwich. *Dill.* III, IV.

TARGIONIA. 1060.

Cal. bivalvis, includens globum.

hypophylla 1. TARGIONIA. *Sp. pl.* 1136.

Targionia minima et vulgaris. *Mich. gen.* 3. t. 3.

Lichen petræus minimus, fructu orobi. *Baub. pin.* 362. *Dill. musc.* 532. t. 78. f. 9.

Lichenastrum capitulo oroboide. *R. Syn.* 110.

Lichen petræus minimus acaulos. *Park.* 1315.

Anglis, Vetch *Targionia*.

Habitat in locis muscosis in comitatu Eboracensi. D. Richardson.

MARCHANTIA. 1061.

Masc. Cal. peltatus, subtus tectus. *Corollis* monopetalis. *Antheræ* multifidæ.

Fem. Cal. sessilis, campanulatis, polyspermus.

polymorpha 1. MARCHANTIA calyce communi decemifido. *Fl. lapp.* 422. *suæc.* 1051. *Sp. pl.* 1137. *Roy. lugdb.* 507.

Marchantia major, capitulo stellato: radiis teretibus.

Mich. gen. 2. t. 1. f. 1

Lichen fontanus major stellatus æque ac umbellatus et cyathophorus. *Dill. musc.* 523. t. 76. f. 6.

Lichen petræus latifolius, sive Hepatica fontana. *Baub. pin.* 362. *R. Syn.* 115.

2 *Marchantia capitulo stellato, radiis teretibus.* *Mich. gen.* 2. t. 1. f. 2.

Lichen domesticus minor stellatus æque ac umbellatus et cyathophorus. *Dill. musc.* 527. t. 77. f. 7.

Lichen petræus stellatus. *Baub. pin.* 362. *R. Syn.* 115.

Anglis, common *Marchantia*.

Habitat

Habitat *a juxta aquas locis umbrosis et ad fontes; β in umbrosis areis et ambulacris hortorum.* VIII.

2. MARCHANTIA calyce communi quadripartito: *cruciata*.
 laciniis tubulosis. *Hort. cliff.* 477. *Sp. pl.* 1137.
 Lichen semifer lunulatus, florifer pileatus tandem
 cruciatus. *Dill. musc.* 521. *t.* 75. *f.* 5.
 Lichen pileatus parvus, capitulo crucis instar se expan-
 dente. *R. Syn.* 115.
 Lunaria vulgaris. *Mich. gen.* 4. *t.* 4.
Anglis, cross-headed Marchantia.
 Habitat *in umbrosis.* VII, VIII.
3. MARCHANTIA calyce communi quinquefido he- *hemisphæ-*
misphærico, perichætio nullo. *Fl. suec.* 1052. *rica.*
lapp. 424. *Sp. pl.* 1138.
 Lichen pileatus parvus, foliis crenatis. *R. Syn.* 114.
Dill. musc. 519. *t.* 75. *f.* 2.
 Hepatica media, capitulo hemisphærico. *Mich. gen.*
 3. *t.* 2. *f.* 2.
Anglis, Marsh Marchantia.
 Habitat *in paludosis; in a small Water-course, near*
Dulwich-wells, Mr. Dandridge. *R. Syn.* IV, V.
4. MARCHANTIA calyce communi quinque loculari *conica.*
subovato. *Sp. pl.* 1138.
 Lichen vulgaris major pileatus et verrucosus. *Dill.*
musc. 516. *t.* 75. *f.* 1.
 Lichen petræus pileatus. *Park.* 1314. *R. Syn.* 114.
 Lichen petræus cauliculo pileolum sustinente. *Bauh.*
pin. 362.
Anglis, Wart Marchantia.
 Habitat *in umbrosis et ad rivulos.* IV.

B L A S I A. 1062.

Masc. *Cal.* cylindricus granis repletus.

Fem. *Cal.* nudus. *Fructu* subrotundo, foliis immer-
 so, polyspermo.

1. BLASIA. *Fl. suec.* 1053. *Sp. pl.* 1138.
 Blasia pusilla, lichenis pyxidati facie. *Mich. gen.* 14. *pusilla.*
t. 7.

Mnium lichenis facie. *Dill. musc.* 237. t. 31. f. 7.
Anglis, Dwarf Blasia.
 Habitat *prope* Manchester. *Dill. musc.*

R I C C I A. 1063.

Masc. nudus. *Antheræ* subulata truncata.
 Fem. nudus. *Caps.* globosa, unilocularis, polysperma.

minima. 1. RICCIA frondibus glabris bipartitis acutis. *Fl. suec.* 1054. *Sp. pl.* 1139.

Riccia minima nitida, segmentis angustioribus acutis.
Mich. gen. 107. t. 57. f. 6.

Lichen omnium minimus, foliolis fissis super terram expansis. *Cat. giff.* 210. *R. Syn.* 115. *Dill. musc.* 534. t. 78. f. 11.

Anglis, small Riccia.

Habitat *in inundatis* ; *in ericeto* Blackheath *dicto.* *R. Syn.* XI.

fluitans. 2. RICCIA frondibus dichotomis lineari-filiformibus.
Sp. pl. 1139.

Lichenastrum aquaticum fluitans tenuifolium furcatum. *Dill. musc.* 514. t. 74. f. 47.

Ulva palustris furcata, angustioribus et firmioribus segmentis. *R. Syn.* 63.

Anglis, Floate Riccia.

Habitat *in aquis stagnantibus.* *In a ditch near* Deptford Dock, Mr. Petiver. *R. Syn.* v. *Barnes con*

innatans. 3. RICCIA frondibus obcordatis ciliatis. *Syst. nat.* 1339.

Lichen parvus vernus cordiformis, ima parte fimbriatus, Lentis palustris modo aquæ innatans. *Buddl. hort. ficc. Vol. II.* 15. *R. Syn.* 116.

Anglis, fringed Riccia.

Habitat *in aquis stagnantibus prope* Henley. *R. Syn.*

glauca. 4. RICCIA frondibus glabris canaliculatis bilobis obtusis. *Sp. pl.* 1139.

Riccia minima pinguis, foliis latiusculis ample sulcatis e glauco virefcentibus. *Mich. gen.* 107. t. 57. f. 4.

Lichen minimus, foliis venosis bifariam vel trifariam se dividendo progredientibus. *R. Syn. ed. II.* 41. *Dill. musc.* 533. t. 78. f. 10.

Lichen f. *Hepatica* foliis crassis, *Rutæ murariæ* aut *Chamædryos* foliis laciniatis quodammodo similibus. *R. Hist. III* 48.

Anglis, Marsh *Riccia*.

Habitat in paludibus; in ericeto Woolwicensi. *Dill. musc.* III.

5. *RICCIA* frondibus ramosis, laciniis sinuatis apice di-sinuata. latatis crenatis.

Lichenoides gelatinosum, foliis latioribus tuniformibus. *Dill. musc.* 142. t. 19. f. 29.

Anglis, Rock *Riccia*.

Habitat ad rupes prope Wigmore in Herefordienfi comitatu. L. Brown. *Dill.*

ANTHOCEROS. 1064.

Masc. *Cal.* sessilis, cylindricus, integer. *Anth.* subulata longissima, trivalvis.

Fem. *Cal.* sexpartitus. *Sem.* tria.

1. *ANTHOCEROS* frondibus indivisis sinuatis puncta-punctatus. tis. *Sp. pl.* 1139.

Anthoceros foliis minoribus magis laciniatis. *Dill. musc.* 476. t. 68. f. 1.

Lichenastrum gramineo pediculo et capitulo, oblongo, bifurco. *R. Syn.* 109.

Anglis, spotted *Anthoceros*.

Habitat in umbrosis sed rarius. Prope Farnhurst in Sufflexia; in ericeto Woolwicensi, &c. *Dill. musc.* IV.

LICHEN. 1065.

Masc. *Receptaculum* subrotundum, planiusculum, nitidum.

Fem. *Farina* foliis adspersa.

* *Leprosi*

* *Leprosi tuberculati.*

- scriptus.* 1. LICHEN leprofus albicans, lineolis nigris ramosis characteriformibus. *Fl. suec.* 1057. *Sp. pl.* 1140.
Lichenoides crusta tenuissima, peregrinis velut litteris inscripta. *R. Syn.* 71. *Dill. musc.* 125. t. 18. f. 1.
Anglis, Lettered Liverwort.
Habitat in corticibus arborum frequens.
- geographicus.* 2. LICHEN leprofus flavescens lineolis nigris mappam referens. *Fl. suec.* 1058. *Sp. pl.* 1140.
Lichenoides nigro-flavum tabulæ geographicæ instar pictum. *Dill. musc.* 126. t. 18. f. 5.
Anglis, Mapp Liverwort.
Habitat in rupibus calcaris in comitatibus Lancastriensi, Eboracensi et Cumberlandico.
- rugosus.* 3. LICHEN leprofus albicans, lineolis simplicibus punctisque nigris confertis. *Sp. pl.* 1140. *Fl. suec.* 1059
Lichenoides punctatum et rugosum nigrum. *Dill. musc.* 125. t. 18. f. 2.
Anglis, wrinkled Liverwort.
Habitat in corticibus arborum.
- sanguineus.* 4. LICHEN leprofus cinereo-virescens, tuberculis atris. *Fl. suec.* 1060. *Sp. pl.* 1140.
Lichenoides leprosum, crusta cinereo-virescente, tuberculis nigerrimis. *Cat. giss.* 205. *R. Syn.* 71.
Dill. musc. 126. t. 18. f. 3.
Anglis, black nobbed Liverwort.
Habitat in rupibus et truncis arborum.
- fusco-ater.* 5. LICHEN leprofus fuscus tuberculis atris. *Sp. pl.* 1140. *Fl. suec.* 1063.
Anglis, Wall Liverwort.
Habitat in muris antiquis passim.
- calcareus.* 6. LICHEN leprofus candidus, tuberculis atris. *Fl. suec.* 1061. *Sp. pl.* 1140.

Lichenoides tartareum tinctorium candidum, tuberculis atris. Dill. musc. 128. t. 18. f. 8.

Anglis, Dyer's Liverwort.

Habitat in rupibus calcareis in comitatu Westmorlandico; in monte Glyder in Arvoniam. Dill.

7. LICHEN leprosus albus, tuberculis albis farinaceis. *fagineus. Sp. pl. 1141. Fl. suec. 1066.*

Anglis, white Liverwort.

Habitat in carpini et ulmi truncis et ramis.

8. LICHEN lichen leprosus cinerius tuberculis albidis *carpinus. rugosus. Sp. pl. 1141. Fl. suec. 1067.*

Lichen geographicus. It. scan. 48.

Anglis, Horn-beam Liverwort.

Habitat in carpini, corylli et Fraxini, truncis.

9. LICHEN leprosus candidus, tuberculis incarnatis. *ericetorum. Fl. suec. 1068. Sp. pl. 1141.*

Lichenoides fungiforme, crusta leprosa candida, capitulis et pediculis incarnatis. R. Syn. 70.

Coralloides fungiforme carneum, basi leprosa. Dill. musc. 76. t. 14. f. 1.

Anglis, Heath Liverwort.

Habitat in ericetis udis in comitatu Eboracensi. D. Richardson. X—IV.

10. LICHEN leprosus fuscus, tuberculis pedunculatis *rufus. planis subfuscis.*

Coralloides fungiforme, saxatile, pallide fuscum. Dill. musc. 78. t. 14. f. 4.

Anglis, brown Liverwort.

Habitat in muris antiquis. XII—II.

11. LICHEN leprosus virescens, tuberculis flavo-ru-
bescentibus. *flavo-rubescens.*

Anglis. Orange Liverwort.

Habitat in truncis arborum inter Hampstead et Highgate.

- ferrugineo-* 12. LICHEN leprosus albicans, tuberculis rufis.
♂s. Lichenoides leprosum, tuberculis fuscis et ferrugineis.
Dill. musc. 126. t. 18. f. 4. t. 55. f. 8.
Anglis, rusty Liverwort.
Habitat in corticibus arborum.

** *Leprosi scutellati.*

- candelarius.* 13. LICHEN crustaceus flavus, scutellis luteis. *Sp. pl.*
 1141. *Fl. suec.* 1069.
Byssus farinacea flava. It. oel. 30. *W-goth.* 159.
Anglis, yellow Liverwort.
Habitat in corticibus arborum et muris. XII—VI.

- tartareus.* 14. LICHEN crustaceus albidus, scutellis luteis : margine albo. *Sp. pl.* 1141. *Fl. suec.* 1070.
 Lichenoides crustaceum et leprosum acetabulis majoribus luteis, limbis argenteis. *R. Syn.* 71. *Dill. musc.* 132. t. 18. f. 13.
Anglis, Welsh Liverwort.
Habitat in rupibus ; in Arvoniam, D. Lhwyd. In comitatu Eboracensi. D. Richardson.

- pallescens.* 15. LICHEN crustaceus albicans, scutellis pallidis. *Fl. suec.* 1071. *Sp. pl.* 1142.
 Lichenoides crustosum orbiculare incanum. *Dill. musc.* 135. t. 18. f. 17.
 Lichenoides cinereum mere crustaceum eleganter expansum. *R. Syn.* 71.
 Lichenoides crustaceum et leprosum, scutellis cinereo-virescenti. *R. Syn.* 71.
Anglis, pale Liverwort.
Habitat in muris, et arborum truncis.

- subfuscus.* 16. LICHEN crustaceus albicans, scutellis subfuscis : margine cinereo subcrenato. *Fl. suec.* 1072. *Sp. pl.* 1142.
 Lichenoides crustaceum et leprosum, scutellis subfuscis. *R. Syn.* 71. *Dill. musc.* 134. t. 18. f. 16.
Anglis, brownish Liverwort.
Habitat in corticibus arborum et muris.

17. LICHEN crustaceus cinereus, scutellis planis albis *albescens*.
 farinaceis.
 Lichen crustaceus albescens, scutis farinaceis. *Pet.*
Vaill. Parif. 116.
 Lichenoides candidum et farinaceum, scutellis fere
 planis. *Dill. musc.* 131. t. 18. f. 11.
 Lichenoides crusta tenuissima, sulcis cochleæ formi-
 bus insignita. *R. Syn.* 71.
Anglis, mealy Liverwort.
 Habitat in muris et corticibus arborum.
18. LICHEN crustaceus cinereus, scutellis virescentibus. *veruco-*
brum. *Dill. musc.* 128. t. 18. f. 9.
Anglis, Warty Liverwort.
 Habitat in corticibus arborum et muris.
19. LICHEN crustaceus albicans, scutellis rugosis ni-
 gris.
 Lichenoides crustaceum et leprosum, scutellis nigri-
 cantibus majoribus et minoribus. *Cat. giff.* 205.
R. Syn. 71. *Dill. musc.* 133. t. 18. f. 15.
Anglis, black cupped Liverwort.
 Habitat in arboribus et muris frequens.
20. LICHEN crustaceus luteo-virescens, scutellis flavis. *flavescent.*
 Lichenoides crustosum, orbiculis et scutellis flavis.
Dill. musc. 136. t. 18. f. 18.
Anglis, yellowish Liverwort.
 Habitat in muris.

*** *Imbricati.*

21. LICHEN imbricatus, foliolis multifidis obsolete læ-centrifu-
 vibus albidis centrifugis, scutellis rufo-fuscis. *Sp. gus.*
pl. 1142. *Fl. suec.* 1074.
 Lichenoides imbricatum viridans, scutellis badis. *Dill.*
musc. 180. t. 24. f. 75.
Anglis, spreading Liverwort.
 Habitat supra saxis grandioribus prope Dolgelle in
 Mervinia et circa Llanberis in Arvoniam. *Dill.*

- saxatilis.* 22. LICHEN imbricatus, foliolis sinuatis scabris lacunosis, scutellis folio concoloribus. *Sp. pl.* 1142. *Fl. suec.* 1075.
 Lichen opere phrygio ornatus. *Vaill. parif. t.* 21. *f.* 5.
 Lichenoides vulgatissimum cinereo-glaucum lacunosum et cirrosum. *Dill. musc.* 188. *t.* 24. *f.* 83.
 Lichenoides crusta foliosa, superne cinereo-glaucæ, inferne nigra et cirrosa, scutellis nigricantibus. *Cat. giff.* 206. *R. Syn.* 72.
 β Lichenoides arboreum purpurascens tenue et læve, scutellis majoribus. *R. Syn.* 73.
Anglis, Stone Liverwort.
Habitat in saxis et truncis arborum frequens.
- omphalodes.* 23. LICHEN imbricatus, foliolis multifidis glabris obtusis incanis, punctis vagis eminentibus. *Sp. pl.* 1143. *Fl. suec.* 1076.
 Lichen nigricans omphalodes. *Vaill. parif. t.* 20. *f.* 10.
 Lichen petræus purpureus derbienfis. *Park.* 1315.
 Lichenoides saxatile tinctorium, foliis pilosis purpureis. *R. Syn.* 74. *Dill. musc.* 185. *t.* 24. *f.* 80.
Anglis, purple Liverwort.
Habitat in monte Steiperstone in Salopiensi comitatu et in plerisque rupibus Cambriæ. Dill.
- olivaceus.* 24. LICHEN imbricatus, foliolis lobatis nitidis lividis. *Sp. pl.* 1143. *Fl. suec.* 1077.
 Lichenoides olivaceum, scutellis amplioribus verrucosis. *Dill. musc.* 184. *t.* 24. *f.* 78.
 β Lichenoides olivaceum, scutellis lævibus. *Dill. musc.* *t.* 24. *f.* 77.
 Lichenoides crusta foliosa scutellata, pullum. *Cat. giff.* 206. *R. Syn.* 72.
 γ Lichenoides arboreum et saxatile, crusta foliosa tenui, fusco-virente, in segmenta latiora plerumque divisa. *R. Syn.* 74.
Anglis, Olive Liverwort.
Habitat in rupibus et arboribus; in syvula prope Shooters-Hill, in Cantio. Dill.

25. LICHEN imbricatus, dentato-ciliatus, scutellis fo- *crifatus*.
lio majoribus. *Sp. pl.* 1143.
Lichenoides gelatinosum, foliis imbricatis et *crifatis*.
Dill. musc. 140. t. 19. f. 26.
Lichenoides gelatinosum tenerius laciniatum, ex plum-
beo colore cærulescens, scutellis fuscis. *R. Syn.*
72.
β Lichenoides foliis pilosis crassioribus, obscure virenti-
bus, scutellis fuscis. *R. Syn.* 74.
Anglis, crested Liverwort.
Habitat in rupibus et muris; circa Ludloam in Salopia.
Littl. Brown, Dill.
26. LICHEN imbricatus, foliis lobatis truncatis crena- *crispus*.
tis atro-viridibus, scutellis folio concoloribus.
Lichenoides gelatinosum atro-virens, crispum et ru-
gosum *Dill. musc.* 139. t. 19. f. 23.
Anglis, curled Lichen.
Habitat in locis umbrosis saxosis et ad murorum bases.
27. LICHEN imbricatus, foliolis crispis fulvis, peltis *parieti-*
concoloribus fulvis. *Sp. pl.* 1143. *Fl. suec.* 1080. *nus*.
Lichenoides vulgare sinuosum, foliis et scutellis luteis.
Dill. musc. 180. t. 24. f. 76.
Lichenoides crusta foliosa scutellata, flavescens. *Cat.*
giff. 206. *R. Syn.* 72.
Anglis, common Liverwort.
Habitat in corticibus arborum et muris.
28. LICHEN imbricatus: laciniis obtusis subinflatis. *physodes*.
Sp. pl. 1144. *Fl. suec.* 1081.
Lichen arboreus albescens, segmentis cornigeris, sub-
tus aterrimus. *Pet. Gaz.* t. 14. f. 6.
Lichenoides ceratophyllon obtusius et minus ramo-
sum. *R. Syn.* 76. *Dill. musc.* 154. t. 20. f. 49.
Anglis, inflated Liverwort.
Habitat in ericetis in comitatibus Eboracensi et West-
morlandico passim.

stellaris. 29. LICHEN imbricatus: foliolis oblongis laciniatis angustis cinereis, scutellis pullis. *Sp. pl.* 1144. *Fl. suec.* 1082.

Lichenoides cinereum, segmentis argutis stellatis, scutellis nigris. *Dill. musc.* 176. *t.* 24. *f.* 70.

Lichenoides arboreum, crusta foliosa albo-cinerea tenuiter et eleganter dissecta, scutellis nigris. *R. Syn.* 74.

Anglis, starry Liverwort.

Habitat in corticibus Fraxini, et Pruni sylvestris frequens.

**** *Foliacei.*

ciliaris. 30. LICHEN foliaceus erectiusculus: laciniis linearibus ciliatis, scutellis pedunculatis crenatis. *Sp. pl.* 1144. *Fl. suec.* 1085.

Lichenoides hispidum majus et rigidius, scutellis nigris. *Dill. musc.* 150. *t.* 20. *f.* 45.

Lichenoides arboreum foliosum cinereum, scutellis nigris foliorum extremitatibus hispidis et pilosis. *Cat. giff.* 207. *R. Syn.* 73.

Musco fungus arboreus cinereus scutellatus, marginibus pilosis. *Hist. ox. III.* 634. *f.* 15. *t.* 7. *f.* 6.

Anglis, Hairy Liverwort.

Habitat in arborum truncis frequens.

perlatus. 31. LICHEN foliaceus repens lobatus glaber, subtus atris, scutellis pedunculatis integris.

Lichenoides glaucum perlatus, subtus nigrum et cirrosum. *Dill. musc.* 147. *t.* 20. *f.* 39.

Musco-fungus lichenoides arborum, angustior, foliis superne cinereis, subtus nigricantibus villosis. *Hist. ox. III.* 634.

Anglis, Pearl Liverwort.

Habitat in truncis arborum passim.

islandicus. 32. LICHEN foliaceus adscendens laciniatus; marginibus elevatis ciliatis. *Sp. pl.* 1145. *Fl. suec.* 1085.

Lichen terrestris, foliis eryngii, Buxb. cent. II. *t.* 6. *f.* 1. 2.

Lichenoides

32. This is used by the Icelanders for food.

- Lichenoides rigidum, eryngii folia referens. *R. Syn.*
 77. *Dill. musc.* 209. *t.* 28. *f.* 111.
- β Lichenoides Eryngii folia referens, tenuioribus et crispioribus foliis. *Gelf. upf.* 31. *Dill. musc.* 212. *t.* 28. *f.* 112.
- γ Coralloides tenuissimum nigricans, mundi muliebris instar textum. *Dill. musc.* 113. *t.* 17. *f.* 32.
Anglis, Eryngo-leaved Liverwort.
 Habitat in ericetis et montosis; in montibus Walliæ.
Dill; in comitatibus Eboracensi et Westmorlandico; frequens.
33. LICHEN foliaceus repens lobatus glaber, lobis sinuatis, scutellis fulvis.
 Lichen pulmonarius arboribus adnascens maximus, inferne obscurus, desuper e glauco-cinereus, receptaculis florum rubris, seminibus nigricantibus. *Mich. gen.* 88. *t.* 46.
 Lichenoides subglaucum cumatile, foliis tenacibus, eleganter laciniatis. *Dill. musc.* 197. *t.* 26. *f.* 99.
Anglis, jagged Liverwort.
 Habitat secus viam inter Carno et Main-Lloÿne in Mervinia. *Dill.*
34. LICHEN foliaceus laciniatus obtusus glaber: supra pulmona-lacunofus; subtus tomentosus. *Sp pl.* 1145. *Fl. riis. suec.* 1087.
 Lichen arborum. *Ger. Em.* 1566.
 Lichenoides pulmoneum reticulatum vulgare, marginibus peltiferis. *Dill. musc.* 212. *t.* 219. *f.* 113.
 Lichenoides peltatum arboreum maximum. *Cat. giff.* 208. *R. Syn.* 76.
 Muscus pulmonarius. *Bauh. pin.* 361. *f.* Lichen arborum. *Park.* 1311.
Anglis, Tree Liverwort, or Lungwort.
 Habitat in arborum truncis præsertim Querci, Fraxini.
35. LICHEN foliaceus decumbens laciniatus, laciniis decumbens.
 apice trifidis, scutellis subfuscis.

CRYPTOGAMIA ALGÆ.

Lichenoides gelatinosum fuscum, Jacobææ maritimæ
divisura. *Dill. musc.* 140. t. 19. f. 25.

Anglis, transparent Liverwort.

Habitat prope Lucton in Herefordia legit Littl. Brown.

nigrescens. 36. LICHEN foliaceus rugosus lobatus atro-virens, scu-
tellis confertis rufis.

Lichenoides gelatinosum membranaceum, tenue, ni-
gricans. *Dill. musc.* 138. t. 19. f. 20.

Lichenoides saxatile membranaceum gelatinosum te-
nue nigrescens. *Cat. giff. R. Syn.* 72.

Muscus licheniformis membranaceus nigricans pellu-
cidus, bracteolis innumeris conglomeratis. *Buddl.*
hort. sicc. Vol. II. f. 6.

Anglis, blackish Liverwort.

Habitat in arborum truncis et muris, in sylva Bagley-
Wood dicta prope Oxonium, in Cambria copiose ad
rupes et arborum truncos. *Dill.*

furfura- 37. LICHEN foliaceus decumbens furfuraceus: laciniis
ceus. acutis: subtus lacunosis atris. *Sp. pl.* 1146. *Fl.*
succ. 1088.

Lichenoides cornutum amarum: desuper cinereum:
inferne nigrum. *Dill. musc.* 157. t. 21. f. 52.

Muscus amarus absinthii folio. *Bauh. hist. III.* 764.

Anglis, Branny Liverwort.

Habitat in Arvoniam et ad palos vivarii Croft-Castle in
Herefordia observavit. Littl. Brown. *Dill.*

ampulla- 38. LICHEN foliaceus planiusculus lobatus crenatus,
ceus. peltis globosis inflatis. *Sp. pl.* 1146.

Lichen foliis laciniatis, marginibus convolutis in ve-
siculam abeuntibus. *Roy. lugdb.* 509.

Lichenoides tinctorium glabrum veliculosum. *Dill.*
musc. 138. t. 24. f. 82.

Lichenoides saxatile tinctorium, foliis latioribus non
pilosis, vesiculas proferens. *R. Syn.* 74.

Anglis, Bladder Liverwort.

Habitat in pascuis montanis Emmot-Pasture dictis
prope Coln in Lancastriensi comitatu, D. Rich-
ardson.

39. LICHEN foliaceus erectus compressus ramosus fari-*farinateus*
naceus : lateribus verrucosis. *Sp. pl.* 1146. *Fl. succ.* 1089.
Lichen cinereus angustior, scutis in marginibus seg-
mentorum. *Vaill. parif. t.* 20. *f.* 13. 14. 15.
Lichenoides segmentis angustioribus ad margines ver-
rucosis et pulverulentis. *Dill. musc.* 172. *t.* 23.
f. 63.
Lichenoides arboreum ramosum cinereo-candicans,
segmentis angustioribus. *R. Syn.* 76.
β Lichenoides arboreum ramosum, cinereo-virescens
segmentis angustioribus, circa margines crispis. *R.*
Syn. 76.
γ Lichenoides non tubulosum *πλατυδαρύφυλλον* tuber-
culis pulverulentis donatum. *R. Syn.* 66.
Anglis, mealy Liverwort.
Habitat in arborum truncis.
40. LICHEN foliaceus erectus linearis ramosus lacuno-*calicaris*.
sus convexus mucronatus. *Sp. pl.* 1146. *Fl. succ.*
1090.
Lichenoides coralliforme rostratum et canaliculatum.
Dill. musc. 170. *t.* 23. *f.* 62.
Lichenoides arboreum ramosum angustioribus cine-
reo-virescentibus. *R. Syn.* 76.
Musco-fungus arboreus, capitulis rostratis. *Hist. ox.*
III. 634. *f.* 15. *t.* 7. *f.* 5.
Anglis, channeled Liverwort.
Habitat in rupibus et corticibus arborum. In Cam-
bria, D. Lhwyd ; circa Slingfoldiam in Suffexia;
et copiose ad rupes montis Steiperstone in Salopiensi
comitatu. *Dill.*
41. LICHEN foliaceus erectus oblongus lanceolatus sub-*fraxineus*.
laciniatus lacunofus glaber, scutellis subpeduncula-
tis. *Sp. pl.* 1146. *Fl. succ.* 1091.
Lichenoides longifolium rugosum rigidum. *Dill. musc.*
165. *t.* 22. *f.* 59.
Lichenoides arboreum ramosum scutellatum majus et
rigidius, colore virescente. *R. Syn.* 75.

CRYPTOGAMIA ALGÆ.

Musco-fungus quernus latifolius cinereus. *Hist. ox.*

III. 634. f. 15. t. 7. f. 14. 3. 4.

Anglis, Ash Liverwort.

Habitat in arborum truncis præsertim Fraxinis et Querci.

prunastri. 42. LICHEN foliaceus erectiusculus lacunofus: subtus tomentosus albus. *Sp. pl. 1147. Fl. suec. 1092.*

Lichenoides cornutum, bronchiale molle subtus incanum. *Dill. musc. 160. t. 21. f. 55.*

Lichenoides arboreum ramosum majus et mollius, colore candicante. *R. Syn. 75.*

β Lichenoides corniculatum candidum molle, segmentis angustis. *Dill. musc. 159. t. 21. f. 54.*

Anglis, Thorn Liverwort.

Habitat in Fraxinis, Salicibus, præsertim in Prunis spinosis aliisque frequens.

capreatus. 43. LICHEN pallide viridis rugosus margine undulatus. *Sp. pl. 1147.*

Lichenoides capreatum; rosaceæ expansum, e sulphureo virens. *Dill. musc. 193. t. 25. f. 97.*

Lichenoides crusta foliosa, ex cinereo et luteo virescente, inferne nigra et lævi. *Cat. giff. 207. R. Syn. 73.*

Musco-fungus lichenoides, crustæ modo adnascens major cinereus. *Hist. ox. III. 633. f. 15. t. 7. f. 1.*

Anglis, Rose-leaved Liverwort.

Habitat in saxis et arboribus passim.

juniperinus. 44. LICHEN foliaceus laciniatus crispus fulvus, peltis lividis. *Sp. pl. 1147. Fl. suec. 1093.*

Lichen fulvus sinubus dædaleis laciniatus. *Fl. lapp. 450. Roy. lugdb. 510.*

Anglis, Juniper Liverwort.

Habitat in ramis et truncis arborum.

Convenit cum L. parietino sed differt colore luteo-viridi; foliis laxis erectiusculis; peltis disco bruneis.

45. LICHEN foliaceus depressus lobatus glaber, marginibus crispo farinaceo. *Sp. pl.* 1148. *Fl. suec.* 1094.

Lichenoides endiviæ foliis crispis splendentibus subtus nigricantibus. *Dill. musc.* 192. t. 25. f. 96.

Musco-fungus lichenoides arborum crispus cinereus subtus nigricans. *Hist. ox.* III. 633. f. 15. t. 7. f. 4.

Anglis, shining Liverwort.

Habitat in pascuis montosis Emmot-moor dictis prope Coln in Lancastria; in ericeto Banstead-downs dicto. Th. Manningham. *Dill. musc.*

***** *Coriacei.*

46. LICHEN foliaceus repens lobatus, peltis marginalibus posticis. *Sp. pl.* 1148. *Fl. suec.* 1096. *resupinatus.*

Lichenoides fuscum, peltis posticis ferrugineis. *Dill. musc.* 206. t. 28. f. 105.

Lichenoides saxatile fuscum, peltis in averfa foliorum superficie locatis. *R. Syn.* 77.

Anglis, livid Lichenoides.

Habitat ad lapides minores Pebbles vocatos in ora maritima Suffexiæ ad rupes circa Bishops-castle, in Salopia. *Dill. musc.*

47. LICHEN foliaceus repens laciniatus lacunofus, peltis marginalibus adscendentibus. *sylvaticus.*

Lichenoides polyschides villosum et scabrum, peltis parvis. *Dill. musc.* 199. t. 27. f. 101.

Anglis, rough Liverwort.

Habitat in sylvis passim.

48. LICHEN foliaceus repens lobatus obtusus glaber, peltis marginalibus horizontalibus planis. *horizontalis.*

Lichenoides subfuscum, peltis horizontalibus planis.

Dill. musc. 205. t. 28. f. 104.

Anglis, flat Sheild Liverwort.

Habitat in sylvis; in sylva Enfieldensi. *Dill.*

49. LICHEN foliaceus repens subrotundus planus peltis immersis. *immersus.*

CRYPTOGAMIA ALGÆ.

Lichenoides lichenis facie, peltis acetabulis immersis.

Dill. musc. 221. t. 30. f. 121.

Anglis, round Liverwort.

Habitat in faucibus rupium, Chedder-rocks dictarum in Somersetia, et in monte Snowdon in Cambria.

Dill.

caninus. 50. LICHEN foliaceus repens lobatus obtusus planus :
subtus venosus villosus, pelta marginali adscendente.

Sp. pl. 1149. *Fl. succ.* 1100.

Lichen terrestris cinereus *R. Cat. Ang.* 185.

Lichenoides digitatum cinereum, lactucæ foliis sinu-
osis. *Dill. musc.* 200. t. 27. f. 102.

Lichenoides peltatum terrestre cinereum majus, foliis
divisis, *R. Syn.* 76.

Anglis, Ash-coloured Ground Liverwort.

Habitat in ericetis.

Pharmac. Lichenis cinerei terrestris.

***** Umbilicati, squalentes quasi fuligine.

minutus. 51. LICHEN umbilicatus gibbus punctatus subtus ful-
vus. *Sp. pl.* 1149.

Lichen pulmonarius alpinus cinereus punctatus. *Hall.*
helv. 75. t. 2. f. 2.

Lichenoides coriaceum nebulosum cinereum puncta-
tum subtus fulvum. *Dill. musc.* 223. t. 30. f.
127.

Anglis, cloudy Liverwort.

Habitat ad rupes Chedderenses in Somersetia, et in
summis rupibus udis Hilsvæ in Cambria. *Dill.*

valleus. 52. LICHEN umbilicatus subtus hirsutissimus. *Sp. pl.*
1150. *Fl. succ.* 1103.

Lichenoides coriaceum, latissimo folio umbilicato et
verrucofo. *Dill. musc.* 545. t. 82. f. 5.

Anglis, fleecy Liverwort.

Habitat in rupibus prope Settle in comitatu Eboracensi.
D. Richardson.

pustulatus 53. LICHEN umbilicatus subtus lacunofus. *Sp. pl.* 1150.
Fl. succ. 1107.

Lichenoides

Lichenoides pustulatum cinereum et veluti ambustum. *Dill. musc.* 226. t. 30. f. 131.

Anglis, finged Liverwort.

Habitat in Cambriæ comitatu Radnorienſi primus et ſolus adhuc obſervavit Little Brown, in rupibus meridiem ſpectantibus, ſubter arcem Keven Lees Caſtle, dictam, copioſe. *Dill. musc.*

54. LICHEN umbilicatus undique lævis. *Sp. pl.* 1150. *deuſtus*.
Fl. ſucc. 1105.

Lichenoides coriaceum cinereum, peltis atris compressis. *Dill. musc.* 218. t. 29. f. 117.

Lichenoides ſaxatile, foliis minus diviſis, cinereo-fuscis. *R. Syn.* 73.

Anglis, Sooty Liverwort.

Habitat in rupe Sancti Vincentii prope Briſtoliā, D. Dare.

55. LICHEN umbilicatus polyphyllus undique lævis *polyphyllus* crenatus. *Sp. pl.* 1150. *Fl. ſucc.* 1104.

Lichenoides tenue pullum, foliis utrinque glabris. *Dill. musc.* 225. t. 30. f. 129.

Anglis, ſmooth Liverwort.

Habitat in monte Snowdon ad rupes Llyn Llydaw dictas. *Dill.*

56. LICHEN umbilicatus polyphyllus utrinque lævis *polyrhizos*. *Sp. pl.* 1151. *Fl. ſucc.* 1108.

Lichenoides pullum ſuperne et glabrum, inferne nigrum et cirroſum. *Dill. musc.* 226. t. 30. f. 130.

Anglis, Fibrous Liverwort.

Habitat cum priore.

***** *Scyphiferi.*

57. LICHEN scyphifer ſimplex integerrimus, ſtipite *cocciferus*, lindrico, tuberculis coccineis. *Sp. pl.* 1151. *Fl.*

ſucc. 1109. *Mat. med.* 496.
Coralloides ſcyphiforme, tuberculis coccineis. *Dill. musc.* 82. t. 14. f. 7.

Lichenoides tubuloſum pyxidatum, tuberculis amæne coccineis. *R. Syn.* 69.

Anglis, red Liverwort.

Habitat in ericetis et nemorosis. XI, II.

cornucopi- 58. LICHEN scyphifer simplex folio brevior, tuberculis
oides. coccineis. *Sp. pl.* 1151. *Fl. suec.* 1110.

Coralloides scyphiforme, marginibus radiatis et foliatis. *Dill. musc.* 85. t. 14. f. 9.

Lichenoides pyxidatum, marginibus eleganter foliatis. *R. Syn.* 69.

Anglis, radiated Liverwort.

Habitat in ericetis Hampstedienfi et nigro prope Londinum. Dill. II.

pyxidatus. 59. LICHEN scyphifer simplex crenulatus, tuberculis fuscis. *Sp. pl.* 1151. *Fl. suec.* 1111.

Lichen pyxidatus major. *Vaill. parif. t.* 21. f. 8.

Mich. gen. 82. t. 41. f. 1.

Coralloides scyphiforme, tuberculis fuscis. *Dill. musc.* 79. t. 14. f. 6.

Muscus pyxoides terrestris. *Baub. pin.* 361.

Lichenoides tubulosum pyxidatum cinereum. *Cat. giff.* 204. *R. Syn.* 68.

β Lichenoides tubulosum pyxidatum proliferum. *R. Syn.* 69.

γ Lichenoides tubulosum pyxidatum, tuberculis fusconigris, proliferum. *R. Syn.* 69.

Anglis, Cup Liverwort, or Moss.

Habitat in ericetis et sylvis.

imbriatus 60. LICHEN scyphifer simplex denticulatus, stipite cylindrico. *Sp. pl.* 1152. *Fl. suec.* 1112.

Lichen pyxidatus minor. *Vail. parif. t.* 21. f. 6.

Mich. gen. 83. t. 41. f. 45.

Coralloides scyphiforme gracile, marginibus ferratis. *Dill. musc.* 84. t. 14. f. 8.

Lichenoides tubulosum pyxidatum, marginibus ferratis. *Cat. giff.* 204. *R. Syn.* 69.

β Lichenoides pyxidatum proliferum, marginibus ferratis. *R. Syn.* 69.

Anglis, indented Cup Liverwort.

Habitat in sylvis et ericetis.

filiformis. 61. LICHEN scyphifer simplex integerrimus filiformis tuberculis fuscis. Li-

Lichenoides tubulosum pyxidatum exiguum, fusco-virens. *R. Syn.* 70.

Coralloides scyphis gracilibus tubiformibus, pedicularis folio. *Dill. musc.* 85. *t.* 14. *f.* 10.

Anglis, Trumpet Liverwort.

Habitat in muris et ericetis.

62. LICHEN foliis sub erectis laciniatis crispis scyphiferis, scyphis brevissimis conicis.

Lichenoides cartilaginofum, tubulis et pyxidulis exiguis. *R. Syn.* 70.

Coralloides scyphiforme, foliis albicorniformibus cartilaginosis. *Dill. musc.* 87. *t.* 14. *f.* 12.

Anglis, Leafy Liverwort.

Habitat in ericetis montosis passim.

63. LICHEN scyphifer ramosus denticulatus filiformis. *gracilis*, *Sp. pl.* 1152. *Fl. suec.* 1113.

Coralloides scyphiforme ferratum elatius, caulibus gracilibus glabris. *Dill. musc.* 88. *t.* 14. *f.* 13.

Lichenoides pyxidatum cinereum elatius, ramulis pyxidatim desinentibus. *R. Syn.* 69.

Anglis, tall Liverwort.

Habitat in ericetis et locis petrosis.

64. LICHEN scyphifer ramosissimus: ramis cylindricis *digitatus*, calycibus integris nodosis. *Sp. pl.* 1152. *Fl. suec.* 1114.

Coralloides ramulosum, tuberculis coccineis. *Dill. musc.* 96. *t.* 15. *f.* 19.

Lichenoides coralliforme, apicibus coccineis. *R. Syn.* 68.

Anglis, Fingered Liverwort.

Habitat in ericetis et sylvis. II.

65. LICHEN scyphifer ramosissimus ventricosus, scyphis dentatis, tuberculis fuscis. *ventricosus*.

Lichenoides tubulosum cinereum, valde crustaceum, ramulis brevioribus et crebrioribus ex acetabulis enascentibus. *R. Syn.* 68.

Coralloides cornu-copioides incanum, scyphis cristatis. *Dill. musc.* 94. *t.* 15. *f.* 17.

Anglis,

Anglis, cruſted Liverwort.

Habitat in ſylvis.

- cornutus*. 66. LICHEN ſcyphifer ſimpliciuſculus ſubventricofus, calycibus integris. *Sp. pl.* 1152. *Fl. ſuec.* 1115. Coralloides vix ramoſum, ſcyphis obſcuris. *Dill. muſc.* 60. t. 15. f. 14.

Lichenoides tubuloſum cinereum, non ramoſum. *R.*

Syn. 27.

Anglis, plain Liverwort.

Habitat in ericetis. *XI.* II.

- difformis*. 67. LICHEN ſcyphifer ſimpliciuſculus ſubventricofus, calycibus dentatis. *Sp. pl.* 1152. *Fl. ſuec.* 1116. Lichen pyxoides teres, acetabulis minoribus repandis. *Mich. gen.* 80. t. 41. f. 1.

Coralloides craſſius ſubincanum, calycibus dentatis. *Dill. muſc.* 95. t. 15. f. 18.

Lichenoides tubuloſum minus ramoſum cauliculis craſſioribus difformibus. *R. Syn.* 68.

Anglis, ferrated Liverwort.

Habitat in ericetis.

***** *Fruticuloſi.*

- rangiferi-nus*. 68. LICHEN fruticuloſus perforatus ramoſiſſimus, ramulis nutantibus. *Sp. pl.* 1153. *Fl. ſuec.* 1117.

Lichenoides tubuloſum ramoſiſſimum, fruticuli ſpecie, candicans. *R. Syn.* 66.

Coralloides montanum fruticuli ſpecie, ubique candicans. *Dill. muſc.* 107. t. 16. f. 29.

Musco-fungus coralloides montanus ramoſiſſimus cinereus vulgaris. *Hiſt. ex. III.* 633. f. 15. t. 7. f. 9.

β Coralloides fruticuli ſpecie candicans, corniculis ruſcentibus. *Dill. muſc.* 110. t. 16. f. 30.

Lichenoides tubuloſum ramoſiſſimum, fruticuli ſpecie, candicans, corniculis ruſcentibus. *R. Syn.* 67.

Anglis, Rhendeer Liverwort.

Habitat in ſylvis et ericetis frequens.

- furcatus*. 69. LICHEN fruticuloſus tubuloſus ramoſus, ramis erectis furcatis.

Liche-

Lichenoides tubulosum cinereum, ramosius et crustaceum. *R. Syn.* 67.

Coralloides corniculis brevioribus et crebrioribus.

Dill. musc. 104. t. 16. f. 27.

Musco-fungus montanus corniculatus. *Hist. ex. III.*

632. f. 15. t. 7. f. 1.

Anglis, forked Liverwort.

Habitat in ericetis et locis montosis.

70. LICHEN fruticulosus tubulosus: ramosissimus spinosus, ramulis tuberculatis subdigitatis.

Lichenoides tubulosum ramosum rigidius, majus et crassius, cinereo fuscum. *R. Syn.* 16.

Coralloides sparsum, caulibus tortuosis et spinosis.

Dill. musc. 101. t. 16. f. 25.

Muscus coralloides saxatilis cervi cornua referens.

Bauh. pin. 361.

Anglis, prickly Liverwort.

Habitat in ericetis sed rarius.

71. LICHEN fruticulosus perforatus, ramulis brevissimis uncialis, acutis. *Sp. pl.* 1153. *Fl. suec.* 1118.

Coralloides perforatum minus, molle et tenue. *Dill. musc.* 99. t. 16. f. 22.

Lichenoides tubulosum, cauliculis mollioribus et crassioribus, minus. *R. Syn.* 67.

Anglis, short Liverwort.

Habitat in ericeto Woolwicensi in Cantio, et in monte Steiperstone prope Norbury in comitatu Salopiensi. *Dill.*

72. LICHEN fruticulosus subdichotomus, ramis simplicibus subulatis. *Sp. pl.* 1153. *Fl. suec.* 1119.

Coralloides corniculis longioribus et rarioribus. *Dill. musc.* 102. t. 16. f. 26.

Lichenoides tubulosum cinereum, minus crustaceum minusque ramosum. *R. Syn.* 67.

Muscus corniculatus. *Tab. ic.* 809. *Ger. Em.* 1560. *B. hist. III.* 767. *Park.* 1308.

Anglis, horned Liverwort.

Habitat in ericetis passim.

- paschalis.* 73. LICHEN fruticulosus solidus tectus foliolis crustaceis. *Sp. pl.* 1153. *Fl. suec.* 1120.
 Coralloides crispum et botryforme alpinum. *Dill. musc.* 114. *t.* 17. *f.* 33.
 Lichenoides non tubulosum cinereum ramosum, totum crustaceum. *R. Syn.* 66.
 Musco-fungus coralloides fruticosior et lignosior. *Hist. ox.* III. 633. *f.* 15. *t.* 7. *f.* 12.
Anglis, Woody Liverwort.
 Habitat in montibus Cambriæ altissimis. *Dill.*
- fragilis.* 74. LICHEN fruticosus solidus, ramulis teretibus obtusis. *Sp. pl.* 1154. *Fl. suec.* 1121.
 Coralloides alpinum, corallinæ minoris facie. *Dill. musc.* 116. *t.* 17. *f.* 34.
 Lichenoides non tubulosum ramosissimum fruticuli specie cinereo fuscum. *R. Syn.* 65.
Anglis, brittle Liverwort.
 Habitat in montibus Cambrobritannicis *Dill.* in ericetis montosis Moors, dictis, in Westmorlandia.
- globosus.* 75. LICHEN fruticulosus solidus ramosissimus ramulis teretibus acutis furcatis tuberculis terminalibus globosis.
 Lichenoides non tubulosum, ramulis scutellis nigris terminatis. *R. Syn.* 66.
 Coralloides cupressiforme, capitulis globosis. *Dill. musc.* 117. *t.* 17. *f.* 35.
 Muscus coralloides tunbrigensis, bracteolis nigerrimis. *Pet. mus. N.* 437. *R. Hist.* III. 30.
Anglis, Round-headed Liverwort.
 Habitat in rupibus et locis saxosis. In rupibus prope Tunbridge, in monte Snowdon. *Dill.* In montibus Eboracensibus passim.
- siliquosus.* 76. LICHEN fruticulosus solidus subramosus, tuberculis lateralibus concavis.
 Coralloides fasciculare verrucosum et veluti siliquosum. *Dill. musc.* 119. *t.* 17. *f.* 38.
 Musco-fungus petræus, cornibus indivisis et incurvatis. *Hist. ox.* III. 633. *f.* 15. *t.* 7. *f.* 4.
Anglis, Poddied Liverwort.

Habitat

Habitat in rupibus; in saxa illa grandia, quæ per campos elatos, Grey-Weathers in Wiltoniensi comitatu dictos, dispersa jacent. Dill.

***** Filamentosi.

77. LICHEN filamentosus pendulus, ramis implexis, plicatus. scutellis radiatis. *Sp. pl.* 1154. *Fl. suec.* 1122. *Mat. med.* 495.

Usnea vulgaris, loris longis implexis. *Dill. musc.* 56. *t.* 11. *f.* 1.

Muscus arboreus, Usnea officinarum. *B. pin.* 361. *R. Syn.* 64.

Muscus quernus. *Ger. Em.* 1558.

Muscus arboreus vulgaris et quercinus. *Park.* 1311. *Anglis*, common hairy Liverwort, or Tree Moss.

Habitat in silvis densis sed rarius.

78. LICHEN filamentosus pendulus subarticulatus, ramis patentibus. *Sp. pl.* 1155. *Fl. suec.* 1123.

Usnea barbata, loris tenuibus fibrosis. *Dill. musc.* 63. *t.* 12. *f.* 6.

Muscus capillaceus longissimus. *Bauh. pin.* 361.

Anglis, bearded Liverwort.

Habitat in sylva dicta Forrest of Dean in Glocestriensi comitatu, et in sylva prope Bishop's-Castle in Salopia. Dill.

79. LICHEN filamentosus pendulus: axillis compressis. *Sp. pl.* 1155. *Fl. suec.* 1124. *jubatus.*

Usnea jubata nigricans. *Dill. musc.* 64. *t.* 12. *f.* 7.

Muscus corallinus faxatilis fœniculaceus. *R. Syn.* 65.

Musco-fungus trichoides e nigridine cinereus, jubæ ad instar e saxis et arboribus dependens. *Hist. ox.*

III. 633.

Anglis, Fennel Liverwort.

Habitat in sylvis et rupibus; in Charley Forest, Leicestershire. Petiver. *In comitatu Eboracensi passim.* D. Richardson.

80. LICHEN filamentosus ramosissimus decumbens implexatus, plicatus opacus. *Sp. pl.* 1155. *Fl. suec.* 1125.

Usnea

Usnea lanæ nigræ instar saxis adhærens. Dill. musc.
66. t. 13. f. 8.

Muscus coralloides, lanæ nigræ instar saxis adhærens.
R. Syn. 65.

Anglis, Woolly Liverwort.

Habitat in rupibus. In Cornubia D. Stephens in rupibus Cambriæ circa Borth, milliaris distantia a Bangoria. S. Brewer; in Westmorlandia frequens.

chalybei-
formis.

81. LICHEN filamentosus subramosus decumbens implicato-flexuosus. *Sp. pl. 1155. Fl. suec. 1127.*

Usnea rigida horsum vorsum extensa. Dill. musc. 66.
t. 13. f. 10.

Muscus caule rigido instar fili chalybei. R. Syn. 65.
Anglis, Wiry Liverwort.

Habitat in truncis arborum et locis saxosis. In monte Salopiæ, Steiperstone dicto. Dill.

hirtus.

82. LICHEN filamentosus ramosissimus erectus, tuberculis farinaceis sparsis. *Fl. lapp. 989. suec. 1128.*
Sp. pl. 1155.

Usnea vulgarissima tenuior et brevior sine orbiculis.
Dill. musc. 67. t. 13. f. 12.

Muscus ramosus. Ger. 1372. R. Syn. 65.
Anglis, rough Liverwort.

Habitat in sylvis et sepibus.

vulpinus.

83. LICHEN filamentosus ramosissimus erectus fastigiatus inæquali-angulosus. *Sp. pl. 1155. Fl. suec. 1129.*

Usnea capillacea citrina, fruticuli specie. Dill. musc.
72. t. 13. f. 16.

Muscus aureus tenuissimus. Mer. pin. R. Syn. 65.
Anglis, Gold Liverwort.

Habitat in sylvis passim. Prope Totteridge in Hartfordia. Dill.

articulatus.

84. LICHEN filamentosus articulatus, ramulis tenuissimis punctatis. *Sp. pl. 1156. Roy. lugdb. 513.*

Usnea

Usnea capillacea et nodosa. Dill. musc. 60. t. 11.
f. 4.

Muscus arboreus nodosus. Bauh. pin. 361. R. Syn.
65. f. geniculatus. Park. 1311.

Anglis, Jointed Liverwort.

Habitat in sylvis sed rarius; in sylva ad villam Stoken-Church; prope Burnley in Lancastria. Dill.

85. *LICHEN filamentosus ramosus erectus, scutellis floridis, radiatis.* Sp. pl. 1156. Fl. suec. 1130.

Usnea vulgatissima tenuior et brevior cum orbiculis.
Dill. musc. 69. t. 13. f. 13.

Muscus arboreus cum orbiculis. Bauh. pin. 361.
R. Syn. 65.

Muscus ramosus floridus. Tab. ic. 808. Ger. Em.
1560.

Anglis, Flowering Liverwort,

Habitat in sylvis et dumetis.

TREMELLA. 1067.

Fruetificationes vix manifestæ, in corpore gelatinoso.

1. *TREMELLA plicata undulata.* Sp. pl. 1157. Fl. Nostoc.
suec. 1136.

Tremella terrestris sinuosa pinguis et fugax. Dill.
musc. 52. t. 10. f. 14.

Ulva terrestris pinguis et fugax. R. Syn. 64.

Nostoc paracelsi. Act. parif. 1708. p. 228.

Anglis, Jelly Tremella.

Habitat in pratis post pluvius.

2. *TREMELLA fessilis membranacea auriformis cinea auricula.*
rea. Sp. pl. 1157. Fl. suec. 1137.

Agaricum auriculæ forma. Mich. gen. 124. t. 66.
f. 1.

Peziza auriculam referens. R. Syn. 18.

Fungus membranaceus auriculam referens, f. sambucinus. Bauh. pin. 372.

Fungus sambucinus, f. auricula Judæ. Ger. 1385.
Park. 1320.

Anglis,

Anglis, Eared Tremella, or Jews-ear.
Habitat in lignis putridis.

Lichenoides. 3. TREMELLA erectis planis : margine crispo lacinato. *Sp. pl.* 1157. *Fl. suec.* 1138.

Lichenoides pellucidum, endiviæ foliis tenuibus crispis. *Dill. musc.* 143. t. 19. f. 31.

Lichenoides tenue rufescens. *R. Syn.* 77.

Anglis, transparent Tremella.

Habitat in ericetis et umbrosis.

verrucosa. 4. TREMELLA tuberculosa solida rugosa. *Sp. pl.* 1158. *Fl. suec.*

Tremella fluviatilis gelatinosa et uterculosa. *Dill. musc.* 54. t. 10. f. 16.

Anglis, Warty Tremella.

Habitat supra lapides in rivulis. In rivulis alpinis in Arvoniam. *Dill. In fluvio Martin River dicto in comitatu Surriensi.*

hemisphærica. 5. TREMELLA hemisphærica sparsa. *Sp. pl.* 1158. *Fl. suec.* 1141.

Anglis, Sea Tremella.

Habitat in Conservis et Fucis marinis.

purpurea. 6. TREMELLA subglobosa sessilis solitaria glabra. *Sp. pl.* 1158. *Fl. suec.* 1142.

Tremella palustris, vesiculis sphæricis fungiformibus. *Dill. musc.* 55. t. 10. f. 17.

Lichenoides fungiforme, capitulis vel vesiculis sphæricis, aqueo humore repletis. *R. Syn.* 70.

Anglis, Purple Tremella.

Habitat ad margines fossarum inque fossis exsiccatis versus Newington et Hackney prope Londinum. *Dill.*

C H A R A. 1066.

Cal. biphyllus. *Anthera* globosa sessilis. *Stigmata tria.* *Sem.* unicum, subrotundum.

tomentosa. 1. CHARA aculeis caulinis ovatis. *Fl. suec.* 1132. *Sp. pl.* 1156.

Chara major subcinerea fragilis. Act. paris. 1719 p.
18. *R. Syn.* 132.

Equisetum fragile majus subcinereum aquis immer-
sium. Hist. ox. III. 621. t. 4. f. 9.

Equisetum f. Hippuris coralloides. Ger. Em. 1115.
Anglis, brittle Chara.

Habitat in fossis et stagnis near Chiffelhurst in Kent.

2. *CHARA* caulibus lævibus, frondibus interne denta-*vulgaris.*
tis. Sp. pl. 1156. Fl. suec. 1131.

Chara vulgaris foetida. Act. paris. 1719. p. 17. t. 3.
f. 1. R. Syn. 132.

Equisetum foetidum sub aqua repens. Ger. Em. 1115.
Bauh. pin. 16. hist. III. 731. Park. 1201.

Anglis, common Chara.

Habitat in fossis et stagnis frequens.

3. *CHARA* aculeis caulinis capillaribus confertis. *Sp. hispida.*
pl. 1156. Fl. suec. 1133.

Chara major caulibus spinosis. Act. paris. 1719. p.
18. *t. 3. f. 3.*

Equisetum f. Hippuris muscosus sub aqua repens.
Pluk. ph. t. 193. f. 6.

Anglis, prickly Chara.

Habitat in fossis et stagnis; in fossis turfosis in comitatu
Westmorlandico.

4. *CHARA* caulium articulis inermibus diaphanis super-*flexilis.*
ne latioribus. Sp. pl. 1157. It. gotl. 215. Fl. suec.
1134.

Chara translucens minor flexilis. R. Syn. 133.
Anglis, smooth Chara.

Habitat in fossis. In the ponds about Ipswich in Suf-
folk. Buddl. In a Bog where the stinking Sparw
empties itself, nigh Knaissborow. Dr. Richardson.
R. Syn.

F U C U S. 1068.

Masc. Vesiculæ villis intertextæ.

Fem. Vesiculæ adspersæ granis immersis, apice promi-
nulis.

Sem. solitaria.

H h

* *Dicho-*

* *Dichotomi frondescentes.*

serratus. γ. FUCUS planus dichotomus serratus ad apices tuberculatus. *Sp. pl.* 1158. *Fl. suec.* 1144.

· Fucus f. Alga latifolia major dentata. *R. Syn.* 42. *Hist. ox. III.* 648. *t. 9. f. 1.*

β Fucus marina humilis, latifoliæ ferratæ similis. *R. Syn.* 42.

Anglis, serrated Fucus, or Sea Wrack.

Habitat in rupibus et saxis marinis frequens.

vesiculosus 2. FUCUS dichotomus integer, caule medium folium transcurrente, vesiculis verrucosis terminalibus. *Sp. pl.* 1158. *Fl. lapp.* 466. *suec.* 1145. *Roy. lugdb.* 514. *Guett. stamp.* II. 401.

Fucus f. Alga marina latifolia vulgarissima. *R. Syn.* 40.

Fucus maritimus vel Quercus maritima vesiculas habens. *Bauh. pin.* 363.

· Quercus marina. *Lob. ic. 2. p.* 252. *Ger. Em.* 1116.

Quercus marina herbacea et varietas. *Park.* 1293.

β Fucus maritimus f. Quercus maritima foliorum extremis tumidis. *Bauh. pin.* 365.

Anglis, common Fucus, or Sea-Oak.

Habitat in rupibus et saxis marinis ubique.

ceranoides 3. FUCUS dichotomus planus integer : apicibus bifidis vesiculosis. *Fl. lapp.* 465. *suec.* 1146. *Sp. pl.* 1158.

Fucus humilis dichotomus ceranoides, latioribus foliis ut plurimum verrucosis. *Hist. ox. III.* 646. *t. 8. f. 13.*

Fucus membranaceus ceranoides varie dissectus. *R. Syn.* 44.

β Fucus parvus cauliculis teretibus, summitatibus membranaceis dilatatis et laceratis. *Buddl. hort. succ. I.* 47. *R. Syn.* 44.

γ Fucus membranaceus ceranoides ramosus, per ficitatem obsolete virescens. *Pluk. alm.* 161. *R. Syn.* 44.

♂ Fucus

- ♂ *Fucus ceranoides albidus*, ramulorum apicibus stellatis. *Pluk. alm.* 161. *R. Syn.* 44.
Anglis, Buckthorn *Fucus*.
Habitat in littoribus marinis ubique.
4. *FUCUS dichotomus integer* caule folium percurren- *spiralis*.
 te : inferne nudo, vesiculis verrucosis terminalibus.
Fl. lapp. 467. *succ.* 1147. *Sp. pl.* 1159.
Fucus spiralis maritimus major. *Tourn.* 568. *R. Syn.* 41.
 ♂ *Fucus spiralis maritimus minor.* *Tourn.* 568. *R. Syn.* 42.
Anglis, twisted *Fucus*.
Habitat in littore Cantiano et Sufflexiano frequens.
5. *FUCUS membranaceus linearis dichotomus divarica-* *divarica-*
 tus integer : ramificationibus vesiculosis. *Sp. pl. tus.* 1159.
Fucus bullatus fruticescens, caule nudo, foliis rectis compressis bifidis. *Hist. ox. III.* 647. *f.* 15. *t.* 8. *f.* 5.
Anglis, shrubby *Fucus*.
Habitat in littore Insulæ Anglesey. Hist. ox. prope Sheerness in Insula Sheepy.
6. *FUCUS dichotomus planus linearis acutus vesiculis linearis,*
ovatis sparsis.
Fucus longo angusto crassoque folio. *Bauh. pin.* 364. *pr.* 155, 5. *R. Syn.* 43.
Fucus marinus secundus. *Dod. pemp.* 479. *Park.* 1293.
Quercus marina 2. *Ger. Em.* 1115.
Anglis, narrow-leaved *Fucus*, or Sea-Thongs.
Habitat in littoribus marinis passim.

** *Dichotomi caulescentes.*

7. *FUCUS dichotomus teres uniformis fastigiatus, ves-* *fastigia-*
culis terminalibus bifidis. *tus.*
Fucus dichotomus teres uniformis fastigiatus. *Sp. pl.* 1162.

CRYPTOGAMIA ALGÆ.

Fucus palmaris tenuis in orbem expansus, in segmenta bifida breviora teretia divisus. *Hist. ox. III.* 649.

t. 9. f. 9.

Fucus i alga exigua dichotomus foliorum segmentis longiusculis, crassis et subrotundis. *R. Syn.* 45.

Anglis, forked *Fucus*.

Habitat in littoribus marinis frequens.

excissus. 8. *FUCUS* dichotomus linearis excavatus: axillis divaricatis. *Sp. pl.* 1159.

Fucus pumilus dichotomus, segmentis ex una parte gibbosis, ex altera excavatis. *R. Syn.* 43.

Fucus dichotomus ceranoides, angulos rotundiusculos efformans. *Hist. ox. III.* 648. *t. 8. f. 11.*

Anglis, furrowed *Fucus*.

Habitat in littoribus marinis.

nodosus. 9. *FUCUS* compressus dichotomus medio ramorum in vesiculam dilatato. *Roy. lugdb.* 514. *Sp. pl.* 1159. *Fl. suec.* 1149.

Fucus maritimus nodosus. *Bauh. pin.* 365. *R. Syn.* 48.

Fucus maritimus, vesiculis majoribus singularibus per intervalla dispositis. *Hist. ox. III.* 647. *t. 8. f. 2.*

Anglis, knotted *Fucus*.

Habitat in rupibus et saxis marinis passim.

*** *Ramosi foliis distinctis.*

caulus. 10. *FUCUS* caule teretiufculo compresso, foliis oblongis integerrimis.

Anglis, Oval-leaved *Fucus*.

Habitat in littore Eboracensi.

**** *Ramosi fronde unita.*

siliquosus. 11. *FUCUS* teres ramosissimus, pedunculis alternis vesiculis oblongis acuminatis. *Fl. lapp.* 463. *suec.* 1150. *Sp. pl.* 1160.

Fucus angustifolius vesiculis longis siliquarum æmulis. *R. Syn.* 48.

[*Fucus*

Fucus maritimus alter, tuberculis paucissimis. *B. pin.*
368.

Quercus marina quarta. *Ger. Em.* 1569.

Anglis, Podded *Fucus*.

Habitat in littoribus marinis.

12. *FUCUS* vesiculis oblongis pedunculatis: terminatis barbatus.
foliolis linearibus. *Sp. pl.* 1161.

Fucus folio tenuissimo diviso filiquatus. *R. Syn.* 48.

Fucus folliculaceus foliis abrotani. *Bauh. pin.* 365.

Fucus maritimus foliis tumidis barbatis. *Bauh. pin.*
365.

Muscus marinus abrotanoides. *Park.* 1290.

Anglis, Bearded *Fucus*.

Habitat in littoribus Suffexiano et Cornubiensi.

13. *FUCUS* caule tereti ramosissimo, foliis alternis su- tamarisci-
bulatis, vesiculis rotundis imbricatis sessilibus axilla- folius.
ribus.

Fucus foliis *Ericæ* f. *Tamarisci*. *R. Syn.* 49.

Fucus tamarisco similis maritima. *Bauh. pin.* 365.

Erica marina quibusdam. *Bauh. hist. III.* 799.

Anglis, Tamarisk-leaved *Fucus*.

Habitat in littoribus marinis. On the coast of Corn-
wal. *R. Syn.* in littore Eboracensi passim.

14. *FUCUS* vesiculis ovatis alternis pedunculatis, fron- foenicula-
dibus multifidis linearibus. *Sp. pl.* 1161. ceus.

Fucus abrotanoides vesiculis parvis rotundis ornatus.

Hist. ox. III. 648. t. 8. f. 17.

Fucus radicibus arborum fibrosis similis. *R. Syn.* 49.

Fucus folliculaceus: foeniculi folio. *B. pin.* 365.

Anglis, Fennel-leaved *Fucus*.

Habitat in littoribus Eboracensi et Lancastriensi.

15. *FUCUS* bipinnatus subfiliformis: apicibus vesiculosis abrotani-
dilatatis. *Sp. pl.* 1161. folius.

Anglis, Jointed *Fucus*.

Habitat in rupibus marinis in littore Suffexiano,

16. *FUCUS* filiformis teres, ramis geminis ramosissimis, usneoides,
fructifi-

fructificationibus innatis concatenatis, foliolis ac-
rofis. *Syst. nat.* 1345 ?

Fucus angustifolius foliis dentatis. *R. Syn.* 48.

Fucus tenuifolius foliis dentatis. *Hist. ox.* III. 648.

t. 9. f. 4

Anglis, prickly *Fucus*.

Habitat in littore Cornubiensi copiose. *R. Syn.*

***** *Fructificationibus non vesicariis.*

muscoïdes. 17. *FUCUS* teretes ramosissimus, ramis sparsis spinis
molliibus alternis. *Sp. pl.* 1161.

Fucus folliculaceus fuscus f. nigricans ex insula cy-
pro. *Pet. mus.* 219. *R. hist.* III. 13.

Anglis, Thorny *Fucus*.

Habitat in littore Eboracensi sed rarius.

incurvus. 18. *FUCUS* teres ramosus, ramulis imbricatis, laciniiis
secundis incurvis.

Pinus maritima f. Fucus teres, cujus ramuli fetis fur-
sum tendentibus sunt obfiti. *R. Syn.* 50.

Anglis, black *Fucus*, or Sea Pine.

Habitat in littore Suffexiano copiose.

plicatus. 19. *FUCUS* capillaris uniformis ramosissimus implicatus
subdiaphanus.

Fucus trichoides nostras aurei coloris, ramulorum a-
picibus furcatis. *Pluk. ph. t.* 184. *f. 2. R. Syn.*
45.

β *Fucus coralloides erectus.* *R. Syn.* 51.

Anglis, Matted *Fucus*.

Habitat in littoribus marinis.

albus. 20. *FUCUS* filiformis teretiusculus subdichotomus, ge-
niculis tumidiusculis, ramis distantibus acutis.

Fucus teres albus tenuissime divisus. *R. Syn.* 50.

Anglis, white *Fucus*.

Habitat in littore Suffexiano. *R. Syn.*

verrucosus 21. *FUCUS* filiformis teres ramosus, ramis simplicibus,
vesiculis rotundis lateralibus.

Fucus

- Fucus marinus purpurascens parvus, caule et ramulis f. foliolis teretibus. R. Syn. 50.*
Anglis, Warty Fucus.
Habitat in littoribus marinis passim.
22. *FUCUS* filiformis teres ramosissimus, ramis alternis, *purpureus*.
 ramulis confertis fructiferis, fructibus globosis.
Fucus teres rubens ramosissimus. R. Syn. 51.
- β *Fucus teres rubens minus ramosus in longum protensus. R. Syn. 51.*
Anglis, Purple Fucus.
Habitat in saxis et rupibus marinis.
23. *FUCUS* caule tereti ramoso, ramis alternis ramo-*scorpioides*
 fissimis apice inflexis.
Fucoides erectum fruticuli specie, summitatibus inflexis. R. Syn. 38. t. 2. f. 6.
Anglis, upright Fucus.
Habitat in littore Suffexiano; at Selfey-Island plentifully. Dr. Dill. R. Syn.
24. *FUCUS* teres subdichotomus uniformis, ramis ob-*rotundus*.
 tusis verrucosis.
Fucus kali geniculato similis, non tamen geniculatus. R. Syn. 43.
Anglis, round Fucus.
Habitat in rupibus marinis prope S. Ives in Cornubia. R. Syn.
25. *FUCUS* teres dichotomus, ramis æqualibus acutis. *lumbrica-*
Fucus parvus segmentis prælongis teretibus acutis. R. lis.
Syn. 45. Hist. ox. III. 648. f. 15. t. 9. f. 4.
Fucus forcellata lumbricalis species. B. pin. 366.
Anglis, sharp-pointed Fucus.
Habitat in littoribus marinis passim.
26. *FUCUS* gelatinosus frondibus teretiusculis ramosis. *gelatino-*
Fucus spongiosus nodosus. Johnsf. it. cant. 3. f. 3. fus.
Ger. Em. 1570. R. Syn. 49.
Spongia ramosa altera anglica. Park. 1304.
Anglis, Gelly Fucus.
Habitat in littore maris prope Sheerness copiose.

filiformis. 27. FUCUS filiformis dichotomus planus.

Anglis, flat Fucus.

Habitat in littore Lancastrienti.

crispus. 28. FUCUS frondibus planis dichotomis, apicibus dilatatis bifidis.

Fucus membranaceus purpureus, variæ ramosus. *R. Syn.* 47.

Alga crispa scabiosa rubra et pallida. *B. hist.* III. 795.

Anglis, curled Fucus.

Habitat in littoribus marinis frequens.

palmatus. 29. FUCUS frondibus planis palmatis. *Roy. lugdb.* 515. *Sp. pl.* 1162.

Fucus membranaceus ceranoides. *B. pr.* 155. *R. Syn.* 46.

Fucus foliaceus humilis, palmam humanam referens. *Hist. ox.* III. 646. *f.* 15. *t.* 8. *f.* 1.

Anglis, Handed Fucus.

Habitat in littoribus marinis passim.

pavonicus. 30. FUCUS fronde sessili reniformi decussatim striata. *Sp. pl.* 1162.

Fucus maritimus gallopavonis pennas referens. *Bauh. pr.* 155. *Hist. ox.* III. 645. *t.* 8 *f.* 7. *R. Syn.* 43.

Anglis, striated Fucus, or Turkey-Feather.

Habitat in rupibus et saxis marinis, ex Insula Anglesæ, et littore Cornubiensi accepit Bobartus; prope Harvicum collegit, D. Buddle; et prope Exmouth Devonix, oppidum copiose nascentem observavit D. Stevens. *R. Syn. in littore Insulæ Walney, sed rarius.*

ciliatus. 31. FUCUS frondibus planis ramosis ciliatis.

Fucus membranaceus rubens angustifolius, marginibus ligulis armatis. *R. Syn.* 47.

Fucus humilis membranaceus acaulos elegantissimus ruber, capillis longis fimbriatis. *Hist. ox.* III. 646. 10.

β Fu-

- ¶ *Fucus membranaceus purpureus latifolius pinnatus.*

R. Syn. 47.

Anglis, ciliated Fucus.

Habitat in littoribus marinis frequens.

32. *FUCUS* frondibus planis ramosis, ramis dentato-pinnatifidus.
pinnatifidis marginibus callosis.

Fucus dealensis pedicularis rubræ folijs. Mus. pet.

405. *R. Syn. 48.*

Alga cervi cornu divisura. B. hist. III. 797.

Anglis, jagged Fucus.

Habitat in saxis et rupibus marinis. Found about Deal by Mr. Dandridge. R. Syn. in littore Devonix frequens.

33. *FUCUS* frondibus planis subdichotomis, ramis al-alatus.
ternis decurrentibus bifidis.

Fucus dichotomus parvus costatus et membranaceus.

Buddl. hort. sic. I. 12. R. Syn. 44.

Fucus purpureus tenuiter divisus non geniculatus.

Hist. ox. III. 646.

Anglis, Winged Fucus.

Habitat in littoribus marinis frequens.

34. *FUCUS* cartilagineus compressus ramosissimus, ramis cartilagineis.
alternatim secundis, fructibus globosis late-neus.
ralibus.

Fucoides rubens varie dissectum. R. Syn. 37.

Anglis, Scarlet Fucus.

Habitat in littoribus marinis frequens.

35. *FUCUS* caule teretiusculo compresso ramoso, ramis plumosus.
duplicato pinnatis coloratis.

Fucoides purpureum eleganter plumosum. R. Syn.

38. t. 2. f. 5.

Anglis, Feathered Fucus.

Habitat in littoribus marinis passim.

36. *FUCUS* cartilagineus compressus, frondibus supra-filicinus.
decompositis pinnatis, laciniis obtusis.

Anglis, Fern-leaved Fucus.

Habitat in littore Lancastriensi sed rarius.

- pinnatus.* 37. FUCUS membranaceus, frondibus triplicato-pinnatis.
Anglis, Pinnated Fucus.
 Habitat in littoribus Eboracensi et Sufflexiano.
- corneus.* 38. FUCUS cartilagineus, caule teretiusculo compresso ramofissimo, ramis subpinnatis, laciniis acutis fructiferis.
 Fucus flavicans teretifolius, ramulis pennatim enascentibus. *R. Syn.* 50.
 Fruticulus parvus pennatus flavicans, corneus et tenax. *Buddl. hort. succ. II.* 2.
Anglis, Horny Fucus.
 Habitat in littore Devonix passim.
- saccharinus.* 39. FUCUS caule tereti brevissimo, fronde ensiformi simplici.
 Fucus caule tereti brevissimo, fronde maxima subsimplici. *Sp. pl.* 1161. *Fl. lapp.* 460. *succ.* 1151.
 Fucus longissimo latissimo crasloque folio. *Bauh. pr.* 154. *R. Syn.* 39.
 β Fucus folio singulari longissimo, lato, in medio rugoso, qui balteiformis dici potest. *R. Syn.* 39.
 γ Fucus latissimus et longissimus oris crispis. *R. Syn.* 38.
Anglis, sweet Fucus.
 Habitat in rupibus et saxi marinis.
- confervoides.* 40. FUCUS caule tereti ramofissimo, ramulis alternis fasciculatis brevissimis tuberculatis scabris.
Anglis, rough Fucus.
 Habitat in littore Eboracensi.
- digitatus.* 41. FUCUS caule tereti brevissimo, fronde palmata laciniis ensiformibus.
 Fucus arboreus polyschides edulis. *B. pin.* 364. *R. Syn.* 46.
 Fucus phasganoides et polyschides. *Ger. Em.* 1570.
 Fucus maximus polyschides. *Park.* 1292.
 β Fucus membranaceus polyphyllus major. *R. Syn.* 46.
 γ Fucus

- γ *Fucus scoticus latissimus edulis dulcis.* *Sibb. pr. 26.*
R. Syn. 46.
Fucus alatus f phasganoides. *Baub. pin. 362.*
Anglis, Fingered Fucus.
Habitat in rupibus marinis.
42. *FUCUS* caule tereti ramoso, frondibus oblongo-*n. rubens.*
dulatis sinuatis. *Roy. lugdb. 514. Sp. pl. 1162.*
Alga minor suave rubens varie divisa. *Mart. cent.*
32. t. 32.
Anglis, red Fucus.
Habitat in littore Cornubiensi.
43. *FUCUS* caule tereti ramoso, frondibus oblongo-*sanguineus*
lanceolatis undulatis integerrimis.
Fucus f. Alga folio membranaceo purpureo, lapathi
sanguinei figura et magnitudine. *Hist. ox. III. 645.*
t. 8. f. 6. R. Syn. 49.
Anglis, Dock-leaved Fucus.
Habitat in littoribus Eboracensi et Cornubiensi fre-
quens.
44. *FUCUS* frondibus ramosis membranaceus, ramis *laciniatus.*
dilatatis palmatis.
Anglis, jagged Fucus.
Habitat in littore Eboracensi copiose.
45. *FUCUS* filiformis simplex subfragilis opacus. *Sp. pl. Filum.*
1162. Fl. suec. 1152.
Fucus chordam referens teres prælongus. *R. Syn. 40.*
Filum marinum germanicum. *B. pr. 155.*
Ligula marina alba Suffexiana. *Pet. gag. t. 91. f. 5.*
Anglis, Thread Fucus.
Habitat in rupibus marinis.

U L V A. 1069.

Fruetificationes in membrana diaphana.

1. *ULVA* tubulosa simplex. *Fl. lapp. 458. suec. 1154. intestinalis.*
Sp. pl. 1163.
Ulva marina tubulosa, intestinorum figuram referens.
R. Syn. 62.

Fucus

Fucus cavus. Bauh. pin. 364. hist. III. 803.
Tremella marina tubulosa intestinorum figura. Dill.
 musc. 47. t. 9. f. 7.
Anglis, Gutt Laver.
Habitat in fossis maritimis.

compressa. 2. *ULVA* tubulosa ramosa compressa. It. gotl. 262.
 Fl. suec. 1155. Sp. pl. 1163.
Ulva marina tenuissima et compressa. R. Syn. 63.
Anglis, flat Laver.
Habitat in faxis et rupibus marinis frequens.

articulata. 3. *ULVA* tubulosa ramosissima articulata, articulis cylindricis, ramis oppositis.
Corallina lenta purpurea compressa. R. Syn. 34.
Fucus purpureus humilis tenuiter divisus geniculatus.
 Hist. ox. III. 646. f. 15. t. 8. f. 14.
Anglis, Jointed Laver.
Habitat in littore Cornubiæ. R. Syn. in littore Insulæ Walney passim.

ramosa. 4. *ULVA* fronde ramosa plana undulata purpurea,
Anglis, Branched Laver.
Habitat in littore Lancastriensi.

dichotoma. 5. *ULVA* fronde dichotoma viridis.
Fucus membranaceus dichotomus gramineus. R. Syn.
 45.
Anglis, Grass-leaved Laver.
Habitat in littore Insulæ Walney passim.

latissima. 6. *ULVA* oblonga plana undulata membranacea viridis. Sp. pl. 1163.
Fucus longissimo latissimo tenuique folio. Bauh. pr.
 154. R. Syn. 40.
Anglis, curled Laver.
Habitat in littoribus marinis; in littore Insulæ Shepey. R. Syn.
Obs. Variat colore purpureo.

7. ULVA palmata prolifera membranacea: ramentis *Lactuca*.
 inferne angustatis. *Sp. pl.* 1163.
 Ulva marina lactucæ similis. *R. Syn.* 62.
 Tremella marina vulgaris, lactucæ similis. *Dill.*
musc. 42. t. 8. f. 1.
 Fucus marinus, lactuca marina dictus. *Park.* 1293.
 Muscus marinus lactucæ similis. *Bauh. pin.* 364.
 Lichen marinus. *Ger. Em.*
- β Ulva palustris lactucæ marinæ similis, sed multo minor et tenerior. *R. Syn.* 63.
 Tremella palustris, vulgari marinæ similis, sed minor et tenerior. *Dill. musc.* 43. t. 8. f. 2.
Anglis, Green Laver, or Oyfter Green.
 Habitat in mari β in fossis pone Newington prope Londinum. *Dill.*
8. ULVA fronde dilatata subfinuata, centro radicata. *umbilicatis*.
Sp. pl. 1163.
 Ulva marina umbilicata. *R. Syn.* 62.
 Fucus umbilicus marinus dictus. *Bauh. pin.* 364.
 Tremella marina umbilicata. *Dill. musc.* 46. t. 8. f. 3.
Anglis, Navel Laver.
 Habitat in saxis et rupibus marinis, pone Sheerness sparsim. *Dill. in littoribus Lancastriensi et Eboracensi passim.*
9. ULVA fronde oblonga bullata. *Sp. pl.* 1163. *Linza*.
 Ulva marina fasciata. *R. Syn.* 62.
 Tremella marina fasciata. *Dill. musc.* 46. t. 9. f. 6.
 Fucus Linza dictus, lactucæ marinæ similis. *B. hist.* III. 801.
 Muscus lactucæ marinæ similis. *Bauh. pin.* 364.
Anglis, Ribbin Laver.
 Habitat in littore marino pone Sheerness. *R. Syn.*
10. ULVA sphærica aggregata. *Fl. suec.* 1160. *Sp. pl. granulata*. 1164.
 Tremella palustris, vesiculis sphæricis fungiformibus.
Dill. musc. 55. t. 10. f. 17.

Lichenoides fungiforme, capitulis vel vesiculis sphæricis, aqueo humore repletis. *R. Syn.* 70.

Anglis, Bladder Laver.

Habitat *ad margines fossarum inque fossis exsiccatis inter Newington et Hackney prope Londinum.* *R. Syn.*

C O N F E R V A. 1070.

Tubercula inæqualia, in fibris capillaribus, longissimis.

* *Filamentis simplicibus, æqualibus, geniculis destitutis.*

rivularis. 1. CONFERVA filamentis simplicissimis æqualibus longissimis. *Fl. suec.* 1161. *Sp. pl.* 1164.

Conferva fluviatilis sericea vulgaris et fluitans. *Dill. musc.* 12. t. 2. f. 1.

Conferva Plinii. *Gesn. Cat.* 23. *R. Syn.* 58.

Conferva palustris sericea, filamentis crassioribus et longioribus. *R. Syn.* 477.

Conferva palustris sericea, crassior et varæ extensa. *Dill. musc.* 13. t. 2. f. 2.

Anglis, River Conferva.

Habitat *in rivulis.*

fontinalis. 2. CONFERVA filamentis simplicissimis æqualibus digito-brevioribus. *Sp. pl.* 1164. *Fl. suec.* 1163.

Conferva minima byssi facie. *Dill. musc.* 14. t. 2. f. 3.

Conferva fontalis fusca, omnium minima mollis. *R. Syn.* 58.

Anglis, Spring Conferva.

Habitat *in rivo novo versus Hornsey.* *Dill.*

** *Filamentis ramosis, æqualibus.*

furcata. 3. CONFERVA filamentis æqualibus extremitatibus ramosis, ramis sub-simplicibus.

Conferva capillacea, filamentis bifidis et trifidis. *Dill. musc.* 16. t. 2. f. 6.

Conferva rivulorum bicornis, filamentis tenuissimis. *R. Syn.* 58.

β Conferva palustris, filamentis brevioribus et crassioribus. *Dill. musc.* 17. t. 3. f. 10.

Conferva

Conferva palustris subhirsuta filamentis brevioribus et crassioribus. R. Syn. 477.

Anglis, forked Conferva.

Habitat in rivulis & in fossis passim.

4. CONFERVA filamentis æqualibus ramosissimis, ramis ramulisque longissimis sparsis. *foeniculacea.*

Conferva marina foeniculacea. Dill. musc. 16. t. 2. f. 8.

Anglis, Fennel-leaved Conferva.

Habitat in rupibus maritimis in Mona Insula. Dill.

5. CONFERVA filamentis æqualibus ramosissimis, ramulis brevissimis alternis. *rigida.*

Conferva fluvialis fibrillosa, subrigida. Dill. musc. 21. t. 4. f. 16.

Anglis, Rough Conferva.

Habitat in fluvio qui ericetum Hounslow percurrit. Dill. in fluvio Lug. prope Mortimer's Cross in Herefordienſi comitatu. L. Brown.

6. CONFERVA filamentis æqualibus ramosis, exsiccatione coeuntibus in aculeos. *amphibia.* *Sp. pl. 1164. Fl. suec. 1164.*

Conferva amphibia, fibrillosa et spongiosa. Dill. musc. 22. t. 4. f. 17.

Conferva terrestris exilis fibrillosa. R. Syn. 59.

Fungus vel potius spongia viridis doliolis adnascenti similis. R. Syn. 57.

Anglis, amphibious Conferva.

Habitat ad ripam Thameſis frequens.

7. CONFERVA filamentis æqualibus ramosis, bullas aëreas includentibus. *bullosa.* *Sp. pl. 1164. Fl. suec. 1166.*

Conferva palustris botabycinia. R. Syn. 60. Dill. musc. 18. t. 3. f. 11.

Alga bombycinia. Bauh. pin. 363. pr. 155. Park. 1261.

Anglis, Cotten Conferva.

Habitat in fossis et aquis stagnantibus, passim.

- canaliculatis.* 8. CONFERVA filamentis æqualibus basin versus ramosioribus. *Sp. pl.* 1164. *Fl. suec.* 1162.
 Conferva rivulorum capillacea, densissime congestis ramulis. *R. Syn.* 58. *Dill. musc.* 21. t. 4. f. 15.
 Alga in tubulis aquam fontanam ducentibus. *B. pin.* 364.
Anglis, Mill Conferva.
 Habitat in rivulis purioribus, inque tubulis et afferibus aquas ad molas deducentibus.
- tomentosa.* 9. CONFERVA filamentis æqualibus ramosissimis, ramis simplicibus confertis fuscis.
 Conferva marina tomentosa, tenerior et albicans. *R. Syn.* 59. *Dill. musc.* 19. t. 3. f. 12.
 Muscus maritimus gossipio similis. *Bauh. pin.* 663. *Hist. ox.* III. 649.
 β Conferva marina tomentosa, minus tenera et ferruginea. *Dill. musc.* 19. t. 3. f. 13.
Anglis, Wool Conferva.
 Habitat in fossis et littoribus marinis passim.
- ærugino-fa.* 10. CONFERVA filamentis ramosis mollibus digitis brevioribus viridissimis. *Sp. pl.* 1165.
 Conferva marina capillacea brevis, viridissima mollis. *Dill. musc.* 23. t. 4. f. 20.
Anglis, Sea-green Conferva.
 Habitat in fucis prope Llanfaethly in Mona insula. *Dill.*
- spongiosa.* 11. CONFERVA filamentis ramosis æqualibus, ramulis simplicibus imbricatis.
 Fucus teretifolius spongiosus pilosissimus. *R. Syn.* 46.
 Fucus teres villis quaquaversum abductus. *R. Syn. ed.* II. 330.
 Muscus marinus hirsutus, flagellis longis ramosis subviridibus. *Hist. ox.* III. 650. f. 15. t. 9. f. 6. 7.
Anglis, Sponge Conferva.
 Habitat in littoribus marinis frequens.

12. CONFERVA filamentis æqualibus ramosis, ramis *nigra*.
 fasciculatis brevissimis.
Anglis, black Conferva.
 Habitat *in littore Eboracensi copiose*.
13. CONFERVA filamentis æqualibus dichotomis. *Sp. dichotoma*.
pl. 1165.
 Conferva dichotoma, setis porcinis similis. *Dill. musc.*
17. t. 3. f. 9.
 Conferva Plinii setis porcinis similis. *Merr. pin. R.*
Syn. 58.
Anglis, Bristle Conferva.
 Habitat *in fossis inter Greenwich et Woolwich. R.*
Syn.
14. CONFERVA filamentis proliferis fastigiatis hirtis. *scoparia*.
Sp. pl. 1156.
 Conferva marina pennata. *Dill. musc.* 24. *t. 4. f. 23.*
Anglis, Broom Conferva.
 Habitat *in littore marino inter Margate et Dover, in*
littore Devonix frequens.
15. CONFERVA filis ramosis : filamentis alternis bre- *cancellata*.
 vibus multipartitis digitatis. *Sp. pl.* 1165. *Fl. suec.*
1171.
 Conferva marina cancellata. *R. Syn.* 59. *Dill. musc.*
24. t. 4. f. 22.
Anglis, Latticed Conferva.
 Habitat *pone Sheerness in insula Sheepy. R. Syn.*
- ** *Filamentis anastomasantibus.*
16. CONFERVA filamentis reticulato-coadunatis. *Sp. reticulata*.
pl. 1165. *Fl. suec.* 1167.
 Conferva reticulata. *R. Syn.* 59. *Dill. musc.* 20. *t.*
4. f. 14.
 Conferva reticulata crispa. *Pluk. ph. t. 24. f. 2.*
Hist. ox. III. 644. t. 4. f. 4.
Anglis, Net Conferva.
 Habitat *in fossis prope Westmonasterium, et rivulis*
in ericeto Hounsflejano, D. Doody.

*** *Filamentis nodosis.*

fluviatilis. 17. CONFERVA filis simplicissimis setiformibus rectis : geniculis crassioribus angulatis. *Sp. pl.* 1165. *Fl. suec.* 1168.

Conferva fluviatilis lubrica setosa, Equiseti facie. *Dill. musc.* 39. t. 7. f. 47.

Anglis, Horse-tail Conferva.

Habitat prope Bangoriam in rivo aquæ dulcis, dicto Afon Cegin, inter molendinum vocatum Panda maes y Geirchan et Melun Escob lapidibus dense adnascentem invenit et misit. *Sam. Brewer. Dill. musc.*

gelatinosa. 18. CONFERVA filis ramosis moniliformibus, articulis globosis gelatinosis. *Sp. pl.* 1166.

Conferva fontana nodosa, spermatis ranarum instar lubrica, major et fusca. *R. Syn.* 62. *Dill. musc.* 36. t. 7. f. 42.

β Conferva fontana nodosa, spermatis ranarum instar lubrica, minor et viridis. *Dill. musc.* 37. t. 7. f. 43.

γ Conferva stagnatilis, globulis virescentibus mucosis. *Dill. musc.* 38. t. 7. f. 44.

δ Conferva alpina lubrica, filamentis nodosis cæruleis. *Dill. musc.* 38. t. 7. f. 45.

ε Conferva fontana nodosa lubrica, filamentis tenuissimis nigris. *Dill. musc.* 39. t. 7. f. 46.

Anglis, Frog-spawn Conferva.

Habitat in fonte amplo et limido vici Godalmin in Surreia, juxta viam regiam, et in fonte quodam prope Cicestriam in Sufflexia; β. in rivulo in saltu Enfieldensi; γ. δ. prope Mancunium; ε. in Mona insula. *Dill. in rivulo prope Martin in Surreia.*

***** *Filamentis geniculatis.*

capillaris. 19. CONFERVA filis geniculatis simplicibus, articulis alternatim compressis. *Sp. pl.* 1166.

Conferva filamentis longis geniculatis simplicibus. *Dill. musc.* 25. t. 5. f. 25.

Conferva palustris, s. *filum marinum anglicum*. R.
Syn. 60.

Conferva geniculata minima nostras. Pluk. ph. t.
84. f. 9. Hist. ox. III. 644. t. 4. f. 4.

Anglis, Thread *Conferva*.

Habitat in fossis et lucubus marinis.

20. *CONFERVA* filis geniculatis ramosis, articulis al- *tubulosa*,
ternatim compressis.

Conferva marina fistulosa. Dill. musc. 34. t. 6. f. 39.

Conferva marina geniculata albicans diaphragmatis
distincta. R. Syn. 60.

Conferva marina nodosa ex albo rubescens mollis, sed
minus lubrica. R. Syn. 51.

Tremella filiformis articulata: articulis alternatim
compressis. Sp. pl. 1163.

Anglis, Hollow *Conferva*.

Habitat in littore marino passim.

21. *CONFERVA* filamentis geniculatis ramosissimis, *pellucida*,
articulis cylindricis, ramis oppositis.

Anglis, transparent *Conferva*.

Habitat in littore insulæ Walney.

22. *CONFERVA* filamentis geniculatis subdichotomis, *polymor-*
ramis fasciculatis nigris. *pha.*

Conferva filamentis geniculatis, ramis fasciculatis. Sp.
pl. 1167. Fl. suec. 1174.

Conferva marina geniculata nigra palmata. Dill. musc.
32. t. 6. f. 35.

Conferva marina geniculata ramosissima lubrica, bre-
vibus et palmatim congestis ramulis. R. Syn. 61.

Musculus capillaceus multifidus niger. B. pin. 363.

Anglis, Palmated *Conferva*.

Habitat in littoribus marinis frequens.

23. *CONFERVA* filis geniculatis dichotomis, filamen- *glomerata*,
tis ramosis, ramulis brevibus fasciculatis scabris.

Conferva fontalis ramosissima, glomeratim congesta.

R. Syn. 59. Dill. musc. 28. t. 5. f. 31.

CRYPTOGAMIA ALGÆ.

Conferva filamentis geniculatis, ramulis brevioribus multifidis. *Sp. pl.* 1167.

Conferva viridis capillacea, brevioribus setis, ramifior, s. *conferva* minor ramosa. *Hist. ox. III.* 644. f. 15. t. 4. f. 2.

Anglis, Cluster *Conferva*.

Habitat in fluviis et littoribus marinis.

Confer. *Conferva* marina trichoides, ramosissima, sparsa. *Dill. musc.* 28. t. 5. f. 30. *R. Syn.* 60.

corallinoi- 24. *CONFERVA* filamentis geniculatis dichotomis.
des. *Sp. pl.* 1166.

Conferva marina gelatinosa, corallinae instar geniculata tenuior. *Dill. musc.* 33. t. 6. f. 37.

Corallina confervoides gelatinosa rubens, ramulis et geniculis perangustis. *R. Syn.* 34.

Muscus marinus capillaris rubens non ramosus. *Musc. Pet.* 274.

β *Conferva* marina gelatinosa, corallinae instar geniculata crassior. *Dill. musc.* 33. t. 5. f. 36.

Corallina confervoides geniculata alba, geniculis crassiusculis pellucidis. *R. Syn.* 34.

Anglis, Coralline *Conferva*.

Habitat in littoribus marinis passim.

elongata. 25. *CONFERVA* filamentis geniculatis ramosissimis, ramulis longissimis distantibus acutis.

Conferva marina geniculata ramosissima lubrica, longis sparsisve ramulis. *R. Syn.* 61. *Dill. musc.* 35. t. 6. f. 38.

Anglis, Pointed *Conferva*.

Habitat in littoribus marinis frequens.

fulva: 26. *CONFERVA* filamentis geniculatis ramosis, ramis ramulisque brevissimis alternis.

Anglis, short *Conferva*.

Habitat in littore Eboracensi.

plicata. 27. *CONFERVA* filis geniculatis ramosis, filamentis ramosissimis implexis.

Conferva

- Conferva marina capillacea longa ramofiffima, mollis. *R. Syn.* 59. *Dill. mufc.* 23. t. 6. f. 19.
Anglis, matted Conferva.
 Habitat in rupibus et faxis marinis paffim.
28. CONFERVA filamentis geniculatis flexuofis, ramis vagabundis ramulifque brevioribus. *Sp. pl.* 1167. *Fl. fuc.* 1177. *da.*
 Conferva marina trichoides, lanæ inflar expanfa. *R. Syn.* 60. *Dill. mufc.* 30. t. 5. f. 32.
Anglis, spreading Conferva.
 Habitat in paludibus maritimis.
29. CONFERVA filamentis geniculatis ramofis, ramis rupeflris alternis ramofis viridibus.
 Conferva filamentis geniculatis ramofiffimis viridibus. *Sp. pl.* 1167. *Fl. fuc.* 1176.
 Conferva marina trichodes ramofior. *Dill. mufc.* 28. t. 5. f. 29.
 Conferva marina trichodes, f. mufcus marinus virens tenuifolius. *Pluk. mant.* 53. t. 182. f. 6. *R. Syn.* 60.
 Conferva Auviatilis trichodes, extremitatibus ramofis. *Cat. giff.* 199.
Anglis, Rock Conferva, or Sea-Beard.
 Habitat in rupibus maritimis et in fluviis, paffim.
30. CONFERVA filamentis geniculatis ramofis ramis fasciculatis confertis viridibus. *sericea.*
 Conferva trichodes virgata, *sericea.* *Dill. mufc.* 31. t. 5. f. 33.
 Conferva marina trichodes, ramulis virgatis longioribus glabris. *R. Syn.* 60.
Anglis, Silk Conferva.
 Habitat in littore infulæ Shepey; in Rivo Novo prope Londinum. *Dill.*
31. CONFERVA filamentis geniculatis ramofiffimis ramulis multifidis fasciculatifque. *fucoides.*
Anglis, Branched Conferva.
 Habitat in littore Eboracenfi.

- fusca.* 32. CONFERVA filamentis geniculatis ramosissimis, ramulis alternis simplicibus.
Anglis, Brown Conferva.
 Habitat in littore Eboracensi.
- vubra.* 33. CONFERVA filamentis geniculatis ramosissimis compressis, ramis supradecompositis pinnatis coloratis.
 Fucus cartilagineus, stipite teretiusculo compresso: frondibus supradecompositis pinnatis: laciniis linearibus coloratis. *Roy. lugdb.* 515. *Sp. pl.* 1161.
 Muscus marinus folio multifido. *Johns. Merc. bot.* II. 27.
 Muscus pelagicus pennatus rubens, ramulis numerosis mollibus latius se spargentibus. *Pluk. ph. t.* 48. f. 2.
 Muscus marinus purpureus parvus, foliis oblongis millefolii fere divisura. *R. Hist.* I. 79.
Anglis, Red Conferva.
 Habitat in littoribus marinis frequens.
- pennata.* 34. CONFERVA filamentis geniculatis ramosis, ramis duplicato pinnatis fuscis.
 Conferva marina pennata. *Pet. hort. succ. R. Syn.* 59.
Anglis, Feathered Conferva.
 Habitat in littore marino inter Margate et Dover, *R. Syn.* In littore insula Walney.
- parasitica.* 35. CONFERVA filis geniculatis pinnatis.
Anglis, Winged Conferva.
 Habitat in fucis in littore Eboracensi passim.
- agagropia.* 36. CONFERVA filamentis geniculatis ramosissimis e centro confertissimis globum constituentibus. *Sp. pl.* 1167. *Fl. suec.* 1169.
 Conferva globosa. *Aët. Angl. Vol. XLI.* 498.
Anglis, Globe Conferva, or Moor Balls.
 Habitat in lacubus montosis, in comitatu Westmorlandico. In Wallingfen-Moor, Yorkshire. *Mr. Tho. Knowlton.*

B Y S S U S. 1071.

*Lanugo aut Pulvis simplicissimus.** *Filamentosæ.*

1. BYSSUS filamentis plumosis natantibus. *Sp. pl. Flos aquæ*
1168.

*Byssus latissima, papyri instar supra aquam expansa.**Dill. musc. 1. R. Syn. 57.**Anglis, Paper Byssus.**Habitat in aquis stagnantibus.*

2. BYSSUS lanuginosa violacea lignis adnascens. *R. phospho-*
Syn. 56. Dill. musc. 4. t. 1. f. 6. Sp. pl. 1168. rea.

*Anglis, Violet-coloured Byssus.**Habitat in lignis putrescentibus.*

3. BYSSUS filamentosa viridis. *Sp. pl. 1168. velutina.*

*Byssus tenerrima viridis velutum referens. R. Syn. 56.**Dill. musc. 7. t. 1. f. 14.**Anglis, Velvet Byssus.**Habitat in corticibus arborum et terris umbrosis.*

4. BYSSUS capillacea pulverulenta, fructificationibus aurea.
sparsis, filamentis simplicibus ramosisque. *Gleditsch.*
fung. 1. Sp. pl. 1168.

Byssus petræa crocea, glomerulis lanuginosis. Dill.
*musc. 8. t. 1. f. 16.**Byssus aurea derbiensis humifusus. Pet. gaz. 1. t. 13.*
*f. 3.**Anglis, Saffron-coloured Byssus.**Habitat in rupibus Arvonix passim. Dill.*

5. BYSSUS filamentis ramosis rigidis atris faxis adnas- *nigra.*
cens.

*Byssus petræa nigerrima fibrosa. R. Syn. 57.**Anglis, Black Byssus.**Habitat in rupibus calcariis prope Ingleborough et*
Settle in comitatu Eboracensi.

6. BYSSUS filamentis ramosis fulvis.

fulva.

Byffus arborea crocea fibrosa. *R. Syn. 57. Dill. musc. 9. t. 1. f. 17.*
Anglis, Brown Byffus.
Habitat in lignis putridis.

barbata. 7. BYSSUS? filamentis erectis, summitatibus ramosis.
 Byffus arborea barbata, fulvi coloris. *R. Syn. 57. Dill. musc. 9. t. 1. f. 19.*
 Pseudo-spongia fungoides fulva lignis adnascens. *R. Syn. ed. II. 334.*
Anglis, Bearded Byffus.
Habitat in lignis putrescentibus.

candida. 8. BYSSUS filamentis ramosissimis ramis fasciculatis albidis.
 Byffus tenerrime villosa et elegantissime ramulosa. *R. Syn. 476. t. 23. Dill. musc. 7. t. 1. f. 15.*
Anglis, Grey Byffus.
Habitat in lignis et foliis putrescentibus.

cryptarum. 9. BYSSUS capillacea perennis cinerea tenax rupi innata. *Fl. lapp. 527. succ. Sp. pl. 1168.*
 Byffus albida brevis setacea. *Dill. musc. 10. t. 1. f. 20.*
Anglis, Bristle Byffus.
Habitat in crypta subteranea in comitatu Herefordiensis. Dill.

** *Pulverulentæ.*

antiquitatis. 10. BYSSUS pulverulenta atra. *It. gotl. 298. Sp. pl. 1168.*
Anglis, Black Powder Byffus.
Habitat in muris antiquissimis frequens.

saxatilis. 11. BYSSUS pulverulenta cinerea rupes operiens. *Sp. pl. 1169.*
Anglis, Stone Byffus.
Habitat in rupibus calcariis in comitatibus Westmorlandico et Eboracensi.

12. BYSSUS pulverulenta flava lignis adnascens. *R. candelaris.*
Syn. 56. Dill. musc. 3. t. 1. f. 4. Sp. pl. 1169.
Anglis, Yellow Byssus.
Habitat in corticibus arborum.
13. BYSSUS pulverulenta viridis. *Sp. pl. 1169. botryoides.*
Byssus botryoides saturate virens. R. Syn. 56. Dill.
musc. 3. t. 1. f. 5.
Anglis, Green Byssus.
Habitat in cortice arborum et terra humida, umbrosa.
14. BYSSUS pulverulenta incana, farinæ instar strata. *incana.*
R. Syn. 56. Dill. musc. 3. t. 1. f. 3. Sp. pl. 1169.
Anglis, Mealy Byssus.
Habitat in solo glareoso, ad latera fossularum, juxta
vias publicas.
15. BYSSUS pulverulenta-crustacea albissima. *Sp. pl. lactea.*
1169.
Byssus candidissima, calcis instar Muscos vestiens.
Dill. musc. 2. t. 1. f. 2.
Anglis, White Byssus.
Habitat in muscis et arborum corticibus.

S P O N G I A. 1072.

Pili intricati intertexti.

* *Tenaces.*

1. SPONGIA tenax ramosissima erecta, ramis teretiusculata.
culis obtusis. Sp. pl. 1170.
Spongia ramosa. Bauh. pin. 368. R. Syn. 29, Bri-
tannica. Park. 1304.
Anglis, Branched Spunge.
Habitat in littoribus marinis passim.
2. SPONGIA erecta dichotoma compressa. *dichotoma.*
Spongia dichotomos teretifolia viridis. R. Syn. 29.
Fucus spongiosus teres ramosior viridis erectis. Hist.
ox. III. 647. t. 8. f. 7.

Fucus

CRYPTOGAMIA FUNGI.

Fucus filiformis compressus dichotomus rectus : geniculis tumidiusculis. *Sp. pl.* 1159.

Anglis, forked Sponge.

Habitat in littoribus marinis. In Cornubia. *R. Syn.*
in littore Eboracensi, sed varius.

* *Fragiles.*

lacustris. 3. SPONGIA repens fragilis, ramis teretibus obtusis.
Sp. pl. 1171. *Fl. suec.* 1190.

Spongia ramosa fluviatilis. *R. Syn.* 30.

Anglis, creeping sponge.

Habitat in fundo lacuum. In comitatu Westmorlandico frequens.

fluviatis. 4. SPONGIA erecta fragilis, ramis teretiusculis. *Sp.*
pl. 1171. *Fl. suec.* 1191.

Spongia fluviatilis ramosa fragilis. *R. Syn.* 30.

Spongia fluviatilis anfractuosa perfragilis ramosissima.

Pluk. ph. t. 112. *f.* 3.

Anglis, River Sponge.

Habitat in fluviis. In the river Thames by Swythens Wyars, near Oxford. In fluvio Camo inter Cantabrigiam et Chesterton. In the river by Waltham Abby, Essex. *R. Syn.*

F U N G I.

A G A R I C U S. 1074.

Fungus horizontalis, subtus lamellosus.

* *Stipitati, pileo orbiculato.*

Chantarellus. 1. AGARICUS stipitatus, lamellis ramosis decurrentibus. *Sp. pl.* 1171. *Fl. suec.* 1207.

Fungus angulosus et velut in lacinias sectus. *Bauh. pin.* 371. *Vaill. paris. t.* 11. *f.* 14, 15.

Fungus luteus f. pallidus, Chanterelle dictus, se contorquens esculentus. *Bauh. hist. III.* 832. *R. Syn.* 2.

Fungi

Fungi lethales. *Ger. Em.* 1385.

Anglis, yellow Agaric, or Champignon.

Habitat in pratis, pascuis et sylvis.

2. AGARICUS stipitatus, lamellis omnibus magnitudi-integer.
ne æqualibus. *Fl. suec.* 1230. *Sp. pl.* 1171.

Agaricus caulescens, pileo plano-concavo purpureo,
lamellis stipiteque albis. *Fl. lapp.* 486.

Fungus major pediculo longo, modice crasso, lamel-
lis albis creberrimis, superne ad margines apparen-
tibus. *R. Syn.* 4.

Anglis, white Agaric.

Habitat in sylvis.

3. AGARICUS stipitatus, lamellis dimidiatis solitariis, *muscarius*.
stipite volvato : apice dilatato, basi ovato. *Fl. suec.*
1236. *Sp. pl.* 1172.

Fungus pileo sanguineo verrucoso, lamellis albis, an-
nulo, fugaci, pediculo bulbofo. *Hall. helv.* 39.

Anglis, reddish Agaric.

Habitat in pratis inter Hampstead et Highgate, sed
rarius.

4. AGARICUS stipitatus, pileo plano carneo lactes-lactifluus.
cente, lamellis rufis, stipite longo carneo. *Sp. pl.*
1172.

Fungus lactescens non acris. *R. Syn.* 4.

Anglis, Milky Agaric.

Habitat in sylvis. In sylva Charlton Wood dicta.

5. AGARICUS stipitatus, pileo planiusculo lactescente : piperatus.
margine deflexo, lamellis incarnato-pallidis. *Sp. pl.*
1173. *Fl. suec.* 1195.

Fungus piperatus albus lacteo succo turgens. *Bauh.*
hist. III. 825. *R. Syn.* 4. *Mich. gen.* 141. *Hall.*
helv. 34.

Fungus albus acris. *Bauh. pin.* 370.

Anglis, Pepper Agaric.

Habitat in sylvis. In sylva Charlton Wood dicta
copiose.

- campestris.* 6. AGARICUS stipitatus, pileo convexo squamato al-bido, lamellis rufis. *Sp. pl.* 1173. *Fl. suec.* 1203.
Fungus campestris albus superne, inferne rubens.
Bauh. hist. III. 824. *R. Syn.* 2. *Mich. gen.* 174.
Fungus esculentus. *Park.* 1317.
Anglis, Field Agaric, or common Mushroom.
Habitat in pratis et pascuis.
- violaceus.* 7. AGARICUS stipitatus, pileo rimoso : margine vio-laceo tomentoso, stipite cærulecente lana ferrugi-neâ. *Sp. pl.* 1173. *Fl. suec.* 1226.
Fungus pileo magno, orbiculari, sublivido, lamellis albis, pediculo brevi bulbiformi violaceo. *R. Syn.* 3.
Fungus esculentus bulbosus dilute purpureus. *Mich. gen.* 149 t. 49. f. 1. *Hall. helv.* 52.
Anglis, Violet Agaric.
Habitat in pratis circa Newington. *Dill. R. Syn.* XII.
- cinnamo-meus.* 8. AGARICUS stipitatus, pileo fordide flavo, lamellis luteo-rufis. *Sp. pl.* 1173. *Fl. suec.* 1205.
Fungus ovinus. *Sterb.* 64 t. 4. *D. E. F. R. Syn.* 2.
Fungus planus orbiculatus aureus. *B. pin.* 371.
Anglis, brown Agaric.
Habitat in sylvis passim.
- viscidus.* 9. AGARICUS stipitatus, pileo purpurascente-fusco vis-cido, lamellis fusco-purpurascens. *Sp. pl.* 1173.
Fl. suec. 1229.
Agaricus caulescens, pileo convexo purpurascens.
lamellis rufis, stipite brevi crassiusculo albo. *Fl. lapp.* 485.
Amanita dura, ex fusco-rubens quercina. *Dill. cat. giff. R. Syn.* 5.
Anglis, clammy Agaric.
Habitat in sylvis.
- equestris.* 10. AGARICUS stipitatus, pileo pallido : disco stellatim luteo, lamellis sulphureis. *Sp. pl.* 1173. *Fl. suec.* 1219.

Agaricus caulescens, pileo convexo fordido: margine inflexo, lamellis pallidis: basi remotis, stipite cylindraceo. *Fl. lapp.* 502.

Fungus parvus parvi galeri formam exprimens rufus.

Bauh. pin. 373. *R. Syn.* 7.

Anglis, Starry Agaric.

Habitat in ericetis passim.

11. AGARICUS stipitatus, pileo convexo acuminato mammo-griseo, lamellis convexis griseis crenatis, stipite nudus. *Sp. pl.* 1174. *Fl. suec.* 1218.

Fungus minor, capitulo pileum nauticum referente, pediculo longiore. *Buxb. cent.* IV. 13. t. 21. f. 1.

Fungus fordide fulvus capitulo in conum fastigiato, pediculo longissimo firmo striato. *R. Syn.* 4.

Anglis, grey Agaric.

Habitat in sylvis.

12. AGARICUS stipitatus, pileo hemisphærico viscido clypeatus. acuminato, lamellis albis, stipite longo cylindrico albo. *Fl. lapp.* 506. *suec.* 1216. *Sp. pl.* 1174.

Fungus major, pediculo longo, modice crasso, lamellis albis creberrimis, superne ad margines apparentibus. *R. Syn.* 4.

Anglis, long-stalked Agaric.

Habitat in sylvis.

13. AGARICUS stipitatus pileo campaniformi albido extincto-lacero, lamellis niveis, stipite subbulboso subulato rufus. nudo. *Sp. pl.* 1174. *Fl. suec.* 1196.

Anglis, Conic Agaric.

Habitat in pratis prope Tottenham.

14. AGARICUS stipitatus, pileo campanulato lacero, fimetari-lamellis nigris lateraliter flexuosis, stipite fistuloso. *Sp. pl.* 1174. *Fl. suec.* 1215.

Fungus albus ovum referens. *R. Syn.* 5. *Buxb. cent.* IV. t. 27. f. 1.

Anglis, Egg Agaric.

Habitat in fimetis frequens.

15. AGA-

- verrucosus* 15. AGARICUS caulescens, pileo convexo cinereo :
verrucis lamellisq̃ue albis, stipite basi crassiusculo :
Fl. lapp 512. *suéc* 1236.
Fungus pileo plano ad oras striato sordido verrucoso,
subtus albus annulatus. *Hall. helv* 43.
Fungi albi venenati viscid. *B. hist. III.* 826. *Sterb.*
fung. 206 t. 2. f. *K. R. Syn.* 7.
Anglis, Warty Agaric.
Habitat in nemorosis.
- campanu-* 16. AGARICUS stipitatus, pileo campanulato striato
latus. pellucido, lamellis adscendentibus, stipite nudo.
Sp. pl. 1175. *Fl. suéc.* 1213.
Fungus perpusillus, pediculo oblongo, pileo tenui
utrinque striato, seu flabelli in modum plicatili. *R.*
Syn. 8.
Anglis, Bell Agaric.
Habitat in pratis et pascuis.
- separatus.* 17. AGARICUS stipitatus, pileo lævi livido, lamellis
nigricantibus separatis, stipite bulboso volvato.
Sp. pl. 1175. *Fl. suéc.* 1220.
Fungus parvus pediculo oblongo, galericulatus, striis
lividis aut nigris. *R. Syn.* 8.
Anglis, blackish Agaric.
Habitat in stercoreis.
- fragilis.* 18. AGARICUS stipitatus, pileo convexo viscido pel-
lucido lamellisq̃ue luteis, stipite nudo. *Sp. pl.*
1175.
Fungus pileo croceo splendoris particeps. *Vaill. pa-*
ris. t. 11. f. 16. 17. 18.
Fungus pratensis minor, externe viscidus, capitulo
præcedentis, striis subtus fulvis s. croceis. *R.*
Syn. 8.
Anglis, brittle Agaric.
Habitat in ericetis.
- umbellife-* 19. AGARICUS stipitatus, pileo plicato membranaceo,
rus. lamellis basi latioribus. *Sp. pl.* 1175. *Fl. suéc.*
1192.

Fungus

Fungus parvus candidissimus lamellatus, pediculo longo gracili. *R. Syn.* 9.

Anglis, Wood Agaric.

Habitat inter semiputrida dejecta folia sylvarum.

20. AGARICUS stipitatus albus, pileo plicato membra- *Androsana-*
naceo, stipite nigro. *Sp. pl.* 1175. *Fl. suec.* 1193. *ceus.*

Fungus minimus adianti aurei capitulis. *R. Syn.* 9.

Fungus caule nigro capillari, androsaces capitulo.

Bocc. musc. 143. *t.* 104.

Anglis, Black stalked Agaric.

Habitat in lignis et foliis putridis. On rotten Sticks in the Bottoms of old Lakes. *R. Syn.*

* *Parasiti acaules, dimidiati.*

21. AGARICUS acaulis, lamellis labyrinthiformibus. *quercinus.*
Sp. pl. 1176.

Agaricus quernus lamellatis coriaceus villosus. *Cat.*

giff. R. Syn. 25.

Anglis, Oak Agaric.

Habitat in truncis arborum.

22. AGARICUS acaulis coriaceus, lamellis sinuatis *pectina-*
simplicibus. *tus.*

Agaricus coriaceus longissimus, pectinatim inferne
divisus. *R. Syn.* 25.

Anglis, Toothed Agaric.

Habitat in lignis putridis.

23. AGARICUS acaulis coriaceus villosus : margine *Betulinus.*
obtus, lamellis ramosis anastomosantibus. *Sp. pl.*

1176. *Fl. suec.* 1244.

Agaricus villosus, lamellis sinuosis et invicem im-
plexis. *Cat. giff. R. Syn.* 24.

Anglis, Birch Agaric.

Habitat in truncos arborum.

24. AGARICUS acaulis, lamellis bifidis pulverulentis. *alneus.*
Sp. pl. 1176. *Fl. suec.* 1242.

Agar-

- Agaricus parvus lamellatus pectunculi forma elegans.
Cat. giff. R. Syn. 25.
Anglis, Alder Agaric.
 Habitat *ad truncos arborum. In the woods near Dul-*
wich. R. Syn.

B O L E T U S. 1075.

Fungus horizontalis, subtus porosus.

* *Parasitici acaules.*

- suberosus.* 1. BOLETUS acaulis pulvinatus albus lævis, poris acutis difformibus. *Sp. pl. 1176. Fl. suec. 1253.*
 Agaricus intibaceus. *Tourn. 562. R. Syn. 23.*
 β Fungus foraminosus arboreus lævis albissimus. *R. Syn. ed. II. 340.*
Anglis, Cork Boletus.
 Habitat *in Fraxinis in comitatu Westmorlandico frequens.*
- fomentarius.* 2. BOLETUS acaulis pulvinatus inæqualis obtusus, poris teretibus æqualibus glaucis. *Sp. pl. 1176. Fl. suec. 1252.*
 Agaricus porosus igniarius Carpini. *Cat. giff. R. Syn. 24.*
Anglis, Spongy Boletus.
 Habitat *in caudicibus arborum.*
- versicolor.* 3. BOLETUS acaulis, fasciis dicoloribus, poris albis. *Sp. pl. 1176. Fl. suec. 1254.*
 Agaricus varii coloris squamosus. *Tourn. 562. R. Syn. 24.*
 Fungus arborum et lignorum putrescentium, coloris varii. *R. Syn. ed. II. 18.*
Anglis, striped Boletus.
 Habitat *in truncis arborum.*
- albus.* 4. BOLETUS acaulis pulvinatus lævis, poris oblongis. Agaricus officinali similis. *Cat. giff. R. Syn. 23.*
 Fungus arboreus albidus maximus, f. Agaricus spurius. *R. Syn. ed. II. 335.*
Anglis, white Boletus.
 Habitat *in salicibus.*

5. BOLETUS acaulis pulvinatus lævis, poris tenuissimis. *Fl. lapp.* 582. *suec.* 1256. *Sp. pl.* 1176.
 Agaricus pedis equini facie. *Tourn.* 562. *R. Syn.* 22.
 Fungus durus five igniarius. *Park.* 1323.
Anglis, hard Boletus, or Touchwood.
 Habitat in arborum truncjs.

** Stipitatis.

6. BOLETUS stipitatus, pileo glabro pulvinato marginato, poris compositis acutis: porulis angulatis brevioribus. *Sp. pl.* 1177. *Fl. suec.* 1246.
 Suillus fulvus inferne ex flavo virescens. *Hall. helv.* 29.
 Fungus porosus crassus. *R. Syn.* 11.
Anglis, brown Boletus.
 Habitat in sylvis.
7. BOLETUS stipitatus, pileo pulvinato subvilcido, poris rotundatis convexis flavissimis, stipite albo. *Sp. pl.* 1177. *Fl. suec.* 1247.
 Boletus luteus. *Cat. giff.* *R. Syn.* 10.
 Suillus annulatus terreus inferne flavescens. *Hall. helv.* 29.
Anglis, yellow Boletus.
 Habitat in sylvis passim.

HYDNUM. 1076.

Fungus horizontalis, subtus echinatus.

1. HYDNUM stipitatum, pileo convexo imbricato. *imbricatum.*
Sp. pl. 1178. *Fl. lapp.* 523. *suec.* 1257.
 Erinaceus esculentus albus crassus. *Mich. gen.* 132.
 t. 72. f. 2. *Hall. helv.* 31.
 Fungus pæne candidus, prona parte erinaceus. *Bauh. hist.* III. 828. *R. Syn.* 11.
Anglis, common Hydnum.
 Habitat in sylvis. In sylva quadam prope Middleton vicum in agro Warwicensi. *R. Syn.*

P H A L L U S. 1077.

*Fungus supra reticulatus, subtus lævis.**esculentus.* 1. PHALLUS pileo ovato, stipite nudo rugoso. *Fl. suec.* 1262. *Sp. pl.* 1178.Boletus esculentus rugosus. 1. 2. 3. *Mich. gen.* 203. t. 85. f. 1. 2.Fungus favaginosus. *Park.* 1317. *R. Syn.* 11.*Anglis*, Esculent Morel.

Habitat in sylvis et nemorosis.

impudicus. 2. PHALLUS volvatus stipitatus, pileo celluloso. *Sp. pl.* 1179. *Fl. suec.* 1261.Phallus vulgaris totus albus. *Mich. gen.* 201. t. 83.Phallus hollandicus. *Park.* 1322.Fungus Phalloides. *B. hist. Ill.* 843. *R. Syn.* 12.Fungus foetidus penis imaginem referens. *Bauh. pin.* 374.*Anglis*, Stinking Morel.

Habitat in sylvis et ad sepes sed rarius.

E L V E L A. 1079.

*Fungus turbinatus, lævis.**mitra.* 1. ELVELA pileo deflexo adnato lobato difformi. *Fl. suec.* 1265. *Sp. pl.* 1180.Fungoides fungiforme crispum laciniatum et varie complicatum, pediculo crasso striato rimoso et fistuloso. *Mich. gen.* 204. t. 86. f. 7.Fungus terrestris pediculo striato et cavernoso, capitulo plicatili subtus plano. *R. Syn.* 8.*Anglis*, curled Elvela.

Habitat in truncis putridis sed rarius.

P E Z I Z A. 1080.

*Fungus campanulatus, sessilis.**lentifera.* 1. PEZIZA campanulata lentifera. *Sp. pl.* 1180. *Fl. suec.* 1271.

Fung

Fungi calyciformes seminiferi. *Mentz. pug. t. 6.*
R. Syn. 20.

Fungus campaniformis niger parvus, multa semina
 in se continens. *Merr. pin. Pluk. ph. t. 184. f. 9.*
Anglis, black Peziza.

Habitat in lignis, agris, arvis hortorum passim.

2. PEZIZA turbinata truncata, disco punctato. *Fl. suec. punctata,*
1275. Sp. pl. 1180.

Fungus minimus infundibuliformis, superne nigris
 punctis notatus. *R. Syn. 12.*

Fungus minimus lignosus, disco punctato. *Bocc. mus.*
11. t. 107.

Anglis, spotted Peziza.

Habitat in stercore equino.

3. PEZIZA infundibuliformis, disco patente sinuato *cornucopia-*
 punctato. *Sp. pl. 1181. oides.*

Peziza tubæ fallopianæ æmula. *Cat. giff. R. Syn.*
20.

Fungoides nigricans majus cornucopiæ forma. *Vaill.*
paris. t. 13. f. 2. 3.

Anglis, Cornucopia Peziza.

Habitat in sylvis; in sylva Hornsey Wood dicta *co-*
piose.

4. PEZIZA cyathiformis, extus angulata venis ramosis. *acetabul-*
Sp. pl. 1181. lum.

Fungoides fuscum acetabuliforme, externe ramifica-
 tum. *Vaill. paris. t. 13. f. 1.*

Peziza subfusca major. *Cat. giff. R. Syn. 17.*

Anglis, Cup Peziza.

Habitat in lignis putridis. *Inter Hampstead et High-*
gate.

5. PEZIZA cyathiformis: margine obtuso erecto. *Fl. cyathoides,*
suec. 1274. Sp. pl. 1181.

Peziza lutea parva: marginibus lævibus. *R. Syn. 18.*
t. 24. f. 4.

Anglis, Smooth Peziza.

Habitat in sylvis. *In Charlton Wood. R. Syn.*

- scutellata*. 6. PEZIZA plana : margine convexo piloso. *Sp. pl.*
1181. *Fl. succ.* 1272.
Peziza lutea parva, marginibus pilosis. *R. Syn.* 18.
t. 24. *f.* 3
Anglis, Hairy Peziza.
Habitat in lignis putridis. In a lane near Chichester,
Southgate. *Dill. R. Syn.*

CLAVARIA. 1081.

Fungus lævis, oblongus.

* *Indivisæ.*

- pistillaris*. 1. CLAVARIA clavæformis simplicissima. *Fl. succ.*
1266. *Sp. pl.* 1182.
Clavaria alba pistilli forma. *Vaill. parif.* 39. *t.* 8.
f. 5.
Fungoides clavatum minus. *Cat. giff. R. Syn.* 14.
Anglis, simple Clavaria.
Habitat in sylvis et ericetis passim.

- ophioglossoides*. 2. CLAVARIA clavata integerrima compressa obtusa.
Guet. stamp. 1. *p.* 8. *Dalib. parif.* 385. *Sp. pl.*
1182.
Clavaria ophioglossoides nigra. *Vaill. parif.* *t.* 7. *f.* 4.
Fungus ophioglossoides. *Pb. br. R. Syn.* 14.
Anglis, black Clavaria.
Habitat in sylvis.

** *Ramosæ.*

- digitata*. 3. CLAVARIA ramosa lignea nigra. *Sp. pl.* 1182.
Fungus piperi æthiopico similis, vel digitatus niger.
Merr. pin. R. Syn. 14.
Lythophyton terrestre digitatum nigrum. *Comm. Ac.*
R. Sc. Ann. 1711.
Anglis, Fingered Clavaria.
Habitat in lignis semiputridis.

- Hypoxylo-*
lon. 4. CLAVARIA ramofo-cornuta compressa. *Sp. pl.*
1182. *Fl. succ.* 1267.

Fungus

- Fungus ramosus, niger, compressus, parvus, apicibus albidis. *R. Syn.* 15.
- * Fungus cornu dorcadis facie. *E. N. C. Dec.* 1. *Ann.* 4. 195.
 Agaricus ramosus cornu rangiferi referens. *D. Mill. Blackf. sp. bot.* 2. t. 1.
Anglis, Flat Clavaria.
 Habitat in lignis putridis.
5. CLAVARIA ramis confertis ramosissimis inæquali- *coralloides*
 bus *Sp. pl.* 1182. *Fl. suec.* 1268.
 Corallo-fungus flavus. *Vaill. parif. t.* 8. f. 4.
 Fungus ramosus flavus et albidus. *Bauh. hist.* III. 837. *R. Syn.* 16.
 Barba caprina. *Sterb. fung.* 96. 97. t. 11. f. A. B. C. D.
Anglis, yellow Clavaria
 Habitat in ericetis et sylvis.
6. CLAVARIA ramis confertis ramosissimis fastigiatis. *fastigiata.*
Sp. pl. 1183. *Fl. suec.* 1269.
 Fungoides fungiforme luteum foetidum et minus ramosum. *R. Syn.* 479. t. 24. f. 5.
Anglis, stinking Clavaria.
 Habitat in sylvis.
7. CLAVARIA ramis ramosis acuminatis inæqualibus *muscoides*,
 luteis. *Sp. pl.* 1183. *Fl. suec.* 1270.
 Fungus parvus ramosus luteus. *R. Syn.* 16. t. 24. f. 7.
Anglis, Pointed Clavaria.
 Habitat in sylvis et ericetis.

LYCOPERDON. 1082.

Fungus subrotundus feminibus farinaceis repletis.

* *Solida, subterranea absque radice.*

- * 1. LYCOPERDON solidum muricatum, radice de- *Tuber.*
 stitutum. *Sp. pl.* 1183. *Fl. suec.* 1282.
 Tuber brumale, pulpa obscura odorata. *Mich. gen.* 221. t. 164.

CRYPTOGAMIA FUNGI.

Tubera. *Baub. pin.* 376. *hist. III.* 849. *R. Syn.* 28.
terræ. *Ger. Em.* 1385. *edulia. Park.* 1319.

Anglis, solid Puff-balls, or Truffles.

Habitat *sub terra*; in Rushton Park, Northamptonshire. *R. Syn.*

cervinum. 2. LYCOPERDON solidiusculum centro farinifero, radice destitutum. *Sp. pl.* 1183. *Fl. suec.* 1276.

Lycoperdastrium tuberosum arrhizon fulvum, cortice duriore crasso et granulato, medulla ex albo purpurascens, semine nigro crassiore. *Mich. gen.* 220. *t.* 99. *f.* 4.

Tubera perniciofa terrestria s. cervina. *Sterb. fung.* 315. *t.* 32. *f.* B. B. B. *R. Syn.* 28.

Tubera cervina. *Baub. pin.* 376.

Anglis, Branny Puff-ball.

Habitat *in sylvis.* In Canewood, Hampstead, *R. Syn.*

**** Pulverulenta, radicata supra terram.**

Bruxia. 3. LYCOPERDON subrotundum, lacerato dehiscens. *Sp. pl.* 1183. *Fl. lapp.* 525. *suec.* 1280.

Lycoperdon vulgare. *Tourn.* 563.

Fungus maximus rotundus pulverulentus, dictus germanis Bofist. *Baub. hist. III.* 848.

Anglis, common Puff-ball.

Habitat *in pratis et pascuis.*

stellatum. 4. LYCOPERDON volva multifida patente, capitulo glabro: ore acuminato dentato. *Sp. pl.* 1184. *Fl. suec.* 1277.

Fungus pulverulentus, crepitus lupi dictus, coronatus et inferne stellatus. *R. Syn.* 27. *t.* 1. *f.* 1.

Anglis, Star Puff-ball.

Habitat *in pratis et pascuis.*

fornicatum. 5. LYCOPERDON volva quadrifida erecta fornicata, capitulo glabro; ore obtuso ciliato.

Geaster volvæ radiis et operculo elevatis. *D. Watson Aët. Angl.* No. 474. *p.* 234.

Fungus

Fungus pulverulentus turriculam fornicatam referens.

D. Rand. Blackf. spec. bot. 24. t. 2.

Anglis, Turret Puff-ball.

Habitat in pratis et pascuis sed rarius. At Bucklebury, ten miles from Reading, Berkshire, and about Wickham, near Bromley in Kent. *Blackf. spec. bot.*

6. LYCOPERDON stipite longo, capitulo globoso pedunculabro: ore cylindrico integerrimo. *Sp. pl. 1184. latum.*

Fl. suec. 1276.

Lycoperdon parisiense minimum pediculo donatum.

Tourn. 563. t. 331. f. E. F.

Fungus pulverulentus minimus, pediculo longo infidens. *R. Syn. 27.*

Anglis, Stalked Puff-ball.

Habitat in pascuis frequens.

MUCOR. 1083.

Fungus vesiculis stipitatis.

* *Perennes.*

1. MUCOR perennis, stipite filiformi nigro, capitulo sphaerocephalo globofo cinereo. *Fl. suec. 1286. Sp. pl. 1185. phalus.*

Sphaerocephalus niger, villo ochroleuco. *Hall. helv.*

III. t. 1. f. 3.

Fungus pileatus minimus, pediculo tenui capillaceo.

R. Syn. 12.

Anglis, Round-headed Mucor.

Habitat in lignis putridis frequens.

2. MUCOR perennis stipite subulato nigro, capitulo Lichenolenticulari cinereo. *Fl. suec. 1287. Sp. pl. 1185. des.*

Coralloides fungiforme arboreum nigrum vix crustosum. Dill. musc. 78. t. 14. f. 3.

Anglis, Grey-headed Mucor.

Habitat in lignis putridis.

3. MUCOR seta nigra, villo fusco. *Sp. pl. 1185. Fl. Embolus. suec. 1288.*

Embolus nigerrimus, villo albo adspersus. *Hall. helv.*
8. t. 1. f. 1.
Anglis, black Mucor.
Habitat in lignis putridis.

** *Fugaces.*

Mucedo. 4. *MUCOR* stipitatus, capsula globosa. *Sp. pl.* 1185.
Fl. succ. 1284.
Mucor vulgaris capitulo lucido per maturitatem nigro, pediculo griseo. *Mich. gen.* 215. t. 95. f. 1.
Hall. helv. 7.
- *Fungus bombycinus murini coloris*, e fimo felino, tenuissimis capillis. *Pluk. alm.* 164. *R. Syn.* 13.
Anglis, Cotten Mucor.
Habitat in variis putridis; pane, plantis, &c.

leprosus. 5. *MUCOR* setaceus, seminibus radicalibus. *Sp. pl.* 1185.
Aspergillus cespitosus densissimus, initio niveus, deinde aureus, seminibus ovatis. *Mich. gen.* 213. t. 91. f. 5.
Anglis, yellowish Mucor.
Habitat in cellulis fornicatis et cavernis.

glaucus. 6. *MUCOR* stipitatus capitulo subrotundo, aggregato. *Sp. pl.* 1186.
Aspergillus capitatus, capitulo glauco, seminibus rotundis. *Mich. gen.* 212. t. 91. f. 1.
Anglis, greenish Mucor.
Habitat in Pomis, Melonibus, aliisque corruptis.

crustaceus. 7. *MUCOR* stipitatus, spicis digitatus. *Sp. pl.* 1186.
Mucor crustaceus albus, capitulis fetisque aqueis. *Fl. succ. ed. I.* 1118.
Botrytis non ramosa alba, seminibus rotundis. *Mich. gen.* 212. t. 91. f. 3.
Anglis, Fingered Mucor.
Habitat in cibis corruptis.





A P P E N D I X.

O M I S S A.

V E R B E N A.

Cor. infundibuliformis subæqualis, curva. *Calycis* unico dente truncato. *Semina* quatuor nuda. *Stamina* quatuor.

1. VERBENA tetrandra, spicis filiformibus paniculatis, *officinalis*, foliis multifido-lanceolatis, caule solitario. *Sp. pl.* 20. *Fl. suec.* 30.

Verbena vulgaris. *Bauh. hist. III.* 443. *R. Syn.* 236.

Verbena mas, f. recta et vulgaris. *Park.* 674.

Verbena communis *Ger. Em.* 664, cæruleo flore. *Bauh. pin.* 269.

Anglis, Vervain.

Habitat in arvis et ad vias publicas.

Pharmac. Verbenæ Herba.

I L E X.

Hermaph. *Cal.* quadridentatus. *Cor.* quadripartite rotata. *Stam.* quatuor. *Stylus* o. *Stig.* quatuor. *Bacca* tetrasperma.

Masc. *Cal.* *Cor.* *Stam.* prioris.

1. ILEX foliis ovatis acutis spinosis. *Hort. cliff.* 40. *aquifolium*. *Sp. pl.* 125.

Ilex aculeata baccifera. *Bauh. pin.* 425.

Agrifolium. *Ger. Em.* 1338. f. Aquifolium. *Park.* 1486. *R. Syn.* 466.

Anglis, Holly-tree.

Habitat in sylvis, ericetis et sepibus. h.

- Alnus.* 2. BETULA pedunculis ramosis. *Sp. pl.* 983. *Fl. suec.* 861.
Alnus. *Ger. Em.* 1249. *R. Syn.* 442. vulgaris. *Park.* 1408.
Alnus rotundifolia glutinosa viridis. *Bauh. pin.* 428.
 β *Alnus vulgaris*, sub-conis ligulis membranaceis rubris donata. *R. Syn.* 442.
Anglis, Common Alder.
 Habitat in aquosis. h. VII.



I N D E X

NOMINUM LATINORUM

Generica nomina Romana Characterē traduntur :

Curſivis Literis Synonyma.

Primus numerus Genus designat ; cæteri Synonyma.

A		Page		Page
ABIES	<i>Abrotanum</i>	361	<i>Agrostis</i>	26—28
	<i>Absinthium</i>	311, 312	<i>Aira</i>	28—31, 374
	<i>Acanthium</i>	308	<i>Ajuga</i>	219
	<i>Acer</i>	378, 379	<i>Aizoon</i>	170, 172
	<i>Acetosa</i>	135, 136	<i>Alcea</i>	269
	<i>Acetosella</i>	136	<i>Alchemilla</i>	59, 60, 160
	<i>Achillea</i>	324, 325	<i>Alga</i>	342, 473, 475, 479
	<i>Acinos</i>	228, 230	<i>Alisma</i>	137, 138
	<i>Aconitum</i>	201	<i>Alliaria</i>	251
	<i>Acorus</i>	128, 13	<i>Allium</i>	120—122
	<i>Acrostichum</i>	383, 384	<i>Alnus</i>	83, 505
	<i>Actæa</i>	201	<i>Aloe</i>	208
	<i>Adiantum</i>	392, 383, 386, 387, 390	<i>Alopecurus</i>	23, 24, 47
	<i>Adonis</i>	209	<i>Alfinanthemos</i>	139
	<i>Adoxa</i>	150	<i>Alfinastrum</i>	151
	<i>Ægilops</i>	41	<i>Alfina</i>	113, 2, 6, 7, 11, 48, 51, 64, 120, 128, 139, 163, 166—169, 175—178, 225, 241
	<i>Ægopodium</i>	111	<i>Alfinella</i>	63, 64
	<i>Æthusa</i>	107	<i>Alsinoides</i>	10
	<i>Agaricus</i>	490—497	<i>Althæa</i>	267
	<i>Agaricum</i>	463	<i>Amanita</i>	492
	<i>Agrifolium</i>	505	<i>Amara dulcis</i>	79
	<i>Agrimonia</i>	180	<i>Amaranthus</i>	356
	<i>Agrostemma</i>	173	<i>Ambrosia</i>	248
			<i>Ana-</i>	

INDEX NOMINUM LATINORUM.

	Page		Page
<i>Anacampteros</i>	170, 370	<i>Asphodelus</i>	125, 128
<i>Anagallis</i>	73, 74, 51, 79	<i>Asplenium</i>	384—386
<i>aquatica</i>	4, 5	<i>Aster</i>	319, 314, 320
<i>Anblatum</i>	233	<i>Astragalus</i>	281, 282, 274
<i>Anchusa</i>	66	<i>Athamanta</i>	100
<i>Andromeda</i>	154	<i>Atriplex</i>	376—378, 90
<i>Androsæmum</i>	291		—92
<i>Anemone</i>	208, 209	<i>Atropa</i>	79
<i>Anethum</i>	110	<i>Avena</i>	41, 42
<i>Angelica</i>	103, 111	<i>Auricula muris</i>	168, 175
<i>Anonis</i>	273	B	
<i>Anthemis</i>	323, 324	<i>Baccharis</i>	314
<i>Anthericum</i>	127, 128	<i>Balladonna</i>	79
<i>Anthoceros</i>	441	<i>Ballota</i>	227
<i>Anthoxanthum</i>	10, 135	<i>Balsamine</i>	332
<i>Anthyllis</i>	273, 120, 167	<i>Barba capræ</i>	191
<i>Antirrhinum</i>	236—239	<i>Barbarea</i>	251
<i>Aparine</i>	57—69	<i>Bardana</i>	304, 356
<i>Aphaca</i>	275	<i>Bartsia</i>	233
<i>Aphanes</i>	60	<i>Becabunga</i>	4
<i>Apium</i>	111, 100, 102	<i>Bellis</i>	320, 321
<i>Aponogeton</i>	344	<i>Berberis</i>	119
<i>Aquilegia</i>	207, 208	<i>Beta</i>	93
<i>Arabis</i>	254	<i>Betonica</i>	220, 240
<i>Aracus</i>	279, 280	<i>Betula</i>	354
<i>Arbutus</i>	155	<i>Betulus</i>	360
<i>Arctium</i>	304	<i>Behen album</i>	163
<i>Arenaria</i>	167—169	<i>Bidens</i>	309, 310
<i>Argemone</i>	202, 204	<i>Bifolium</i>	338, 339
<i>Argentina</i>	195	<i>Biforta</i>	146, 147
<i>Aria</i>	187	<i>Blackstonia</i>	146
<i>Armeria</i>	161, 162, 166	<i>Blafia</i>	439
<i>Armerius</i>	174	<i>Blattaria</i>	77
<i>Artemisia</i>	311, 312	<i>Blitum</i>	89—93, 356
<i>Arum</i>	342	<i>Boletus</i>	496—498
<i>Arundo</i>	43, 21	<i>Bonus Henricus</i>	89
<i>Asarum</i>	179	<i>Borrage</i>	68, 66, 69
<i>Ascyron</i>	291, 292	<i>Batrytis</i>	504
<i>Asparagus</i>	126	<i>Brassica</i>	252—255, 261
<i>Aspergilla</i>	504	<i>Briza</i>	32
<i>Asperugo</i>	69	<i>Bromus</i>	39—41
<i>Asperula</i>	55	Bry-	

INDEX NOMINUM LATINORUM.

	Page		Page
Bryonia	373, 369	Carum	110
Bryum	403—417, 397	<i>Caryophyllata</i>	198, 199
<i>Bucca-ferrea</i>	63	<i>Caryophyllus</i>	64, 114, 161,
Bufonia	10		162, 166, 174, 176
<i>Buglossum</i>	66, 68, 69, 294	<i>Cassida</i>	231, 232
Bugula	219, 220	<i>Catanance</i>	275
<i>Bulbocastanum</i>	105	Castanea	359
<i>Bulbocodium</i>	122	Caucalis	98, 99
<i>Bulbosa alpina</i>	122	<i>Cauda muris</i>	118
Bunium	105	<i>equina</i>	1
<i>Bupthalmum</i>	324	Centaurea	325, 326
Bupleurum	97	<i>Centaurium</i>	88, 89, 146
<i>Bursa pastoris</i>	243, 247,	Centunculus	51
	249	Cerastium	175—177
Butomus	152	Cerasus	186, 187
Buxus	354, 355	Ceratophyllum	357
Byffus	487—489	<i>Cerefolium</i>	108
		<i>Cerinthæ</i>	67, 68
		<i>Ceterach</i>	385
C		<i>Chærefolium</i>	109
<i>Akile</i>	261	<i>Chærophyllo similis</i>	60
<i>Calamagrostis</i>	25, 43	<i>Chærophyllo</i>	108, 109
<i>Calamaria</i>	393	<i>Chamæcissus</i>	205, 206
<i>Calamintha</i>	224, 230	<i>Chamædryas</i>	218, 5, 199
<i>Calamus aromaticus</i>	128	<i>Chamæfilix</i>	386
<i>Calceolus</i>	341	<i>Chamælea</i>	145
Callitriche	2	<i>Chamælinum</i>	116, 117
Caltha	214, 215	<i>Chamæmelum</i>	322—324
<i>Camelina</i>	243, 251	<i>Chamænerion</i>	140, 142
<i>Cameraria</i>	48	<i>Camæorchis</i>	339
Campanula	80—82	<i>Chamæperichlymenum</i>	58
<i>Canabina</i>	309	<i>Chamæpitius</i>	219
<i>Cannabis</i>	226	<i>Chamæpitea</i>	365
<i>Capillus veneris</i>	391	Chara	464, 465
<i>Caprifolium</i>	80	Cheiranthus	250
Cardamine	255—258, 245	Chenopodium	89—93
<i>Cardiaca</i>	228	Chelidonium	201, 202, 214
Carduus	305—308, 49,	<i>Chondrilla</i>	296
	326	<i>Christophoriana</i>	201
Carex	346—354	Chrysanthemum	321
Carlina	309	Chrysofenium	156, 241
Carpinus	360		Cicho-
<i>Caruifolia</i>	106		

INDEX NOMINUM LATINORUM.

	Page		Page
<i>Cichorium</i>	303, 294, 301	<i>Cuscuta</i>	89
<i>Cicuta</i>	106, 100, 107	<i>Cyanus</i>	325
<i>Cicutaria</i>	107, 108	<i>Cymbalaria</i>	237
<i>Circæa</i>	9, 10	<i>Cynocrambe</i>	371
<i>Cirsium</i>	305—307	<i>Cynoglossum</i>	67, 68
<i>Cistus</i>	204—206, 155	<i>Cynosorchis</i>	333—335
<i>Clavaria</i>	500, 501	<i>Cynosurus</i>	47
<i>Clematis</i>	291	<i>Cyperus</i>	15, 14, 19
<i>Clynopodium</i>	228, 230, 233	<i>Cypripedium</i>	341
<i>Clymenum</i>	291	D	
<i>Cnicus</i>	309	<i>Dactylis</i>	25
<i>Cochlearia</i>	247, 248	<i>Damafonium</i>	137, 138
<i>Colchicum</i>	136	<i>Daphne</i>	145
<i>Conferva</i>	478—486	<i>Datura</i>	78
<i>Conium</i>	100	<i>Daucus</i>	99, 100
<i>Consolida</i>	207, 219, 220	<i>Delphinium</i>	207
<i>Convallaria</i>	126, 127	<i>Dens leonis</i>	297, 301
<i>Convolvulus</i>	74, 75, 149	<i>Dentaria</i>	249
<i>Conyza</i>	314, 320, 327	<i>Dianthus</i>	161, 162
<i>Corallina</i>	479	<i>Diapensia</i>	96
<i>Coralloides</i>	443, 449, 455—460, 503	<i>Digitalis</i>	240
<i>Corallo-fungus</i>	501	<i>Dipsacus</i>	49
<i>Coriandrum</i>	107	<i>Dörtmanna</i>	329
<i>Cornu Cervi</i>	53	<i>Draba</i>	243, 244
<i>Coronopus</i>	52, 53, 248	<i>Dracunculus</i>	325
<i>Corrigiola</i>	86	<i>Drosera</i>	117
<i>Corylus</i>	360	<i>Dryas</i>	199
<i>Cotyledon</i>	169, 96, 157	<i>Dryopteris</i>	384, 388, 391
<i>Cracca</i>	277, 280	E	
<i>Crambe</i>	260	<i>EBulus</i>	113
<i>Crassula</i>	170	<i>Echium</i>	69, 70, 65, 68
<i>Cratægus</i>	187, 188	<i>Echinophora</i>	97, 99
<i>Cratægonon</i>	235, 236	<i>Elatine</i>	151, 237
<i>Crepis</i>	301, 297	<i>Elleborine</i>	341
<i>Crithmum</i>	97, 101, 320	<i>Eleoselinum</i>	111
<i>Cristagalli</i>	234	<i>Elichrysium</i>	313
<i>Crocus</i>	13	<i>Elymus</i>	44, 45
<i>Cruciata</i>	89, 375	<i>Embolus</i>	504
<i>Cucubalus</i>	163, 164, 174	<i>Elvela</i>	498
<i>Cuminum</i>	110	<i>Empetrum</i>	367
		<i>Epi-</i>	

INDEX NOMINUM LATINORUM.

	Page		Page
<i>Epilobium</i>	140—142	<i>Frangula</i>	83
<i>Equisetum</i>	380, 381, 1, 151	<i>Frankenia</i>	119, 120
<i>Erica</i>	144, 145, 119, 143, 155, 367, 469	<i>Fraxinus</i>	379, 189
<i>Ericoides</i>	151	<i>Fritillaria</i>	125
<i>Erigeron</i>	314, 316	<i>Fucus</i>	465—477, 342, 343, 465, 480, 486, 490
<i>Erinaceus</i>	497	<i>Fucoides</i>	473
<i>Eriophorum</i>	19, 20	<i>Fumaria</i>	270
<i>Eruca</i>	243, 250, 251, 253, 258, 259, 261	<i>Fungoides</i>	498—501
<i>Ervum</i>	279, 280, 275	<i>Fungus</i>	463, 479, 490—504
<i>Eryngium</i>	95	G	
<i>Erysimum</i>	250, 251, 150, 259	G <i>ALE</i>	368
<i>Erysimo similis</i>	254	<i>Galeopsis</i>	225—227
<i>Esula</i>	182	<i>Galium</i>	55—58, 375
<i>Euonymus</i>	84	<i>Geaster</i>	502
<i>Eupatorium</i>	310, 180, 309	<i>Genista</i>	271, 272
<i>Euphorbia</i>	181—185	<i>Gentiana</i>	87, 88, 146
<i>Euphrasia</i>	234, 235, 233	<i>Gentianella</i>	88
F		<i>Geranium</i>	262—267
F <i>Agus</i>	359, 360	<i>Geum</i>	198, 199, 157, 158
<i>Fegopyrum</i>	149, 150	<i>Gladiolus</i>	13, 152, 329
<i>Festuca</i>	36—40	<i>Glastrum</i>	261
<i>Ferrum equinum</i>	281	<i>Glaux</i>	86, 281, 282
<i>Ficaria</i>	213, 214	<i>Glecoma</i>	224
<i>Filago</i>	328, 329, 313	<i>Glycyrrhiza</i>	281
<i>Filipendula</i>	105, 190	<i>Gnaphalium</i>	312, 313, 328, 329
<i>Filix</i>	382—392	<i>Gramen</i>	14—48, 114, 130—133, 166, 346, 354, 371, 374
<i>Filicula</i>	384	<i>Graminifolia</i>	242, 393
<i>Filum marinum</i>	475	<i>Grossularia</i>	84, 85
<i>Fœniculum</i>	100	<i>Gratiola</i>	180
<i>Fontalis</i>	147	<i>Gypsophylla</i>	232
<i>Fontinalis</i>	60, 398	H	
<i>Flos adonis</i>	209	H <i>Alimus</i>	376
<i>cuculi</i>	256	<i>Hedera</i>	85
<i>Fœnumgræcum</i>	283	<i>terrestris</i>	224
<i>Fœnum burgundiacum</i>	288	<i>Hederula</i>	344
<i>Fragaria</i>	194, 195	<i>He-</i>	
<i>Franca</i>	119		

INDEX NOMINUM LATINORUM.

	Page				Page
Hedysarum	281	J			
Helenium	319				
Helianthemum	205, 206	JACEA			325, 226
Helleborine	341, 342	<i>Jacobæa</i>		316, 317,	327
Helleboraster	215	Jasione			329
Helleborus	215, 214	Iberis			249
Helxine	149	Ilex			505
Hepatica	439	Illecebrum		85,	171
Heracleum	102	Impatiens			332
Herba dysenterica	320	Inula		319,	320
Gerardi	111	Iris			259
Paris	150	Iris			13
trientalis	139	Ifatis			261
Herniaria	94, 11	Ischæmon			22
Hesperis	251	Isoetes			393
Hieracium	297—303,	<i>Juncellus</i>		14, 16—	18
	295	<i>Juncifolia</i>			242
Hippophaë	368	Juncus		129—132,	14, 16,
Hipposelinum	109			18,	20
Hippocrepis	281	<i>Juncus Floridus</i>			152
Hippuris	1	Jungermannia		430—	438
Holcus	374	Juniperus			372
Holosteum	118, 176				
Hordeum	46	K			
Horminum	9	KALI		92, 93	
Hottonia	72	<i>geniculatum</i>			1
Humulus	369	Keiri			250
Hyacinthus	123, 124	Knawel			160
Hydnum	497				
Hydroceratophyllum	357	L			
Hydrocharis	372	L Actuca		295, 296,	13
Hydrocotyle	96	<i>Lagopodium</i>			285
Hydrolapathum	135	<i>Lagopus</i>			285
Hyoscyamus	77	Lamium		225—227,	231
Hyoseris	302	Lampfana			303
Hypericum	290—292	Lantana			112
Hypnum	418—430	Lapathum		89, 133—	136
Hypochæris	302, 303	Lappa		304,	356
Hypopitys	153	Lapsana			303
Hyssopifolia	180	Lathræa			233
		La			

INDEX NOMINUM LATINORUM.

	Page		Page
Lathyrus	274—277	Lychnoides	173
Lavatera	267	Lycoperdon	501—503
Laureola	145	Lycopodium	393—395
Ledum	155	Lycopsis	69, 70
Lemna	344, 345	Lycopus	8
Lens palustris	345	Lythamchia	72, 73, 140—
Lentibularia	8		142, 179, 180, 232
Lenticula	344, 345	Lythrum	179, 180
Leontodon	296, 297	Lythrophyton	500
Leonurus	228, 226		
Lepidium	244, 245		
Leucoium	250, 330		
Leucanthemum	321		
Lichen	437—463, 477		
Lichenastrum	402, 430—		
	438		
Lichenoides	441—460, 464		
Ligusticum	102		
Ligustrum	3		
Ligula marina	475		
Lilium convallium	126		
Limnapeuce	1		
Limodorum	336		
Limonium	114, 115		
Limosella	241		
Linagrostis	19		
Linaria	86, 237, 238		
Lingua cervina	385		
Linum	115—117		
Lithospermum	65, 66		
Lobelia	329		
Lolium	44		
Lonchitis	383, 388		
Lonicera	80		
Lotus	288, 289		
Loto affinis	274		
Lunaria	244, 382, 439		
Lupulus	369		
Luteola	181		
Lychnis	174, 119, 160,		
	163—166, 173		

INDEX NOMINUM LATINORUM.

	Page		Page
<i>Morsus ranæ</i>	372	<i>Ophioglossum</i>	382
<i>Moscatellina</i>	150	<i>Ophrys</i>	338—340
<i>Mucor</i>	503	<i>Opulus</i>	112
<i>Mucipula</i>	163, 174	<i>Orchis</i>	333—340
<i>Muscus</i>	394—432, 449, 450, 459, 477, 480, 483, 486	<i>Origanum</i>	229
<i>Musco-fungus</i>	448, 451— 463	<i>Ornithopus</i>	280
<i>Myagrum</i>	241, 251	<i>Ornithopodium</i>	280
<i>Myosotis</i>	65	<i>Ornithogalum</i>	124, 125
<i>Myosurus</i>	118	<i>Ornus</i>	189
<i>Myriophyllum</i>	357, 358	<i>Orobanche</i>	232, 233, 153
<i>Myrrhis</i>	108	<i>Orobis</i>	274, 275
<i>Myrtus</i>	368	<i>Osmunda</i>	282, 283
		<i>Ostrya</i>	360
		<i>Othonna</i>	327
		<i>Oxalis</i>	173, 136
		<i>Oxyacantha</i>	188
		<i>Oxycoccus</i>	143
		<i>Oxys</i>	173
N			
NAPUS	253		
<i>Narcissus</i>	122, 123		
<i>Nardus</i>	20, 28, 374		
<i>Narthesium</i>	127, 128		
<i>Nasturtium</i>	243—245, 248, 249, 255—257, 259		
<i>Nepeta</i>	220		
<i>Nidus avis</i>	336, 338		
<i>Noli me tangere</i>	256		
<i>Nostoc paracelsi</i>	463		
<i>Nucula terrestris</i>	106		
<i>Nummularia</i>	73		
<i>Nux vesicaria</i>	113		
<i>Nymphæa</i>	206, 270, 71, 373		
O			
OCYMUM	230		
<i>Oenanthe</i>	104, 105		
<i>Onobrychis</i>	281		
<i>Ononis</i>	273		
<i>Onopordon</i>	308		
<i>Onopteris</i>	387		
P			
PADUS	187		
<i>Palma Christi</i>	335, 336		
<i>Panicum</i>	21, 22, 26, 47		
<i>Panax coloni</i>	227		
<i>Papaver</i>	201—204, 163		
<i>Paralysis</i>	71		
<i>Parietaria</i>	376		
<i>Paris</i>	150		
<i>Parnassia</i>	114		
<i>Paronichia</i>	159, 243—245		
<i>Pecten veneris</i>	108		
<i>Pedicularis</i>	236, 234		
<i>Peloria</i>	238		
<i>Pentaphylloides</i>	118, 195, 196, 200		
<i>Pentaphyllum</i>	59, 195— 198		
<i>Pentapterophyllum</i>	358		
<i>Peplis</i>	128, 181		
<i>Peplus</i>	182		
<i>Percepier</i>	60		
			<i>Per-</i>

INDEX NOMINUM LATINORUM.

	Page		Page
<i>Perfoliata</i>	97, 252	<i>Polymonium</i>	165
<i>Perichlymenum</i>	58	<i>Polytrichum</i>	399—401, 416
<i>Perficaria</i>	147, 148, 332	<i>Populago</i>	215
<i>Pes anserinus</i>	90	<i>Populus</i>	370
<i>Petasites</i>	315	<i>Portulaca</i>	48, 128, 376
<i>Petrofelinum</i>	109	<i>Potamogeton</i>	60—63, 147, 357, 358
<i>Peucedanum</i>	101	<i>Potentilla</i>	195—197
<i>Peziza</i>	498—500, 463	<i>Poterium</i>	358
<i>Phalangium</i>	127, 128	<i>Prenanthes</i>	296
<i>Phalaris</i>	20, 21, 32	<i>Primula</i>	70, 71
<i>Phallus</i>	498	<i>Prunella</i>	231
<i>Phascum</i>	396, 397	<i>Prunus</i>	186, 187
<i>Phellandrium</i>	107	<i>Pseudo-asphodelus</i>	127
<i>Phleum</i>	22, 23, 21	<i>Cyperus</i>	352
<i>Phyllitis</i>	385	<i>Helleborus</i>	214
<i>Phyteuma</i>	83	<i>melanthium</i>	173
<i>Picea</i>	361	<i>narcissus</i>	122
<i>Picris</i>	294	<i>orchis</i>	337
<i>Pilularia</i>	393	<i>Ptarmica</i>	325
<i>Pilosella</i>	255, 290, 299	<i>Pteris</i>	384
<i>Pimpinella</i>	110, 111, 54, 358, 359	<i>Pulegium</i>	224
<i>Pinguicula</i>	7	<i>Pulmonaria</i>	67, 299, 300
<i>Pinus</i>	361, 470	<i>Pulsatilla</i>	209
<i>Piperitis</i>	244	<i>Pyrola</i>	153, 154, 139
<i>Piftacea</i>	113	<i>Pyrus</i>	189
<i>Pifum</i>	274	<i>Pyrafter</i>	189
<i>Plantaginella</i>	241		
<i>Plantago</i>	51—53, 137, 138		
<i>Pneumonanthe</i>	87		
<i>Poa</i>	32—35, 25, 29, 36		
<i>Podagraria</i>	111		
<i>Polemonium</i>	75		
<i>Polygala</i>	271, 86		
<i>Polygon</i>	281		
<i>Polygonatum</i>	126, 127		
<i>Polygonum</i>	146—150, 60, 86, 94, 117, 119, 120, 160		

Q *Quercus* 358, 466
Quinquefolium 196, 197, 200

R *Radiola* 117
Radix cava 150
Ranunculus 208—214, 150, 344
Rapa 253
L l 2 Ra-

INDEX NOMINUM LATINORUM.

	Page		Page
<i>Rapum</i>	253	Sanguisorba	53, 54, 358, 359
<i>geniflæ</i>	232	Sanicula	96, 7, 157
Raphanus	252, 244, 248, 258, 261	<i>Santolina</i>	328
<i>Raphanistrum</i>	252	Saponaria	160, 161
Rapistrum	252, 258, 260	Satyrion	336—338
<i>Rapunculus</i>	81, 83, 96, 329	Saxifraga	156—159, 55, 63, 64, 102, 106, 178
<i>Rapuntium</i>	81	Scabiosa	50, 51, 326, 329
<i>Regina prati</i>	199	Scandix	107, 108
Refeda	181	Schoenus	14, 15
Rhamnoides	368	Scilla	123, 124
<i>Rhamnus</i>	83, 368	Scirpus	16—19
Rhinanthus	234	Sclarea	9
<i>Rhodia</i>	371	Scleranthus	160
Rhodiola	370	<i>Scordium</i>	120
<i>Rhus</i>	368	<i>Scorodonia</i>	219
Ribes	84, 85	<i>Scrodostrasum</i>	120
Riccia	440, 441	Scrophularia	239, 240
<i>Rorella</i>	117	Scutellaria	231, 232
<i>Ros folis</i>	117	Secale	46, 44
Rosa	191, 192	Sedum	170—173, 93, 156, 160, 185
<i>Rosmarinum</i>	155	<i>Selaginoides</i>	395
<i>Rubeola</i>	54, 55	<i>Selago</i>	395
Rubus	193, 194	<i>Selinum</i>	100, 104
Rumex	133—136	Sempervivum	185, 171, 172
Ruppia	63	Senecio	315—318, 301
Ruscus	373	Serapias	341, 336, 337
<i>Ruta muraria</i>	386	<i>Serpyllum</i>	229, 230
S		Serratula	304, 305
<i>Sagina</i>	63, 169	<i>Sesamoides</i>	163
<i>Sagitta</i>	358	<i>Sesamum</i>	243
<i>Salcaria</i>	17 180	Seseli	106
Salicornia	1	Sherardia	54
Salix	362—366	Sibbaldia	118
Salsola	93	Sibthorpia	241
Salvia	9	Sideritis	226, 227, 220, 224
<i>S. mbucus</i>	112, 113	Silene	
<i>S. molus</i>	79		

INDEX NOMINUM LATINORUM.

	Page
Silene	164, 165, 174
<i>Sinapi</i>	260
Sinapis	260
Sison	104
Sisymbrium	257 — 259, 223
Sium	103, 104, 106, 256
<i>Smilax</i>	74
Smyrnium	109, 102
Solanum	78, 79, 150
<i>Solanifolia</i>	9, 10
Solidago	317—319, 327
<i>Soldanella</i>	75
Sonchus	294—296
<i>Sophia</i>	259
Sorbus	188, 189
Sparganium	346
<i>Spartium</i>	271
<i>Spartum</i>	20, 24, 25, 44
Spergula	177, 178, 169
Sphagnum	395—399, 403
<i>Spherocephalus</i>	503
<i>Sphondylium</i>	102
<i>Spina alba</i>	308
<i>acida</i>	119
<i>cervina</i>	83
Splachnum	399
Spongia	489—490, 471
Stachys	226, 227
<i>Staphylodendron</i>	113
Statyce	114, 115
Staphylæa	113
Stipa	24
Stramonium	78
Stratiotes	208, 372
Subularia	242, 393
<i>Suillus</i>	497
Swertia	87
Symphytum	68, 67
<i>Synanchica</i>	55

T

	Page
TAMUS	369
<i>Tamnus</i>	369
Tanacetum	310
Targionia	438
Taxus	372
<i>Telephium</i>	170
Teucrium	218, 219, 199
Thalictrum	216
<i>Thapsus</i>	75
Thesium	86
Thlaspi	245—248, 243
Thymus	229, 230
Tilia	204
<i>Tithymalus</i>	181—185
Tordilyum	98, 99
Tormentilla	197, 198, 59
<i>Trachelium</i>	81, 82
Tragopogon	293
<i>Tragorchis</i>	334, 337
Tremella	463, 464, 476, 477, 483
<i>Tribulus</i>	61
Trichomanes	385
Trifolium	282—288, 71, 173
Triglochin	183
<i>Triorchis</i>	338
<i>Tripolium</i>	319
Triticum	45
Trollius	214
<i>Tuber</i>	501, 502
<i>Tunica</i>	162
Turritis	254, 255
Tuffilago	315
Typha	345, 346

V

Stellaria 166

INDEX NOMINUM LATINORUM.

		Page
V		
V Accinium	142, 143, 155	
Valantia	375	
Valeriana	12, 13, 75	
Valerianella	12, 13	
Vella	243	
Verbascum	75, 76	
Verbasculum	70, 71	
Verbena	505	
Verbesina	309, 310	
Vermicularis	93, 170, 171	
Veronica	3—7, 237	
Viburnum	112	
Vicia	277—280, 275	
Vinca	77	
Viola	330—332, 7, 72, 250	
Virga aurea	317—319	
pastoris	5	
Viscaria	161	
Viscago	164	
Viscum		367
Vitis idæa	142, 143, 155	
Volubilis	145	
Vulneraria	273	
U		
U LEX		272
Ulmaria		191
Ulmus		94, 95
Ulva	475, 437, 440, 463	
Umbilicus veneris		169
Urtica		355, 226
Usnea		461—463
Utricularia		8
X		
X Anthium		356
Xyris		13
Z		
Z Anichellia		344
Zostera		342



I N D E X

NOMINUM ANGLICANORUM.

A		B	
	Page		Page
A dders tongue	382	B ALM, bastard	231
Agaric	491—496	Bane-berries	201
Agrimony	180	Barberry	119
Dutch	310	Barley	46
Hemp	309, 310	Basil	228
Alder, common	505	Beam-tree, white	188
berry bearing	84	Bears-foot	215
Alexanders	109	Bedstraw	56, 57
Alkanet, bastard	66	Beech-tree	360
All-heal	226, 227	Beet	93
feed	117	Bell-flower	80—82
Aloe water	208	Betony, Paul's	4
Angelica	103	water	240
Anthoceros	441	wood	220
Anemone	208, 209	Bill-berries	142, 143
Apple-tree	190	Bind-weed	74, 75
thorny	78	black	149
Archangel	225	Birch-tree	354
Arrow-head	358	Birds-cherry	187
Arsmart	147, 148	eye	71
Afarabacca	179	foot	280
Ash-tree	379	nest	99, 152, 336
mountain	189	tongue	318
Ashweed	111	Biftort	147
Afp	370	Bitter-sweet	79
Asphodel	127, 128	Blafia	440
Avens	198, 199	Blite	90—92, 356
Awl-wort	242	Bloodwort	133
		Blue-	

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Blue-bottle	325	Caraways	100
Boletus	496, 497	Cat-mint	220
Borage	68	Cats-tail	345, 346
Box-tree	355	Catchfly	163—166, 174
Bramble	193, 194	Celandine, greater	202
Brank	150	lesser	214
Briar, common	193	Centaury	88, 89, 146
sweet	191	Champignon	491
Brooklime	4	Charlock	252, 260
Broom	271	Chara	465
rape	232	Cheese-rening	56
Briony, black	369	Cherry-tree	187
white	372	Chervil, wild	108
Bryum	403—417	Chestnut-tree	359
Buck-bean	71	Chickweed, bastard	11
wheat	150	berry-bearing	163
Bugle	219, 220	breakstone	64
Bugloss	69, 70	common	113, 167
Cowslips	67	—	169
Bullace-tree	186	germander	6
Burdock	304, 345	mouse-ear	176, 177
lesser	356	star-headed	2
Burnet	54, 359	water	48
Bur reed	347	winter-green	139
Butchers-broom	373	Cicely, wild	108
Butter-cups	211	Cinquefoil	195—197
wort	7	bastard	118
Byssus	487—489	marsh	200
		Cistus, dwarf	205, 206
		marsh	155
		Clary	9
C		Clavaria	500, 501
Cabbage	253	Claver	288
Calamint	230	Climber	217
Calamus	128	Clot-bur	304
Caltrops	61	Cloud-berry	194
Calves-snout	239	Clover	283, 284
Cammock	273	Cockle	173
Camomile	222, 224	Cock's-comb	234
Campion	163, 164, 174	head	281
Canterbury-bells	82	Codlings and Cream	141
Carex	346—354	Colewort	
Carrot	99		

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Colewort	262	Devil's guts	89
Coltsfoot	315	Dew-berry	193
Columbines	207, 208	Dier's-weed	181, 243
Comfrey	68	Dock	134, 135
Conferva	478—486	Dodder	89
Coral-wort	249	Dropwort	190
Coriander	107	water	105
Cows lungwort	46	Ducks meat	345
weed	108	Dwale	79
wheat	233—236		
Cowslips	71		
Crab-tree	190	E	
Cranberries	143	ELDER	112, 113
Cranesbill	262—267	water	113
Crefs, bastard	245	Elecampane	319
bitter	256	Elm	95
penny	246	Elvela	498
rocket	243	Eryngo	95
rock	249	Eyebright	234, 235
swines	248	cow-wheat	233, 234
water	257		
winter	251	F	
Crosswort	58, 374	FEllwort	88
Crow-foot	211—213	Fennel	110
berries	367	hogs	101
Cudweed	212—214, 328,	Fern	382—384
	329	Feverfew	322
Currants	84, 85	Figwort	239, 240
Cyperus, bastard	14, 19	Finckle	110
millet	19	Flag, sweet smelling	128
sweet	15	Flax	115—117
		Fleabane	314, 320
		Flixweed	259
		Flos adonis	209
D		Flower de luce	13
DAffodil	123	Fluellin	3, 237
chequered	125	Fool-stones	333, 334
Daisy	321	Fox-glove	240
Dandelion	297	Frambose	193
Danewort	113	Fritillary	125
Deptford pink	161	Frog-bit	372
Devil's bit	50, 298		
		M m	
			Fucus

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Fucus	466—475	Grafs, fescue	36—38
Fumitory	270	five-leaved	197
Furze	272	fox tail	23, 24
		hard	375
		hair	28—31
		knot	160
		lyme	45
		meadow	33—35
		melic	32
		mat	20
		millet	26
		oat	41—43
		panic	21, 22
		parnassus	114
		poly	180
		quaking	32
		quick	45
		rye	46
		reed	43, 44
		soft	374
		spring	10
		sea	63
		scurvy	75
		spiked sea	133
		vernal	10
		wheat	45
		Greymill	66
		Gromill	66
		Gromwell	66
		Ground-ivy	224
		pine	219
		Groundfel	316
			H
		Hard-beam	360
		Hare-bells	123
		strong	101
		Hares-car	97
		Harts-tongue	385
		Hafel-nut-tree	360
		wych	95
		Hawk.	
G			
G Alingale	15		
G Garlick	120, 121		
G Gentian	87, 88		
G Germander	5, 218		
G Gillyflower, stock	250		
sea	114		
G Gladdon, stinking	13		
G Gladiole	152, 330		
G Gladwyn	13		
G Glasswort	1, 92, 93		
G Globe-flower	214		
G Goats-beard	293		
stones	337		
G Goldilocks	211		
G Gold of Pleasure	242		
G Golden rod	319		
G Goose-corn	130		
foot	90		
grafs	56—59		
tongue	325		
G Gorfe	272		
G Goule	368		
G Gout-weed	111		
G Grafs, arrow-headed	133		
barley	46		
bent	26—28		
brome	39—41		
canary	20, 21		
cats-tail	23		
cocks-foot	25		
cotton	19		
couch	45		
darnel	44		
dogs	45		
tail	47, 48		
feather	24		

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Hawk-weed	294, 298—	J	
Hawthorn	303	Jack-by-the-hedge	251
Hearts-ease	188	Jacob's ladder	75
Heath	331	Jews-ear	464
berried	144, 145	Jungermannia	430—438
sea	367	Juniper	372
Hedge hyssop	120	Ivy	85
Hellebore, bastard	180		
	215,	K	
	341, 342	Kidney-wort	157, 169
Hellweed	84	vetch	275
Hemlock	100	Knapweed	326
water	106, 107	Knawel	160
Henbane	78	Knot-berries	194
Henbit	225	grafs	149
Hep-tree	192	German	160
Herb Christopher	201	verticillate	86
Gerard	111	Knout-berries	194
Paris	150		
Robert	269	L	
High-taper	76	Ladies bedftraw	56, 57
Hindberry	193	finger	274
Holly-tree	505	hair	32
knee	373	mantle	59
Honeyfuckle	80	flipper	341
Hops	369	smock	255—257
Horehound, bafe	226	traces	338
black	228	Lambs lettuce	13
ftinking	228	Larkspur	207
water	9	Laver	476—478
white	228	Laurel, fpurge	145
Horn-beam	360	Lavender, fea	115
beech	360	Lettuce	296
Horfe-beech	360	frogs	61
radifh	249	Lilly, of the valley	126
tail	380, 381	water	71, 206, 207
Houfeleck	185	Lime-tree	204
Hounds-tongue	67	wort	161
Hyacinth	123, 124	Linden-tree	204
Hydnum	497	Livelong	170
Hypnum	428—430	M in 2	Liver-

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Liverwort	442, 463	Missel, or Misseltoe	367
Lizard flower	337	Mnium	401—403
Locker-gowls	214	Money-wort	73
Looking-lafs, Venus's	82	bastard	272
Loofestriſe, ſpiked	180	baud	100
yellow	72, 73	Moonwort	382
Louſewort	236	Moorberries	143
Lucern	288	Morel	498
Lungwort, cows	76	Mofchatel	151
French	300	Mofs-berries	143
golden	300	bog	396
tree	449	cup	456
		club	394, 395
		water	358, 359
M		Mother of thyme	230
Madder, baſtard	56	wort	228
croſſwort	58	Mucor	503, 504
field	54	Mouſe-ear, coddled	255
wild	54, 56	creeping	298
Madwort	69	ſcorpion-graſs	65
Maidenhair	285—287	tail	118
golden	400	Mug-wort	312
true	391, 392	weed	375
Maithes	209	Mullein	76
Mallow	267—269	Muſhroom	494
Maple	379	Muſcovy	262
Marjoram, wild	229	Muſtard, common	260
Mares-tail	2	hedge	250—259
Marigold, corn	321	mithridate	245
marſh	215	tower	254
Maſſellon	326	treacle	246
Mayweed	323	white	260
Meadowſweet	191	wild	260
Medick	289	Myrtle	368
Medlar	190		
Melilot	282		
Mercury	89, 371		
Milfoil	325	N	
water	357, 358	NAVEW	253
Milkwort	271	Navelwort	169
ſea	86	Needle, ſhepherd's	108
purple	282	Nep mint	225
		Nettle, common	355
		Nettle	

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Nettle, hemp	226	Peafe-heath	274
hedge	227	fea	274
roman	355	Pellitory, wall	378
Nightshade	78, 79	Penny-royal	224
enchanters	9, 10	wort, marsh	66
Nipplewort	303	wall	169
Nut, earth	106	Pepper-grafs	392
bladder	113	wall	178
		water	148
		wort	244
O		Periwinkle	77
OAK	359	Peziza	499, 500
fea	466	Phascum	397
Orach	376—378	Pheasant's eye	209
wild	91	Pilcorn	41
Orpine	170	Pilewort	214
Orchis	333—336	Pimpernel, bastard	51
Ofier	366	male	74
Osmund royal	382	water	79
Ox-eye	321	yellow	73
tongue	294	Pink	161, 162
Oyster-green	477	Pine, fea	470
		Pipperidge-bush	119
		Plantain	51—53
P		bastard	241
PAGIL	71	water	137, 138
Pansies	331	Plowman's spikenard	314
Park-leaves	291	Polypody	387—389
Parsley, bastard	98, 99	Polytrichum	400, 401
corn	98, 104	Pondweed	60—63
fools	107	horned	357
hedge	99	Poplar	370
knotted	98	Poppy	202—204
piert	60	Prick-madam	170
fea	102	wood	84
stone	100, 104	Prim	3
Parsnep, cows	102	Primrose	70
fea	97	peerless	123
water	103, 104	Privet	3
wild	109	Puff-balls	502, 503
Pasque flower	209	Purging-thorn	85
Pearwort	64	Pur-	
Pear-tree	189		

INDEX NOMINUM ANGLICANORUM.

Purslane, water	Page 128		
fea	379	S	Page
		SAGE of Jerusale	67
		wood	219
Q		Saffron	13, 122, 136
Quick in hand	332	Saintfoin	281
Quicken-tree	189	St. John's wort	290, 291
Quill-wort	393	St. Peter's wort	292
		Sallow	366
R		Saltwort	I
R Adish, horse	249	black.	86
water	258	Sampire	I, 97, 101, 320
Ragwort	317, 318, 327	Sanicle	96
Rampions	81, 83	Yorkshire	7
Ramsons	122	Satyrion	337
Rasperry	193	Sauce alone	251
Rattle, yellow	234	Saw-wort	304, 305
Reed	43, 345	Saxifrage, burnet	110, 111
Rest-harrow	273	Cornwall	102
Ribwort	52	English	178
Riccia	440, 441	golden	156
Rocket, base	181	meadow	106
wall	258	white	159
fea	261	Scabious	50, 51
water	258	sheeps	329
wild	254	Scurvygrafs	75, 247, 248
winter	251	Self-heal	231
Ros folis	191, 192	Sea-beard	485
Rosewort	371	Sengreen	157
Rosemary, wild	155	Septfoil	198
Rue, meadow	216	Service	188
Rupture-wort	94	Setterwort	215
least	117	Shepherd's-purse	247
Rush	129, 132	Sicklewort	219
bog	14	Silverweed	195
bull	17	Sloe-tree	186
club	16—18	Smellage	111
flowering	152	Snakeweed	147
grafs	15	Snapdragon	239
hare's-tail	20	Sneczewort	325
		Soapwort	

INDEX NOMINUM ANGLICANORUM.

	Page
Soapwort	161
Soldier, fresh-water	208
Solomon's seal	126, 127
Sorb	188, 189
Sorrel	136
wood	173
Southernwood	311
Sow-thistle	295
bane	90
Sparagus	126
Spearwort	210
Speedwell	3—7
Sperage	126
Spiguel	100
Spikenard	314
Spindle-tree	84
Splachnum	399
Spleenwort	383, 385, 388
Sponge	489, 490
Spurge	182—185
laurel	145
olive	145
Spurrey	169, 177, 178
Squinancy-berries	85
wort	55
Star of Bethlehem	124, 125
of the Earth	53
wort	2, 319
Stitchwort	64, 166
Stonecrop	170—173
Strawberry	195
tree	155
Succory, fwines	302
wild	304
yellow	294
Sun-dew	117
flower	205
Sycamore	379

T	Page
TANSEY	311
wild	195
Tare	279, 280
Targionia	438
Teasel	49, 50
Thistle	305—309, 326
Thongs, fea	467
Thorn, black	186
buck	83
fea	368
fallow	368
white	188
Thorow-wax	97
Thrift	114
Throatwort	81, 82
Thyme	230
Toadflax, bastard	86
common	237, 238
Toothwort	233
Touch me not	332
Touchwood	497
Tormentil	198
Travellers Joy	217
Trefoil	282—289
marsh	71
Tremella	463, 469
Trichomanes	392
True-love	150
Truffles	502
Turneps	253
Tutfan	291
Twablade	338, 339

V	
V alerian	12, 13
greek	75
Venus's comb	108
Ver-	

INDEX NOMINUM ANGLICANORUM.

	Page		Page
Vervain	505	Williams, wild	174
Vetch	275—282	Willow	362—366
Violet	330—332	herb podded	141, 142
calathian	87	hooded	232
corn	82	purple	180
dame's	252	rose-bay	140
water	72	yellow	72
		Winter-green	139, 154
		Woad, wild	181
		Wood-binde	80
		green	272
		roof	55
		waxen	272
		Worm-feed, treacle	251
		wood	311, 312
		Wrack, grafs	343
		fea	466
		Y	
		Y Arrow	325
		Yellow-weed	181
		Yew-tree	374



F I N I S.









